(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 25 May 2001 (25.05.2001)

PCT

(10) International Publication Number WO 01/35726 A1

(51) International Patent Classification7: 5/00, C12N 5/14, 15/82

A01H 1/00.

(21) International Application Number: PCT/US00/31418

(22) International Filing Date: 14 November 2000 (14.11.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

 60/166,228
 17 November 1999 (17.11.1999)
 US

 60/197,899
 17 April 2000 (17.04.2000)
 US

 60/227,439
 22 August 2000 (22.08.2000)
 US

(71) Applicant (for all designated States except US): MENDEL BIOTECHNOLOGY, INC. [US/US]; 21375 Cabot Boulevard, Hayward, CA 94541 (US).

(71) Applicants and

(72) Inventors: HEARD, Jacqueline [US/US]; 810 Guildford Avenue, San Mateo, CA 94402 (US). RATCLIFFE, Oliver [GB/US]; 814 East 21st Street, Oakland, CA 94606 (US). CREELMAN, Robert [US/US]; 2801 Jennifer Drive, Castro Valley, CA 94546 (US). JIANG, Cai-Zhong [CN/US]; 34495 Heathrow Terrace, Fremont, CA 94555 (US). PINEDA, Omaira [CO/US]; 19563 Helen Place, Castro Valley, CA 94546 (US). **REUBER**, **Lynne** [US/US]; 2000 Walnut Avenue, Fremont, CA 94538 (US). **ADAM**, **Luc** [CA/US]; 25800 Industrial Boulevard, L403, Hayward, CA 94545 (US).

(74) Agent: GUERRERO, Karen; Mendel Biotechnology, Inc., 21375 Cabot Boulevard, Hayward, CA 94545 (US).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

1/35726 A

(54) Title: PATHOGEN TOLERANCE GENES

PATHOGEN TOLERANCE GENES

RELATED APPLICATION INFORMATION

The present invention claims the benefit from US Provisional Patent Application Serial Nos. 60/166,228 filed November 17, 1999 and 60/197,899 filed April 17, 2000 and "Plant Trait Modification III" filed August 22, 2000.

5

10

15

20

25

30

35

FIELD OF THE INVENTION

This invention relates to the field of plant biology. More particularly, the present invention pertains to compositions and methods for phenotypically modifying a plant.

BACKGROUND OF THE INVENTION

Transcription factors can modulate gene expression, either increasing or decreasing (inducing or repressing) the rate of transcription. This modulation results in differential levels of gene expression at various developmental stages, in different tissues and cell types, and in response to different exogenous (e.g., environmental) and endogenous stimuli throughout the life cycle of the organism.

Because transcription factors are key controlling elements of biological pathways, altering the expression levels of one or more transcription factors can change entire biological pathways in an organism. For example, manipulation of the levels of selected transcription factors may result in increased expression of economically useful proteins or metabolic chemicals in plants or to improve other agriculturally relevant characteristics. Conversely, blocked or reduced expression of a transcription factor may reduce biosynthesis of unwanted compounds or remove an undesirable trait. Therefore, manipulating transcription factor levels in a plant offers tremendous potential in agricultural biotechnology for modifying a plant's traits.

The present invention provides novel transcription factors useful for modifying a plant's phenotype in desirable ways, such as modifying a plant's pathogen tolerance.

SUMMARY OF THE INVENTION

In a first aspect, the invention relates to a recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-29, or a complementary nucleotide sequence thereof; (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a); (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID Nos. 2N-1, where N=1-29, or a complementary nucleotide sequence thereof; (d) a nucleotide sequence

comprising silent substitutions in a nucleotide sequence of (c); (e) a nucleotide sequence which hybridizes under stringent conditions over substantially the entire length of a nucleotide sequence of one or more of: (a), (b), (c), or (d); (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e); (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide having a biological activity that modifies a plant's pathogen tolerance; (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g); (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide sequence of any of (a)-(g); (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29; (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29; and (1) a nucleotide sequence which encodes a conserved domain of a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-29. The recombinant polynucleotide may further comprise a constitutive, inducible, or tissue-active promoter operably linked to the nucleotide sequence. The invention also relates to compositions comprising at least two of the above described polynucleotides.

10

15

20

25

30

35

In a second aspect, the invention is an isolated or recombinant polypeptide comprising a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotide described above.

In another aspect, the invention is a transgenic plant comprising one or more of the above described recombinant polynucleotides. In yet another aspect, the invention is a plant with altered expression levels of a polynucleotide described above or a plant with altered expression or activity levels of an above described polypeptide. Further, the invention may be a plant lacking a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-29.

The plant may be a soybean, wheat, corn, potato, cotton, rice, oilseed rape, sunflower, alfalfa, sugarcane, turf, banana, blackberry, blueberry, strawberry, raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits, or vegetable brassicas plant.

In a further aspect, the invention relates to a cloning or expression vector comprising the isolated or recombinant polynucleotide described above or cells comprising the cloning or expression vector.

In yet a further aspect, the invention relates to a composition produced by incubating a polynucleotide of the invention with a nuclease, a restriction enzyme, a polymerase; a polymerase and a primer; a cloning vector, or with a cell.

Furthermore, the invention relates to a method for producing a plant having improved pathogen tolerance. The method comprises altering the expression of an isolated or recombinant polynucleotide of the invention or altering the expression or activity of a polypeptide of the invention in a plant to produce a modified plant, and selecting the modified plant for modified pathogen tolerance.

5

10

15

20

25

30

35

In another aspect, the invention relates to a method of identifying a factor that is modulated by or interacts with a polypeptide encoded by a polynucleotide of the invention. The method comprises expressing a polypeptide encoded by the polynucleotide in a plant; and identifying at least one factor that is modulated by or interacts with the polypeptide. In one embodiment the method for identifying modulating or interacting factors is by detecting binding by the polypeptide to a promoter sequence, or by detecting interactions between an additional protein and the polypeptide in a yeast two hybrid system, or by detecting expression of a factor by hybridization to a microarray, subtractive hybridization or differential display.

In yet another aspect, the invention is a method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest. The method comprises placing the molecule in contact with a plant comprising the polynucleotide or polypeptide encoded by the polynucleotide of the invention and monitoring one or more of the expression level of the polynucleotide in the plant, the expression level of the polypeptide in the plant, and modulation of an activity of the polypeptide in the plant.

In yet another aspect, the invention relates to an integrated system, computer or computer readable medium comprising one or more character strings corresponding to a polynucleotide of the invention, or to a polypeptide encoded by the polynucleotide. The integrated system, computer or computer readable medium may comprise a link between one or more sequence strings to a modified plant pathogen tolerance phenotype.

In yet another aspect, the invention is a method for identifying a sequence similar or homologous to one or more polynucleotides of the invention, or one or more polypeptides encoded by the polynucleotides. The method comprises providing a sequence database; and, querying the sequence database with one or more target sequences corresponding to the one or more polynucleotides or to the one or more polypeptides to identify one or more sequence members of the database that display sequence similarity or homology to one or more of the one or more target sequences.

The method may further comprise of linking the one or more of the polynucleotides of the invention, or encoded polypeptides, to a modified plant pathogen tolerance phenotype.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a table of exemplary polynucleotide and polypeptide sequences of the invention. The table includes from left to right for each sequence: the SEQ ID No., the internal code reference number (GID), whether the sequence is a polynucleotide or polypeptide sequence, and identification of any conserved domains for the polypeptide sequences.

5

10

15

20

25

30

35

Figure 2 provides a table of exemplary sequences that are homologous to other sequences provided in the Sequence Listing and that are derived from *Arabidopsis thaliana*. The table includes from left to right: the SEQ ID No., the internal code reference number (GID), identification of the homologous sequence, whether the sequence is a polynucleotide or polypeptide sequence, and identification of any conserved domains for the polypeptide sequences.

Figure 3 provides a table of exemplary sequences that are homologous to the sequences provided in Figures 1 and 2 and that are derived from plants other than *Arabidopsis thaliana*. The table includes from left to right: the SEQ ID No., the internal code reference number (GID), the unique GenBank sequence ID No. (NID), the probability that the comparison was generated by chance (P-value), and the species from which the homologous gene was identified.

DETAILED DESCRIPTION

The present invention relates to polynucleotides and polypeptides, e.g. for modifying phenotypes of plants.

In particular, the polynucleotides or polypeptides are useful for modifying traits associated with a plant's pathogen tolerance when the expression levels of the polynucleotides or expression levels or activity levels of the polypeptides are altered. Specifically, the polynucleotides and polypeptides are useful for modifying traits associated with a plant's pathogen tolerance, such as alterations in cell wall composition, trichome number or structure, callose induction, phytoalexin induction, alterations in the cell death response, or the like. Transgenic plants employing the polynucleotides or polypeptides of the invention are more tolerant to biotrophic or necrotrophic pathogens such as fungi, bacteria, mollicutes, viruses, nematodes, parasitic higher plants or the like.

The polynucleotides of the invention encode plant transcription factors. The plant transcription factors are derived, e.g., from *Arabidopsis thaliana* and can belong, e.g., to one or more of the following transcription factor families: the AP2 (APETALA2) domain transcription factor family (Riechmann and Meyerowitz (1998) <u>J. Biol. Chem.</u> 379:633-646); the MYB transcription factor family (Martin and Paz-Ares (1997) <u>Trends Genet.</u> 13:67-73); the MADS domain transcription factor family (Riechmann and Meyerowitz (1997) <u>J. Biol.</u>

Chem. 378:1079-1101); the WRKY protein family (Ishiguro and Nakamura (1994) Mol. Gen. Genet. 244:563-571); the ankyrin-repeat protein family (Zhang et al. (1992) Plant Cell 4:1575-1588); the miscellaneous protein (MISC) family (Kim et al. (1997) Plant J. 11:1237-1251); the zinc finger protein (Z) family (Klug and Schwabe (1995) FASEB J. 9: 597-604); the homeobox (HB) protein family (Duboule (1994) Guidebook to the Homeobox Genes, Oxford University Press); the CAAT-element binding proteins (Forsburg and Guarente (1989) Genes Dev. 3:1166-1178); the squamosa promoter binding proteins (SPB) (Klein et al. (1996) Mol. Gen. Genet. 1996 250:7-16); the NAM protein family; the IAA/AUX proteins (Rouse et al. (1998) Science 279:1371-1373); the HLH/MYC protein family (Littlewood et al. (1994) Prot. Profile 1:639-709); the DNA-binding protein (DBP) family (Tucker et al. (1994) EMBO J. 13:2994-3002); the bZIP family of transcription factors (Foster et al. (1994) FASEB J. 8:192-200); the BPF-1 protein (Box P-binding factor) family (da Costa e Silva et al. (1993) Plant J. 4:125-135); and the golden protein (GLD) family (Hall et al. (1998) Plant Cell 10:925-936).

In addition to methods for modifying a plant phenotype by employing one or more polynucleotides and polypeptides of the invention described herein, the polynucleotides and polypeptides of the invention have a variety of additional uses. These uses include their use in the recombinant production (i.e, expression) of proteins; as regulators of plant gene expression, as diagnostic probes for the presence of complementary or partially complementary nucleic acids (including for detection of natural coding nucleic acids); as substrates for further reactions, e.g., mutation reactions, PCR reactions, or the like, of as substrates for cloning e.g., including digestion or ligation reactions, and for identifying exogenous or endogenous modulators of the transcription factors.

DEFINITIONS

A "polynucleotide" is a nucleic acid sequence comprising a plurality of polymerized nucleotide residues, e.g., at least about 15 consecutive polymerized nucleotide residues, optionally at least about 30 consecutive nucleotides, at least about 50 consecutive nucleotides. In many instances, a polynucleotide comprises a nucleotide sequence encoding a polypeptide (or protein) or a domain or fragment thereof. Additionally, the polynucleotide may comprise a promoter, an intron, an enhancer region, a polyadenylation site, a translation initiation site, 5' or 3' untranslated regions, a reporter gene, a selectable marker, or the like. The polynucleotide can be single stranded or double stranded DNA or RNA. The polynucleotide optionally comprises modified bases or a modified backbone. The polynucleotide can be, e.g., genomic DNA or RNA, a transcript (such as an mRNA), a cDNA, a PCR product, a cloned DNA, a synthetic DNA or RNA, or the like. The polynucleotide can comprise a sequence in either sense or antisense orientations.

A "recombinant polynucleotide" is a polynucleotide that is not in its native state, e.g., the polynucleotide comprises a nucleotide sequence not found in nature, or the polynucleotide is in a context other than that in which it is naturally found, e.g., separated from nucleotide sequences with which it typically is in proximity in nature, or adjacent (or contiguous with) nucleotide sequences with which it typically is not in proximity. For example, the sequence at issue can be cloned into a vector, or otherwise recombined with one or more additional nucleic acid.

5

10

15

20

25

30

35

An "isolated polynucleotide" is a polynucleotide whether naturally occurring or recombinant, that is present outside the cell in which it is typically found in nature, whether purified or not. Optionally, an isolated polynucleotide is subject to one or more enrichment or purification procedures, e.g., cell lysis, extraction, centrifugation, precipitation, or the like.

A "recombinant polypeptide" is a polypeptide produced by translation of a recombinant polypucleotide. An "isolated polypeptide," whether a naturally occurring or a recombinant polypeptide, is more enriched in (or out of) a cell than the polypeptide in its natural state in a wild type cell, e.g., more than about 5% enriched, more than about 10% enriched, or more than about 20%, or more than about 50%, or more, enriched, i.e., alternatively denoted: 105%, 110%, 120%, 150% or more, enriched relative to wild type standardized at 100%. Such an enrichment is not the result of a natural response of a wild type plant. Alternatively, or additionally, the isolated polypeptide is separated from other cellular components with which it is typically associated, e.g., by any of the various protein purification methods herein.

The term "transgenic plant" refers to a plant that contains genetic material, not found in a wild type plant of the same species, variety or cultivar. The genetic material may include a transgene, an insertional mutagenesis event (such as by transposon or T-DNA insertional mutagenesis), an activation tagging sequence, a mutated sequence, a homologous recombination event or a sequence modified by chimeraplasty. Typically, the foreign genetic material has been introduced into the plant by human manipulation.

A transgenic plant may contain an expression vector or cassette. The expression cassette typically comprises a polypeptide-encoding sequence operably linked (i.e., under regulatory control of) to appropriate inducible or constitutive regulatory sequences that allow for the expression of polypeptide. The expression cassette can be introduced into a plant by transformation or by breeding after transformation of a parent plant. A plant refers to a whole plant as well as to a plant part, such as seed, fruit, leaf, or root, plant tissue, plant cells or any other plant material, e.g., a plant explant, as well as to progeny thereof, and to *in vitro* systems that mimic biochemical or cellular components or processes in a cell.

The phrase "ectopically expression or altered expression" in reference to a polynucleotide indicates that the pattern of expression in, e.g., a transgenic plant or plant

tissue, is different from the expression pattern in a wild type plant or a reference plant of the same species. For example, the polynucleotide or polypeptide is expressed in a cell or tissue type other than a cell or tissue type in which the sequence is expressed in the wild type plant, or by expression at a time other than at the time the sequence is expressed in the wild type plant, or by a response to different inducible agents, such as hormones or environmental signals, or at different expression levels (either higher or lower) compared with those found in a wild type plant. The term also refers to altered expression patterns that are produced by lowering the levels of expression to below the detection level or completely abolishing expression. The resulting expression pattern can be transient or stable, constitutive or inducible. In reference to a polypeptide, the term "ectopic expression or altered expression" further may relate to altered activity levels resulting from the interactions of the polypeptides with exogenous or endogenous modulators or from interactions with factors or as a result of the chemical modification of the polypeptides.

The term "fragment" or "domain," with respect to a polypeptide, refers to a subsequence of the polypeptide. In some cases, the fragment or domain, is a subsequence of the polypeptide which performs at least one biological function of the intact polypeptide in substantially the same manner, or to a similar extent, as does the intact polypeptide. For example, a polypeptide fragment can comprise a recognizable structural motif or functional domain such as a DNA binding domain that binds to a DNA promoter region, an activation domain or a domain for protein-protein interactions. Fragments can vary in size from as few as 6 amino acids to the full length of the intact polypeptide, but are preferably at least about 30 amino acids in length and more preferably at least about 60 amino acids in length. In reference to a nucleotide sequence, "a fragment" refers to any subsequence of a polynucleotide, typically, of at least consecutive about 15 nucleotides, preferably at least about 30 nucleotides, more preferably at least about 50, of any of the sequences provided herein.

The term "trait" refers to a physiological, morphological, biochemical or physical characteristic of a plant or particular plant material or cell. In some instances, this characteristic is visible to the human eye, such as seed or plant size, or can be measured by available biochemical techniques, such as the protein, starch or oil content of seed or leaves or by the observation of the expression level of genes, e.g., by employing Northern analysis, RT-PCR, microarray gene expression assays or reporter gene expression systems, or by agricultural observations such as stress tolerance, yield or pathogen tolerance.

"Trait modification" refers to a detectable difference in a characteristic in a plant ectopically expressing a polynucleotide or polypeptide of the present invention relative to a plant not doing so, such as a wild type plant. In some cases, the trait modification can be evaluated quantitatively. For example, the trait modification can entail at least about a 2%

increase or decrease in an observed trait (difference), at least a 5% difference, at least about a 10% difference, at least about a 20% difference, at least about a 30%, at least about a 50%, at least about a 70%, or at least about a 100%, or an even greater difference. It is known that there can be a natural variation in the modified trait. Therefore, the trait modification observed entails a change of the normal distribution of the trait in the plants compared with the distribution observed in wild type plant.

5

10

15

20

25

30

35

Trait modifications of particular interest include those to seed (such as embryo or endosperm), fruit, root, flower, leaf, stem, shoot, seedling or the like, including: enhanced tolerance to environmental conditions including freezing, chilling, heat, drought, water saturation, radiation and ozone; improved tolerance to microbial, fungal or viral diseases; improved tolerance to pest infestations, including nematodes, mollicutes, parasitic higher plants or the like; decreased herbicide sensitivity; improved tolerance of heavy metals or enhanced ability to take up heavy metals; improved growth under poor photoconditions (e.g., low light and/or short day length), or changes in expression levels of genes of interest. Other phenotype that can be modified relate to the production of plant metabolites, such as variations in the production of taxol, tocopherol, tocotrienol, sterols, phytosterols, vitamins, wax monomers, anti-oxidants, amino acids, lignins, cellulose, tannins, prenyllipids (such as chlorophylls and carotenoids), glucosinolates, and terpenoids, enhanced or compositionally altered protein or oil production (especially in seeds), or modified sugar (insoluble or soluble) and/or starch composition. Physical plant characteristics that can be modified include cell development (such as the number of trichomes), fruit and seed size and number, yields of plant parts such as stems, leaves and roots, the stability of the seeds during storage, characteristics of the seed pod (e.g., susceptibility to shattering), root hair length and quantity, internode distances, or the quality of seed coat. Plant growth characteristics that can be modified include growth rate, germination rate of seeds, vigor of plants and seedlings, leaf and flower senescence, male sterility, apomixis, flowering time, flower abscission, rate of nitrogen uptake, biomass or transpiration characteristics, as well as plant architecture characteristics such as apical dominance, branching patterns, number of organs, organ identity, organ shape or size.

POLYPEPTIDES AND POLYNUCLEOTIDES OF THE INVENTION

The present invention provides, among other things, transcription factors (TFs), and transcription factor homologue polypeptides, and isolated or recombinant polynucleotides encoding the polypeptides. These polypeptides and polynucleotides may be employed to modify a plant's pathogen tolerance.

Exemplary polynucleotides encoding the polypeptides of the invention were identified in the *Arabidopsis thaliana* GenBank database using publicly available sequence

analysis programs and parameters. Sequences initially identified were then further characterized to identify sequences comprising specified sequence strings corresponding to sequence motifs present in families of known transcription factors. Polynucleotide sequences meeting such criteria were confirmed as transcription factors.

Additional polynucleotides of the invention were identified by screening Arabidopsis thaliana and/or other plant cDNA libraries with probes corresponding to known transcription factors under low stringency hybridization conditions. Additional sequences, including full length coding sequences were subsequently recovered by the rapid amplification of cDNA ends (RACE) procedure, using a commercially available kit according to the manufacturer's instructions. Where necessary, multiple rounds of RACE are performed to isolate 5' and 3' ends. The full length cDNA was then recovered by a routine end-to-end polymerase chain reaction (PCR) using primers specific to the isolated 5' and 3' ends. Exemplary sequences are provided in the Sequence Listing.

The polynucleotides of the invention were ectopically expressed in overexpressor or knockout plants and changes in the pathogen tolerance of the plants was observed. Therefore, the polynucleotides and polypeptides can be employed to improve the pathogen resistance of plants.

Making polynucleotides

5

10

15

20

25

30

35

The polynucleotides of the invention include sequences that encode transcription factors and transcription factor homologue polypeptides and sequences complementary thereto, as well as unique fragments of coding sequence, or sequence complementary thereto. Such polynucleotides can be, e.g., DNA or RNA, e.g., mRNA, cRNA, synthetic RNA, genomic DNA, cDNA synthetic DNA, oligonucleotides, etc. The polynucleotides are either double-stranded or single-stranded, and include either, or both sense (i.e., coding) sequences and antisense (i.e., non-coding, complementary) sequences. The polynucleotides include the coding sequence of a transcription factor, or transcription factor homologue polypeptide, in isolation, in combination with additional coding sequences (e.g., a purification tag, a localization signal, as a fusion-protein, as a pre-protein, or the like), in combination with non-coding sequences (e.g., introns or inteins, regulatory elements such as promoters, enhancers, terminators, and the like), and/or in a vector or host environment in which the polynucleotide encoding a transcription factor or transcription factor homologue polypeptide is an endogenous or exogenous gene.

A variety of methods exist for producing the polynucleotides of the invention. Procedures for identifying and isolating DNA clones are well known to those of skill in the art, and are described in, e.g., Berger and Kimmel, <u>Guide to Molecular Cloning Techniques</u>, <u>Methods in Enzymology</u> volume 152 Academic Press, Inc., San Diego, CA ("Berger");

Sambrook et al., Molecular Cloning - A Laboratory Manual (2nd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989 ("Sambrook") and Current Protocols in Molecular Biology, F.M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2000) ("Ausubel").

Alternatively, polynucleotides of the invention, can be produced by a variety of in vitro amplification methods adapted to the present invention by appropriate selection of specific or degenerate primers. Examples of protocols sufficient to direct persons of skill through in vitro amplification methods, including the polymerase chain reaction (PCR) the ligase chain reaction (LCR), Qbeta-repliçase amplification and other RNA polymerase mediated techniques (e.g., NASBA), e.g., for the production of the homologous nucleic acids of the invention are found in Berger, Sambrook, and Ausubel, as well as Mullis et al., (1987) PCR Protocols A Guide to Methods and Applications (Innis et al. eds) Academic Press Inc. San Diego, CA (1990) (Innis). Improved methods for cloning in vitro amplified nucleic acids are described in Wallace et al., U.S. Pat. No. 5,426,039. Improved methods for amplifying large nucleic acids by PCR are summarized in Cheng et al. (1994) Nature 369: 684-685 and the references cited therein, in which PCR amplicons of up to 40kb are generated. One of skill will appreciate that essentially any RNA can be converted into a double stranded DNA suitable for restriction digestion, PCR expansion and sequencing using reverse transcriptase and a polymerase. See, e.g., Ausubel, Sambrook and Berger, all supra.

Alternatively, polynucleotides and oligonucleotides of the invention can be assembled from fragments produced by solid-phase synthesis methods. Typically, fragments of up to approximately 100 bases are individually synthesized and then enzymatically or chemically ligated to produce a desired sequence, e.g., a polynucleotide encoding all or part of a transcription factor. For example, chemical synthesis using the phosphoramidite method is described, e.g., by Beaucage et al. (1981) Tetrahedron Letters 22:1859-69; and Matthes et al. (1984) EMBO J. 3:801-5. According to such methods, oligonucleotides are synthesized, purified, annealed to their complementary strand, ligated and then optionally cloned into suitable vectors. And if so desired, the polynucleotides and polypeptides of the invention can be custom ordered from any of a number of commercial suppliers.

HOMOLOGOUS SEQUENCES

5

10

15

20

25

30

35

Sequences homologous, i.e., that share significant sequence identity or similarity, to those provided in the Sequence Listing, derived from Arabidopsis thaliana or from other plants of choice are also an aspect of the invention. Homologous sequences can be derived from any plant including monocots and dicots and in particular agriculturally important plant species, including but not limited to, crops such as soybean, wheat, corn,

potato, cotton, rice, oilseed rape (including canola), sunflower, alfalfa, sugarcane and turf; or fruits and vegetables, such as banana, blackberry, blueberry, strawberry, and raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits (such as apple, peach, pear, cherry and plum) and vegetable brassicas (such as broccoli, cabbage, cauliflower, brussel sprouts and kohlrabi). Other crops, fruits and vegetables whose phenotype can be changed include barley, rye, millet, sorghum, currant, avocado, citrus fruits such as oranges, lemons, grapefruit and tangerines, artichoke, cherries, nuts such as the walnut and peanut, endive, leek, roots, such as arrowroot, beet, cassava, turnip, radish, yam, and sweet potato, and beans. The homologous sequences may also be derived from woody species, such pine, poplar and eucalyptus.

Transcription factors that are homologous to the listed sequences will typically share at least about 31% amino acid sequence identity. More closely related transcription factors can share at least about 50%, about 60%, about 65%, about 70%, about 75% or about 80% or about 90% or about 95% or about 98% or more sequence identity with the listed sequences. Factors that are most closely related to the listed sequences share, e.g., at least about 85%, about 90% or about 95% or more % sequence identity to the listed sequences. At the nucleotide level, the sequences will typically share at least about 40% nucleotide sequence identity, preferably at least about 50%, about 60%, about 70% or about 80% sequence identity, and more preferably about 85%, about 90%, about 95% or about 97% or more sequence identity to one or more of the listed sequences. The degeneracy of the genetic code enables major variations in the nucleotide sequence of a polynucleotide while maintaining the amino acid sequence of the encoded protein. Conserved domains within a transcription factor family may exhibit a higher degree of sequence homology, such as at least 65% sequence identity including conservative substitutions, and preferably at least 80% sequence identity.

Identifying Nucleic Acids by Hybridization

Polynucleotides homologous to the sequences illustrated in the Sequence Listing can be identified, e.g., by hybridization to each other under stringent or under highly stringent conditions. Single stranded polynucleotides hybridize when they associate based on a variety of well characterized physico-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. The stringency of a hybridization reflects the degree of sequence identity of the nucleic acids involved, such that the higher the stringency, the more similar are the two polynucleotide strands. Stringency is influenced by a variety of factors, including temperature, salt concentration and composition, organic and non-organic additives, solvents, etc. present in both the hybridization and wash solutions and incubations (and number), as described in more detail in the references cited above.

An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is about 5°C to 20°C lower than the thermal melting point (Tm) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Nucleic acid molecules that hybridize under stringent conditions will typically hybridize to a probe based on either the entire cDNA or selected portions, e.g., to a unique subsequence, of the cDNA under wash conditions of 0.2x SSC to 2.0 x SSC, 0.1% SDS at 50-65° C, for example 0.2 x SSC, 0.1% SDS at 65° C. For identification of less closely related homologues washes can be performed at a lower temperature, e.g., 50° C. In general, stringency is increased by raising the wash temperature and/or decreasing the concentration of SSC.

As another example, stringent conditions can be selected such that an oligonucleotide that is perfectly complementary to the coding oligonucleotide hybridizes to the coding oligonucleotide with at least about a 5-10x higher signal to noise ratio than the ratio for hybridization of the perfectly complementary oligonucleotide to a nucleic acid encoding a transcription factor known as of the filing date of the application. Conditions can be selected such that a higher signal to noise ratio is observed in the particular assay which is used, e.g., about 15x, 25x, 35x, 50x or more. Accordingly, the subject nucleic acid hybridizes to the unique coding oligonucleotide with at least a 2x higher signal to noise ratio as compared to hybridization of the coding oligonucleotide to a nucleic acid encoding known polypeptide. Again, higher signal to noise ratios can be selected, e.g., about 5x, 10x, 25x, 35x, 50x or more. The particular signal will depend on the label used in the relevant assay, e.g., a fluorescent label, a colorimetric label, a radio active label, or the like.

Alternatively, transcription factor homologue polypeptides can be obtained by screening an expression library using antibodies specific for one or more transcription factors. With the provision herein of the disclosed transcription factor, and transcription factor homologue nucleic acid sequences, the encoded polypeptide(s) can be expressed and purified in a heterologous expression system (e.g., *E. coli*) and used to raise antibodies (monoclonal or polyclonal) specific for the polypeptide(s) in question. Antibodies can also be raised against synthetic peptides derived from transcription factor, or transcription factor homologue, amino acid sequences. Methods of raising antibodies are well known in the art and are described in Harlow and Lane (1988) Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, New York. Such antibodies can then be used to screen an expression library produced from the plant from which it is desired to clone additional transcription

factor homologues, using the methods described above. The selected cDNAs can be confirmed by sequencing and enzymatic activity.

SEQUENCE VARIATIONS

5

10

It will readily be appreciated by those of skill in the art, that any of a variety of polynucleotide sequences are capable of encoding the transcription factors and transcription factor homologue polypeptides of the invention. Due to the degeneracy of the genetic code, many different polynucleotides can encode identical and/or substantially similar polypeptides in addition to those sequences illustrated in the Sequence Listing.

For example, Table 1 illustrates, e.g., that the codons AGC, AGT, TCA, TCC, TCG, and TCT all encode the same amino acid: serine. Accordingly, at each position in the sequence where there is a codon encoding serine, any of the above trinucleotide sequences can be used without altering the encoded polypeptide.

Table 1

Amino acids		•	Codon			•		
Alanine	Ala	A	GCA	GCC	GCG	GCU	·	
Cysteine	Cys	С	TGC	TGT				
Aspartic acid	Asp	D	GAC	GAT				
Glutamic acid	Glu	E	GAA	GAG			•	
Phenylalanine	Phe	F	TTC	TTT				
Glycine	Gly	G	GGA	GGC	GGG	GGT		
Histidine	His	H	CAC	CAT				
Isoleucine	Ile	I	ATA	ATC	ATT			
Lysine	Lys	K	AAA -	AAG				
Leucine	Leu	L	TTA	TTG	CTA	CTC	CTG	CTT
Methionine	Met	M	ATG					
Asparagine	Asn	N	AAC	AAT			•	
Proline	Pro	P	CCA	CCC	CCG	CCT		
Glutamine	Gln	Q	CAA	CAG				,
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGT
Serine	Ser	S	AGC	AGT	TCA	TCC	TCG	TCT
Threonine	Thr	T	ACA	ACC	ACG	ACT		
Valine	Val	V	GTA	GTC	GTG	GTT		
Tryptophan	Тгр	W	TGG					
Tyrosine	Tyr	Y	TAC	TAT				

15

20

Sequence alterations that do not change the amino acid sequence encoded by the polynucleotide are termed "silent" variations. With the exception of the codons ATG and TGG, encoding methionine and tryptophan, respectively, any of the possible codons for the same amino acid can be substituted by a variety of techniques, e.g., site-directed mutagenesis, available in the art. Accordingly, any and all such variations of a sequence selected from the above table are a feature of the invention.

In addition to silent variations, other conservative variations that alter one, or a few amino acids in the encoded polypeptide, can be made without altering the function of the polypeptide, these conservative variants are, likewise, a feature of the invention.

For example, substitutions, deletions and insertions introduced into the sequences provided in the Sequence Listing are also envisioned by the invention. Such sequence modifications can be engineered into a sequence by site-directed mutagenesis (Wu (ed.) Meth. Enzymol. (1993) vol. 217, Academic Press) or the other methods noted below. Amino acid substitutions are typically of single residues; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. In preferred embodiments, deletions or insertions are made in adjacent pairs, e.g., a deletion of two residues or insertion of two residues. Substitutions, deletions, insertions or any combination thereof can be combined to arrive at a sequence. The mutations that are made in the polynucleotide encoding the transcription factor should not place the sequence out of reading frame and should not create complementary regions that could produce secondary mRNA structure. Preferably, the polypeptide encoded by the DNA performs the desired function.

Conservative substitutions are those in which at least one residue in the amino acid sequence has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the Table 2 when it is desired to maintain the activity of the protein. Table 2 shows amino acids which can be substituted for an amino acid in a protein and which are typically regarded as conservative substitutions.

25

5

10

15

20

30

Table 2

Residue	Conservative Substitutions
Ala	Ser
Arg	Lys
Asn	Gln; His
Asp	Glu
Gln	Asn
Cys	Ser
Glu	Asp
Gly	Pro
His	Asn; Gln
Ile	Leu, Val
Leu	Ile; Val
Lys	Arg; Gln
Met	Leu; Ile
Phe	Met; Leu; Тут
Ser	Thr; Gly
Thr	Ser;Val
Ттр	Тут
Tyr	Trp; Phe
Val	Ile; Leu

Substitutions that are less conservative than those in Table 2 can be selected by picking residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in protein properties will be those in which (a) a hydrophilic residue, e.g., seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g., leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

5

FURTHER MODIFYING SEQUENCES OF THE INVENTION—MUTATION/ FORCED EVOLUTION

In addition to generating silent or conservative substitutions as noted, above, the present invention optionally includes methods of modifying the sequences of the Sequence Listing. In the methods, nucleic acid or protein modification methods are used to alter the given sequences to produce new sequences and/or to chemically or enzymatically modify given sequences to change the properties of the nucleic acids or proteins.

5

10

15

20

25

30

35

Thus, in one embodiment, given nucleic acid sequences are modified, e.g., according to standard mutagenesis or artificial evolution methods to produce modified sequences. For example, Ausubel, *supra*, provides additional details on mutagenesis methods. Artificial forced evolution methods are described, e.g., by Stemmer (1994) Nature 370:389-391, and Stemmer (1994) Proc. Natl. Acad. Sci. USA 91:10747-10751. Many other mutation and evolution methods are also available and expected to be within the skill of the practitioner.

Similarly, chemical or enzymatic alteration of expressed nucleic acids and polypeptides can be performed by standard methods. For example, sequence can be modified by addition of lipids, sugars, peptides, organic or inorganic compounds, by the inclusion of modified nucleotides or amino acids, or the like. For example, protein modification techniques are illustrated in Ausubel, *supra*. Further details on chemical and enzymatic modifications can be found herein. These modification methods can be used to modify any given sequence, or to modify any sequence produced by the various mutation and artificial evolution modification methods noted herein.

Accordingly, the invention provides for modification of any given nucleic acid by mutation, evolution, chemical or enzymatic modification, or other available methods, as well as for the products produced by practicing such methods, e.g., using the sequences herein as a starting substrate for the various modification approaches.

For example, optimized coding sequence containing codons preferred by a particular prokaryotic or eukaryotic host can be used e.g., to increase the rate of translation or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, as compared with transcripts produced using a non-optimized sequence. Translation stop codons can also be modified to reflect host preference. For example, preferred stop codons for *S. cerevisiae* and mammals are TAA and TGA, respectively. The preferred stop codon for monocotyledonous plants is TGA, whereas insects and *E. coli* prefer to use TAA as the stop codon.

The polynucleotide sequences of the present invention can also be engineered in order to alter a coding sequence for a variety of reasons, including but not limited to, alterations which modify the sequence to facilitate cloning, processing and/or expression of

the gene product. For example, alterations are optionally introduced using techniques which are well known in the art, e.g., site-directed mutagenesis, to insert new restriction sites, to alter glycosylation patterns, to change codon preference, to introduce splice sites, etc.

Furthermore, a fragment or domain derived from any of the polypeptides of the invention can be combined with domains derived from other transcription factors or synthetic domains to modify the biological activity of a transcription factor. For instance, a DNA binding domain derived from a transcription factor of the invention can be combined with the activation domain of another transcription factor or with a synthetic activation domain. A transcription activation domain assists in initiating transcription from a DNA binding site. Examples include the transcription activation region of VP16 or GAL4 (Moore et al. (1998) Proc. Natl. Acad. Sci. USA 95: 376-381; and Aoyama et al. (1995) Plant Cell 7:1773-1785), peptides derived from bacterial sequences (Ma and Ptashne (1987) Cell 51; 113-119) and synthetic peptides (Giniger and Ptashne, (1987) Nature 330:670-672).

EXPRESSION AND MODIFICATION OF POLYPEPTIDES

5

10

15

20

25

30

35

Typically, polynucleotide sequences of the invention are incorporated into recombinant DNA (or RNA) molecules that direct expression of polypeptides of the invention in appropriate host cells, transgenic plants, in vitro translation systems, or the like. Due to the inherent degeneracy of the genetic code, nucleic acid sequences which encode substantially the same or a functionally equivalent amino acid sequence can be substituted for any listed sequence to provide for cloning and expressing the relevant homologue.

Vectors, Promoters and Expression Systems

The present invention includes recombinant constructs comprising one or more of the nucleic acid sequences herein. The constructs typically comprise a vector, such as a plasmid, a cosmid, a phage, a virus (e.g., a plant virus), a bacterial artificial chromosome (BAC), a yeast artificial chromosome (YAC), or the like, into which a nucleic acid sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available.

General texts which describe molecular biological techniques useful herein, including the use and production of vectors, promoters and many other relevant topics, include Berger, Sambrook and Ausubel, *supra*. Any of the identified sequences can be incorporated into a cassette or vector, e.g., for expression in plants. A number of expression vectors suitable for stable transformation of plant cells or for the establishment of transgenic plants have been described including those described in Weissbach and Weissbach, (1989) Methods for Plant Molecular Biology, Academic Press, and Gelvin et al., (1990) Plant

Molecular Biology Manual, Kluwer Academic Publishers. Specific examples include those derived from a Ti plasmid of Agrobacterium tumefaciens, as well as those disclosed by Herrera-Estrella et al. (1983) Nature 303: 209, Bevan (1984) Nucl Acid Res. 12: 8711-8721, Klee (1985) Bio/Technology 3: 637-642, for dicotyledonous plants.

5

10

15

20

25

30

35

Alternatively, non-Ti vectors can be used to transfer the DNA into monocotyledonous plants and cells by using free DNA delivery techniques. Such methods can involve, for example, the use of liposomes, electroporation, microprojectile bombardment, silicon carbide whiskers, and viruses. By using these methods transgenic plants such as wheat, rice (Christou (1991) Bio/Technology 9: 957-962) and corn (Gordon-Kamm (1990) Plant Cell 2: 603-618) can be produced. An immature embryo can also be a good target tissue for monocots for direct DNA delivery techniques by using the particle gun (Weeks et al. (1993) Plant Physiol 102: 1077-1084; Vasil (1993) Bio/Technology 10: 667-674; Wan and Lemeaux (1994) Plant Physiol 104: 37-48, and for Agrobacterium-mediated DNA transfer (Ishida et al. (1996) Nature Biotech 14: 745-750).

Typically, plant transformation vectors include one or more cloned plant coding sequence (genomic or cDNA) under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant transformation vectors typically also contain a promoter (e.g., a regulatory region controlling inducible or constitutive, environmentally-or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, an RNA processing signal (such as intron splice sites), a transcription termination site, and/or a polyadenylation signal.

Examples of constitutive plant promoters which can be useful for expressing the TF sequence include: the cauliflower mosaic virus (CaMV) 35S promoter, which confers constitutive, high-level expression in most plant tissues (see, e.g., Odel et al. (1985) Nature 313:810); the nopaline synthase promoter (An et al. (1988) Plant Physiol 88:547); and the octopine synthase promoter (Fromm et al. (1989) Plant Cell 1: 977).

A variety of plant gene promoters that regulate gene expression in response to environmental, hormonal, chemical, developmental signals, and in a tissue-active manner can be used for expression of a TF sequence in plants. Choice of a promoter is based largely on the phenotype of interest and is determined by such factors as tissue (e.g., seed, fruit, root, pollen, vascular tissue, flower, carpel, etc.), inducibility (e.g., in response to wounding, heat, cold, drought, light, pathogens, etc.), timing, developmental stage, and the like. Numerous known promoters have been characterized and can favorable be employed to promote expression of a polynucleotide of the invention in a transgenic plant or cell of interest. For example, tissue specific promoters include: seed-specific promoters (such as the napin, phaseolin or DC3 promoter described in US Pat. No. 5,773,697), fruit-specific promoters that are active during fruit ripening (such as the dru 1 promoter (US Pat. No. 5,783,393), or the

2A11 promoter (US Pat. No. 4,943,674) and the tomato polygalacturonase promoter (Bird et al. (1988) Plant Mol Biol 11:651), root-specific promoters, such as those disclosed in US Patent Nos. 5,618,988, 5,837,848 and 5,905,186, pollen-active promoters such as PTA29, PTA26 and PTA13 (US Pat. No. 5,792,929), promoters active in vascular tissue (Ringli and Keller (1998) Plant Mol Biol 37:977-988), flower-specific (Kaiser et al, (1995) Plant Mol Biol 28:231-243), pollen (Baerson et al. (1994) Plant Mol Biol 26:1947-1959), carpels (Ohl et al. (1990) Plant Cell 2:837-848), pollen and ovules (Baerson et al. (1993) Plant Mol Biol 22:255-267), auxin-inducible promoters (such as that described in van der Kop et al. (1999) Plant Mol Biol 39:979-990 or Baumann et al. (1999) Plant Cell 11:323-334), cytokinininducible promoter (Guevara-Garcia (1998) Plant Mol Biol 38:743-753), promoters responsive to gibberellin (Shi et al. (1998) Plant Mol Biol 38:1053-1060, Willmott et al. (1998) 38:817-825) and the like. Additional promoters are those that elicit expression in response to heat (Ainley et al. (1993) Plant Mol Biol 22: 13-23), light (e.g., the pea rbcS-3A promoter, Kuhlemeier et al. (1989) Plant Cell 1:471, and the maize rbcS promoter, Schaffner and Sheen (1991) Plant Cell 3: 997); wounding (e.g., wunI, Siebertz et al. (1989) Plant Cell 1: 961); pathogens (such as the PR-1 promoter described in Buchel et al. (1999) Plant Mol. Biol. 40:387-396, and the PDF1.2 promoter described in Manners et al. (1998) Plant Mol. Biol. 38:1071-80), and chemicals such as methyl jasmonate or salicylic acid (Gatz et al. (1997) Plant Mol Biol 48: 89-108). In addition, the timing of the expression can be controlled by using promoters such as those acting at senescence (An and Amazon (1995) Science 270: 1986-1988); or late seed development (Odell et al. (1994) Plant Physiol 106:447-458).

Plant expression vectors can also include RNA processing signals that can be positioned within, upstream or downstream of the coding sequence. In addition, the expression vectors can include additional regulatory sequences from the 3'-untranslated region of plant genes, e.g., a 3' terminator region to increase mRNA stability of the mRNA, such as the PI-II terminator region of potato or the octopine or nopaline synthase 3' terminator regions.

Additional Expression Elements

5

10

15

20

25

30

35

Specific initiation signals can aid in efficient translation of coding sequences. These signals can include, e.g., the ATG initiation codon and adjacent sequences. In cases where a coding sequence, its initiation codon and upstream sequences are inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only coding sequence (e.g., a mature protein coding sequence), or a portion thereof, is inserted, exogenous transcriptional control signals including the ATG initiation codon can be separately provided. The initiation codon is provided in the correct reading frame to facilitate transcription. Exogenous transcriptional elements and initiation

codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use.

Expression Hosts

5

10

15

20

25

30

35

The present invention also relates to host cells which are transduced with vectors of the invention, and the production of polypeptides of the invention (including fragments thereof) by recombinant techniques. Host cells are genetically engineered (i.e, nucleic acids are introduced, e.g., transduced, transformed or transfected) with the vectors of this invention, which may be, for example, a cloning vector or an expression vector comprising the relevant nucleic acids herein. The vector is optionally a plasmid, a viral particle, a phage, a naked nucleic acids, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the relevant gene. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to those skilled in the art and in the references cited herein, including, Sambrook and Ausubel.

The host cell can be a eukaryotic cell, such as a yeast cell, or a plant cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Plant protoplasts are also suitable for some applications. For example, the DNA fragments are introduced into plant tissues, cultured plant cells or plant protoplasts by standard methods including electroporation (Fromm et al., (1985) Proc. Natl. Acad. Sci. USA 82, 5824, infection by viral vectors such as cauliflower mosaic virus (CaMV) (Hohn et al., (1982) Molecular Biology of Plant Tumors, (Academic Press, New York) pp. 549-560; US 4,407,956), high velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface (Klein et al., (1987) Nature 327, 70-73), use of pollen as vector (WO 85/01856), or use of Agrobacterium tumefaciens or A. rhizogenes carrying a T-DNA plasmid in which DNA fragments are cloned. The T-DNA plasmid is transmitted to plant cells upon infection by Agrobacterium tumefaciens, and a portion is stably integrated into the plant genome (Horsch et al. (1984) Science 233:496-498; Fraley et al. (1983) Proc. Natl. Acad. Sci. USA 80, 4803).

The cell can include a nucleic acid of the invention which encodes a polypeptide, wherein the cells expresses a polypeptide of the invention. The cell can also include vector sequences, or the like. Furthermore, cells and transgenic plants which include any polypeptide or nucleic acid above or throughout this specification, e.g., produced by transduction of a vector of the invention, are an additional feature of the invention.

For long-term, high-yield production of recombinant proteins, stable expression can be used. Host cells transformed with a nucleotide sequence encoding a polypeptide of the invention are optionally cultured under conditions suitable for the

expression and recovery of the encoded protein from cell culture. The protein or fragment thereof produced by a recombinant cell may be secreted, membrane-bound, or contained intracellularly, depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides encoding mature proteins of the invention can be designed with signal sequences which direct secretion of the mature polypeptides through a prokaryotic or eukaryotic cell membrane.

Modified Amino Acids

5

10

15

20

25

30

35

Polypeptides of the invention may contain one or more modified amino acids. The presence of modified amino acids may be advantageous in, for example, increasing polypeptide half-life, reducing polypeptide antigenicity or toxicity, increasing polypeptide storage stability, or the like. Amino acid(s) are modified, for example, co-translationally or post-translationally during recombinant production or modified by synthetic or chemical means.

Non-limiting examples of a modified amino acid include incorporation or other use of acetylated amino acids, glycosylated amino acids, sulfated amino acids, prenylated (e.g., farnesylated, geranylgeranylated) amino acids, PEG modified (e.g., "PEGylated") amino acids, biotinylated amino acids, carboxylated amino acids, phosphorylated amino acids, etc. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature.

IDENTIFICATION OF ADDITIONAL FACTORS

A transcription factor provided by the present invention can also be used to identify additional endogenous or exogenous molecules that can affect a phentoype or trait of interest. On the one hand, such molecules include organic (small or large molecules) and/or inorganic compounds that affect expression of (i.e., regulate) a particular transcription factor. Alternatively, such molecules include endogenous molecules that are acted upon either at a transcriptional level by a transcription factor of the invention to modify a phenotype as desired. For example, the transcription factors can be employed to identify one or more downstream gene with which is subject to a regulatory effect of the transcription factor. In one approach, a transcription factor or transcription factor homologue of the invention is expressed in a host cell, e.g, a transgenic plant cell, tissue or explant, and expression products, either RNA or protein, of likely or random targets are monitored, e.g., by hybridization to a microarray of nucleic acid probes corresponding to genes expressed in a tissue or cell type of interest, by two-dimensional gel electrophoresis of protein products, or by any other method known in the art for assessing expression of gene products at the level of RNA or protein. Alternatively, a transcription factor of the invention can be used to identify promoter sequences (i.e., binding sites) involved in the regulation of a downstream target. After

identifying a promoter sequence, interactions between the transcription factor and the promoter sequence can be modified by changing specific nucleotides in the promoter sequence or specific amino acids in the transcription factor that interact with the promoter sequence to alter a plant trait. Typically, transcription factor DNA binding sites are identified by gel shift assays. After identifying the promoter regions, the promoter region sequences can be employed in double-stranded DNA arrays to identify molecules that affect the interactions of the transcription factors with their promoters (Bulyk et al. (1999) Nature Biotechnology 17:573-577).

The identified transcription factors are also useful to identify proteins that modify the activity of the transcription factor. Such modification can occur by covalent modification, such as by phosphorylation, or by protein-protein (homo or-heteropolymer) interactions. Any method suitable for detecting protein-protein interactions can be employed. Among the methods that can be employed are co-immunoprecipitation, cross-linking and co-purification through gradients or chromatographic columns, and the two-hybrid yeast system.

The two-hybrid system detects protein interactions in vivo and is described in Chien, et al., (1991), Proc. Natl. Acad. Sci. USA 88, 9578-9582 and is commercially available from Clontech (Palo Alto, Calif.). In such a system, plasmids are constructed that encode two hybrid proteins: one consists of the DNA-binding domain of a transcription activator protein fused to the TF polypeptide and the other consists of the transcription activator protein's activation domain fused to an unknown protein that is encoded by a cDNA that has been recombined into the plasmid as part of a cDNA library. The DNA-binding domain fusion plasmid and the cDNA library are transformed into a strain of the yeast Saccharomyces cerevisiae that contains a reporter gene (e.g., lacZ) whose regulatory region contains the transcription activator's binding site. Either hybrid protein alone cannot activate transcription of the reporter gene. Interaction of the two hybrid proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is detected by an assay for the reporter gene product. Then, the library plasmids responsible for reporter gene expression are isolated and sequenced to identify the proteins encoded by the library plasmids. After identifying proteins that interact with the transcription factors, assays for compounds that interfere with the TF protein-protein interactions can be preformed.

IDENTIFICATION OF MODULATORS

5

10

15

20

25

30

35

In addition to the intracellular molecules described above, extracellular molecules that alter activity or expression of a transcription factor, either directly or indirectly, can be identified. For example, the methods can entail first placing a candidate molecule in contact with a plant or plant cell. The molecule can be introduced by topical administration, such as spraying or soaking of a plant, and then the molecule's effect on the

expression or activity of the TF polypeptide or the expression of the polynucleotide monitored. Changes in the expression of the TF polypeptide can be monitored by use of polyclonal or monoclonal antibodies, gel electrophoresis or the like. Changes in the expression of the corresponding polynucleotide sequence can be detected by use of microarrays, Northerns, quantitative PCR, or any other technique for monitoring changes in mRNA expression. These techniques are exemplified in Ausubel et al. (eds) <u>Current Protocols in Molecular Biology</u>, John Wiley & Sons (1998). Such changes in the expression levels can be correlated with modified plant traits and thus identified molecules can be useful for soaking or spraying on fruit, vegetable and grain crops to modify traits in plants.

Essentially any available composition can be tested for modulatory activity of expression or activity of any nucleic acid or polypeptide herein. Thus, available libraries of compounds such as chemicals, polypeptides, nucleic acids and the like can be tested for modulatory activity. Often, potential modulator compounds can be dissolved in aqueous or organic (e.g., DMSO-based) solutions for easy delivery to the cell or plant of interest in which the activity of the modulator is to be tested. Optionally, the assays are designed to screen large modulator composition libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (e.g., in microtiter formats on microtiter plates in robotic assays).

In one embodiment, high throughput screening methods involve providing a combinatorial library containing a large number of potential compounds (potential modulator compounds). Such "combinatorial chemical libraries" are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as target compounds.

A combinatorial chemical library can be, e.g., a collection of diverse chemical compounds generated by chemical synthesis or biological synthesis. For example, a combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (e.g., in one example, amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound of a set length). Exemplary libraries include peptide libraries, nucleic acid libraries, antibody libraries (see, e.g., Vaughn et al. (1996) Nature Biotechnology, 14(3):309-314 and PCT/US96/10287), carbohydrate libraries (see, e.g., Liang et al. Science (1996) 274:1520-1522 and U.S. Patent 5,593,853), peptide nucleic acid libraries (see, e.g., U.S. Patent 5,539,083), and small organic molecule libraries (see, e.g., benzodiazepines, Baum C&EN Jan 18, page 33 (1993); isoprenoids, U.S. Patent 5,569,588; thiazolidinones and metathiazanones, U.S. Patent 5,549,974; pyrrolidines, U.S. Patents 5,525,735 and 5,519,134; morpholino compounds, U.S. Patent 5,506,337) and the like.

Preparation and screening of combinatorial or other libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (see, e.g., U.S. Patent 5,010,175, Furka, Int. J. Pept. Prot. Res. 37:487-493 (1991) and Houghton et al. Nature 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used.

5

10

15

20

25

30

35

In addition, as noted, compound screening equipment for high-throughput screening is generally available, e.g., using any of a number of well known robotic systems that have also been developed for solution phase chemistries useful in assay systems. These systems include automated workstations including an automated synthesis apparatus and robotic systems utilizing robotic arms. Any of the above devices are suitable for use with the present invention, e.g., for high-throughput screening of potential modulators. The nature and implementation of modifications to these devices (if any) so that they can operate as discussed herein will be apparent to persons skilled in the relevant art.

Indeed, entire high throughput screening systems are commercially available. These systems typically automate entire procedures including all sample and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. Similarly, microfluidic implementations of screening are also commercially available.

The manufacturers of such systems provide detailed protocols the various high throughput. Thus, for example, Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like. The integrated systems herein, in addition to providing for sequence alignment and, optionally, synthesis of relevant nucleic acids, can include such screening apparatus to identify modulators that have an effect on one or more polynucleotides or polypeptides according to the present invention.

In some assays it is desirable to have positive controls to ensure that the components of the assays are working properly. At least two types of positive controls are appropriate. That is, known transcriptional activators or inhibitors can be incubated with cells/plants/ etc. in one sample of the assay, and the resulting increase/decrease in transcription can be detected by measuring the resulting increase in RNA/ protein expression, etc., according to the methods herein. It will be appreciated that modulators can also be combined with transcriptional activators or inhibitors to find modulators which inhibit transcriptional activation or transcriptional repression. Either expression of the nucleic acids and proteins herein or any additional nucleic acids or proteins activated by the nucleic acids or proteins herein, or both, can be monitored.

In an embodiment, the invention provides a method for identifying compositions that modulate the activity or expression of a polynucleotide or polypeptide of the invention. For example, a test compound, whether a small or large molecule, is placed in contact with a cell, plant (or plant tissue or explant), or composition comprising the polynucleotide or polypeptide of interest and a resulting effect on the cell, plant, (or tissue or explant) or composition is evaluated by monitoring, either directly or indirectly, one or more of: expression level of the polynucleotide or polypeptide, activity (or modulation of the activity) of the polynucleotide or polypeptide. In some cases, an alteration in a plant phenotype can be detected following contact of a plant (or plant cell, or tissue or explant) with the putative modulator, e.g., by modulation of expression or activity of a polynucleotide or polypeptide of the invention.

SUBSEQUENCES

5

10

15

20

25

30

35

Also contemplated are uses of polynucleotides, also referred to herein as oligonucleotides, typically having at least 12 bases, preferably at least 15, more preferably at least 20, 30, or 50 bases, which hybridize under at least highly stringent (or ultra-high stringent or ultra-ultra- high stringent conditions) conditions to a polynucleotide sequence described above. The polynucleotides may be used as probes, primers, sense and antisense agents, and the like, according to methods as noted *supra*.

Subsequences of the polynucleotides of the invention, including polynucleotide fragments and oligonucleotides are useful as nucleic acid probes and primers. An oligonucleotide suitable for use as a probe or primer is at least about 15 nucleotides in length, more often at least about 18 nucleotides, often at least about 21 nucleotides, frequently at least about 30 nucleotides, or about 40 nucleotides, or more in length. A nucleic acid probe is useful in hybridization protocols, e.g., to identify additional polypeptide homologues of the invention, including protocols for microarray experiments. Primers can be annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, and then extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other nucleic-acid amplification methods. See Sambrook and Ausubel, *supra*.

In addition, the invention includes an isolated or recombinant polypeptide including a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotides of the invention. For example, such polypeptides, or domains or fragments thereof, can be used as immunogens, e.g., to produce antibodies specific for the polypeptide sequence, or as probes for detecting a sequence of interest. A

subsequence can range in size from about 15 amino acids in length up to and including the full length of the polypeptide.

PRODUCTION OF TRANSGENIC PLANTS

5

10

15

20

25

30

35

Modification of Traits

The polynucleotides of the invention are favorably employed to produce transgenic plants with various traits, or characteristics, that have been modified in a desirable manner, e.g., to improve the pathogen resistance of a plant. For example, alteration of expression levels or patterns (e.g., spatial or temporal expression patterns) of one or more of the transcription factors (or transcription factor homologues) of the invention, as compared with the levels of the same protein found in a wild type plant, can be used to modify a plant's traits. An illustrative example of trait modification, improved pathogen tolerance, by altering expression levels of a particular transcription factor is described further in the Examples and the Sequence Listing.

Antisense and Cosuppression Approaches

In addition to expression of the nucleic acids of the invention as gene replacement or plant phenotype modification nucleic acids, the nucleic acids are also useful for sense and anti-sense suppression of expression, e.g., to down-regulate expression of a nucleic acid of the invention, e.g., as a further mechanism for modulating plant phenotype. That is, the nucleic acids of the invention, or subsequences or anti-sense sequences thereof, can be used to block expression of naturally occurring homologous nucleic acids. A variety of sense and anti-sense technologies are known in the art, e.g., as set forth in Lichtenstein and Nellen (1997) Antisense Technology: A Practical Approach IRL Press at Oxford University, Oxford, England. In general, sense or anti-sense sequences are introduced into a cell, where they are optionally amplified, e.g., by transcription. Such sequences include both simple oligonucleotide sequences and catalytic sequences such as ribozymes.

For example, a reduction or elimination of expression (i.e., a "knock-out") of a transcription factor or transcription factor homologue polypeptide in a transgenic plant, e.g., to modify a plant trait, can be obtained by introducing an antisense construct corresponding to the polypeptide of interest as a cDNA. For antisense suppression, the transcription factor or homologue cDNA is arranged in reverse orientation (with respect to the coding sequence) relative to the promoter sequence in the expression vector. The introduced sequence need not be the full length cDNA or gene, and need not be identical to the cDNA or gene found in the plant type to be transformed. Typically, the antisense sequence need only be capable of hybridizing to the target gene or RNA of interest. Thus, where the introduced sequence is of shorter length, a higher degree of homology to the endogenous transcription factor sequence will be needed for effective antisense suppression. While antisense sequences of various

lengths can be utilized, preferably, the introduced antisense sequence in the vector will be at least 30 nucleotides in length, and improved antisense suppression will typically be observed as the length of the antisense sequence increases. Preferably, the length of the antisense sequence in the vector will be greater than 100 nucleotides. Transcription of an antisense construct as described results in the production of RNA molecules that are the reverse complement of mRNA molecules transcribed from the endogenous transcription factor gene in the plant cell.

5

10

15

20

25

30

35

Suppression of endogenous transcription factor gene expression can also be achieved using a ribozyme. Ribozymes are RNA molecules that possess highly specific endoribonuclease activity. The production and use of ribozymes are disclosed in U.S. Patent No. 4,987,071 and U.S. Patent No. 5,543,508. Synthetic ribozyme sequences including antisense RNAs can be used to confer RNA cleaving activity on the antisense RNA, such that endogenous mRNA molecules that hybridize to the antisense RNA are cleaved, which in turn leads to an enhanced antisense inhibition of endogenous gene expression.

Vectors in which RNA encoded by a transcription factor or transcription factor homologue cDNA is over-expressed can also be used to obtain co-suppression of a corresponding endogenous gene, e.g., in the manner described in U.S. Patent No. 5,231,020 to Jorgensen. Such co-suppression (also termed sense suppression) does not require that the entire transcription factor cDNA be introduced into the plant cells, nor does it require that the introduced sequence be exactly identical to the endogenous transcription factor gene of interest. However, as with antisense suppression, the suppressive efficiency will be enhanced as specificity of hybridization is increased, e.g., as the introduced sequence is lengthened, and/or as the sequence similarity between the introduced sequence and the endogenous transcription factor gene is increased.

Vectors expressing an untranslatable form of the transcription factor mRNA, e.g., sequences comprising one or more stop codon, or nonsense mutation) can also be used to suppress expression of an endogenous transcription factor, thereby reducing or eliminating it's activity and modifying one or more traits. Methods for producing such constructs are described in U.S. Patent No. 5,583,021. Preferably, such constructs are made by introducing a premature stop codon into the transcription factor gene. Alternatively, a plant trait can be modified by gene silencing using double-strand RNA (Sharp (1999) Genes and Development 13: 139-141).

Another method for abolishing the expression of a gene is by insertion mutagenesis using the T-DNA of Agrobacterium tumefaciens. After generating the insertion mutants, the mutants can be screened to identify those containing the insertion in a transcription factor or transcription factor homologue gene. Plants containing a single

transgene insertion event at the desired gene can be crossed to generate homozygous plants for the mutation (Koncz et al. (1992) Methods in Arabidopsis Research, World Scientific).

Alternatively, a plant phenotype can be altered by eliminating an endogenous gene, such as a transcription factor or transcription factor homologue, e.g., by homologous recombination (Kempin et al. (1997) Nature 389:802).

5

10

15

20

25

30

35

A plant trait can also be modified by using the cre-lox system (for example, as described in US Paent No. 5,658,772). A plant genome can be modified to include first and second lox sites that are then contacted with a Cre recombinase. If the lox sites are in the same orientation, the intervening DNA sequence between the two sites is excised. If the lox sites are in the opposite orientation, the intervening sequence is inverted.

The polynucleotides and polypeptides of this invention can also be expressed in a plant in the absence of an expression cassette by manipulating the activity or expression level of the endogenous gene by other means. For example, by ectopically expressing a gene by T-DNA activation tagging (Ichikawa et al. (1997) Nature 390 698-701; Kakimoto et al. (1996) Science 274: 982-985). This method entails transforming a plant with a gene tag containing multiple transcriptional enhancers and once the tag has inserted into the genome, expression of a flanking gene coding sequence becomes deregulated. In another example, the transcriptional machinery in a plant can be modified so as to increase transcription levels of a polynucleotide of the invention (See, e.g., PCT Publications WO 96/06166 and WO 98/53057 which describe the modification of the DNA binding specificity of zinc finger proteins by changing particular amino acids in the DNA binding motif).

The transgenic plant can also include the machinery necessary for expressing or altering the activity of a polypeptide encoded by an endogenous gene, for example by altering the phosphorylation state of the polypeptide to maintain it in an activated state.

Transgenic plants (or plant cells, or plant explants, or plant tissues) incorporating the polynucleotides of the invention and/or expressing the polypeptides of the invention can be produced by a variety of well established techniques as described above. Following construction of a vector, most typically an expression cassette, including a polynucleotide, e.g., encoding a transcription factor or transcription factor homologue, of the invention, standard techniques can be used to introduce the polynucleotide into a plant, a plant cell, a plant explant or a plant tissue of interest. Optionally, the plant cell, explant or tissue can be regenerated to produce a transgenic plant.

The plant can be any higher plant, including gymnosperms, monocotyledonous and dicotyledonous plants. Suitable protocols are available for *Leguminosae* (alfalfa, soybean, clover, etc.), *Umbelliferae* (carrot, celery, parsnip), *Cruciferae* (cabbage, radish, rapeseed, broccoli, etc.), *Curcurbitaceae* (melons and cucumber), *Gramineae* (wheat, corn, rice, barley, millet, etc.), *Solanaceae* (potato, tomato, tobacco,

peppers, etc.), and various other crops. See protocols described in Ammirato et al. (1984) Handbook of Plant Cell Culture -Crop Species. Macmillan Publ. Co. Shimamoto et al. (1989) Nature 338:274-276; Fromm et al. (1990) Bio/Technology 8:833-839; and Vasil et al. (1990) Bio/Technology 8:429-434.

5

10

15

20

25

30

35

Transformation and regeneration of both monocotyledonous and dicotyledonous plant cells is now routine, and the selection of the most appropriate transformation technique will be determined by the practitioner. The choice of method will vary with the type of plant to be transformed; those skilled in the art will recognize the suitability of particular methods for given plant types. Suitable methods can include, but are not limited to: electroporation of plant protoplasts; liposome-mediated transformation; polyethylene glycol (PEG) mediated transformation; transformation using viruses; microinjection of plant cells; micro-projectile bombardment of plant cells; vacuum infiltration; and Agrobacterium tumeficiens mediated transformation. Transformation means introducing a nucleotide sequence in a plant in a manner to cause stable or transient expression of the sequence.

Successful examples of the modification of plant characteristics by transformation with cloned sequences which serve to illustrate the current knowledge in this field of technology, and which are herein incorporated by reference, include: U.S. Patent Nos. 5,571,706; 5,677,175; 5,510,471; 5,750,386; 5,597,945; 5,589,615; 5,750,871; 5,268,526; 5,780,708; 5,538,880; 5,773,269; 5,736,369 and 5,610,042.

Following transformation, plants are preferably selected using a dominant selectable marker incorporated into the transformation vector. Typically, such a marker will confer antibiotic or herbicide resistance on the transformed plants, and selection of transformants can be accomplished by exposing the plants to appropriate concentrations of the antibiotic or herbicide.

After transformed plants are selected and grown to maturity, those plants showing a modified trait are identified. The modified trait can be any of those traits described above. Additionally, to confirm that the modified trait is due to changes in expression levels or activity of the polypeptide or polynucleotide of the invention can be determined by analyzing mRNA expression using Northern blots, RT-PCR or microarrays, or protein expression using immunoblots or Western blots or gel shift assays.

INTEGRATED SYSTEMS—SEQUENCE IDENTITY

Additionally, the present invention may be an integrated system, computer or computer readable medium that comprises an instruction set for determining the identity of one or more sequences in a database. In addition, the instruction set can be used to generate or identify sequences that meet any specified criteria. Furthermore, the instruction set may

be used to associate or link certain functional benefits, such improved pathogen tolerance, with one or more identified sequence.

For example, the instruction set can include, e.g., a sequence comparison or other alignment program, e.g., an available program such as, for example, the Wisconsin Package Version 10.0, such as BLAST, FASTA, PILEUP, FINDPATTERNS or the like (GCG, Madision, WI). Public sequence databases such as GenBank, EMBL, Swiss-Prot and PIR or private sequence databases such as PhytoSeq (Incyte Pharmaceuticals, Palo Alto, CA) can be searched.

5

10

15

20

25

30

35

Alignment of sequences for comparison can be conducted by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math. 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443, by the search for similarity method of Pearson and Lipman (1988) Proc. Natl. Acad. Sci. U.S.A. 85: 2444, by computerized implementations of these algorithms. After alignment, sequence comparisons between two (or more) polynucleotides or polypeptides are typically performed by comparing sequences of the two sequences over a comparison window to identify and compare local regions of sequence similarity. The comparison window can be a segment of at least about 20 contiguous positions, usually about 50 to about 200, more usually about 100 to about 150 contiguous positions. A description of the method is provided in Ausubel et al., supra.

A variety of methods of determining sequence relationships can be used, including manual alignment and computer assisted sequence alignment and analysis. This later approach is a preferred approach in the present invention, due to the increased throughput afforded by computer assisted methods. As noted above, a variety of computer programs for performing sequence alignment are available, or can be produced by one of skill.

One example algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al. J. Mol. Biol 215:403-410 (1990). Software for performing BLAST analyses is publicly available, e.g., through the National Center for Biotechnology Information (http://www.ncbi.nlm.nih.gov/). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters

M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff (1989) Proc. Natl. Acad. Sci. USA 89:10915).

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-5787). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence (and, therefore, in this context, homologous) if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, or less than about 0.01, and or even less than about 0.001. An additional example of a useful sequence alignment algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. The program can align, e.g., up to 300 sequences of a maximum length of 5,000 letters.

The integrated system, or computer typically includes a user input interface allowing a user to selectively view one or more sequence records corresponding to the one or more character strings, as well as an instruction set which aligns the one or more character strings with each other or with an additional character string to identify one or more region of sequence similarity. The system may include a link of one or more character strings with a particular phenotype or gene function. Typically, the system includes a user readable output element which displays an alignment produced by the alignment instruction set.

The methods of this invention can be implemented in a localized or distributed computing environment. In a distributed environment, the methods may implemented on a single computer comprising multiple processors or on a multiplicity of computers. The computers can be linked, e.g. through a common bus, but more preferably the computer(s) are nodes on a network. The network can be a generalized or a dedicated local or

wide-area network and, in certain preferred embodiments, the computers may be components of an intra-net or an internet.

Thus, the invention provides methods for identifying a sequence similar or homologous to one or more polynucleotides as noted herein, or one or more target polypeptides encoded by the polynucleotides, or otherwise noted herein and may include linking or associating a given plant phenotype or gene function with a sequence. In the methods, a sequence database is provided (locally or across an inter or intra net) and a query is made against the sequence database using the relevant sequences herein and associated plant phenotypes or gene functions.

5

10

15

25

30

35

Any sequence herein can be entered into the database, before or after querying the database. This provides for both expansion of the database and, if done before the querying step, for insertion of control sequences into the database. The control sequences can be detected by the query to ensure the general integrity of both the database and the query. As noted, the query can be performed using a web browser based interface. For example, the database can be a centralized public database such as those noted herein, and the querying can be done from a remote terminal or computer across an internet or intranet.

EXAMPLES

The following examples are intended to illustrate but not limit the present invention.

20 EXAMPLE I. FULL LENGTH GENE IDENTIFICATION AND CLONING

Putative transcription factor sequences (genomic or ESTs) related to known transcription factors were identified in the *Arabidopsis thaliana* GenBank database using the tblastn sequence analysis program using default parameters and a P-value cutoff threshold of -4 or -5 or lower, depending on the length of the query sequence. Putative transcription factor sequence hits were then screened to identify those containing particular sequence strings. If the sequence hits contained such sequence strings, the sequences were confirmed as transcription factors.

Alternatively, Arabidopsis thaliana cDNA libraries derived from different tissues or treatments, or genomic libraries were screened to identify novel members of a transcription family using a low stringency hybridization approach. Probes were synthesized using gene specific primers in a standard PCR reaction (annealing temperature 60° C) and labeled with ³²P dCTP using the High Prime DNA Labeling Kit (Boehringer Mannheim). Purified radiolabelled probes were added to filters immersed in Church hybridization medium (0.5 M NaPO₄ pH 7.0, 7% SDS, 1 % w/v bovine serum albumin) and hybridized overnight at 60 °C with shaking. Filters were washed two times for 45 to 60 minutes with 1xSCC, 1% SDS at 60° C.

To identify additional sequence 5' or 3' of a partial cDNA sequence in a cDNA library, 5' and 3' rapid amplification of cDNA ends (RACE) was performed using the MarathonTM cDNA amplification kit (Clontech, Palo Alto, CA). Generally, the method entailed first isolating poly(A) mRNA, performing first and second strand cDNA synthesis to generate double stranded cDNA, blunting cDNA ends, followed by ligation of the MarathonTM Adaptor to the cDNA to form a library of adaptor-ligated ds cDNA.

Gene-specific primers were designed to be used along with adaptor specific primers for both 5' and 3' RACE reactions. Nested primers, rather than single primers, were used to increase PCR specificity. Using 5' and 3' RACE reactions, 5' and 3' RACE fragments were obtained, sequenced and cloned. The process can be repeated until 5' and 3' ends of the full-length gene were identified. Then the full-length cDNA was generated by PCR using primers specific to 5' and 3' ends of the gene by end-to-end PCR.

EXAMPLE II. CONSTRUCTION OF EXPRESSION VECTORS

5

10

30

35

The sequence was amplified from a genomic or cDNA library using primers 15 specific to sequences upstream and downstream of the coding region. The expression vector was pMEN20 or pMEN65, which are both derived from pMON316 (Sanders et al, (1987) Nucleic Acids Research 15:1543-58) and contain the CaMV 35S promoter to express transgenes. To clone the sequence into the vector, both pMEN20 and the amplified DNA fragment were digested separately with SalI and NotI restriction enzymes at 37° C for 2 hours. 20 The digestion products were subject to electrophoresis in a 0.8% agarose gel and visualized by ethidium bromide staining. The DNA fragments containing the sequence and the linearized plasmid were excised and purified by using a Qiaquick gel extraction kit (Qiagen, CA). The fragments of interest were ligated at a ratio of 3:1 (vector to insert). Ligation reactions using T4 DNA ligase (New England Biolabs, MA) were carried out at 16° C for 16 25 hours. The ligated DNAs were transformed into competent cells of the E. coli strain DH5alpha by using the heat shock method. The transformations were plated on LB plates containing 50 mg/l kanamycin (Sigma).

Individual colonies were grown overnight in five milliliters of LB broth containing 50 mg/l kanamycin at 37° C. Plasmid DNA was purified by using Qiaquick Mini Prep kits (Qiagen, CA).

EXAMPLE III. TRANSFORMATION OF AGROBACTERIUM WITH THE EXPRESSION VECTOR

After the plasmid vector containing the gene was constructed, the vector was used to transform *Agrobacterium tumefaciens* cells expressing the gene products. The stock of *Agrobacterium tumefaciens* cells for transformation were made as described by Nagel et al. (1990) FEMS Microbiol Letts. 67: 325-328. *Agrobacterium* strain ABI was grown in 250 ml

LB medium (Sigma) overnight at 28°C with shaking until an absorbance (A_{600}) of 0.5 – 1.0 was reached. Cells were harvested by centrifugation at 4,000 x g for 15 min at 4° C. Cells were then resuspended in 250 μ l chilled buffer (1 mM HEPES, pH adjusted to 7.0 with KOH). Cells were centrifuged again as described above and resuspended in 125 μ l chilled buffer. Cells were then centrifuged and resuspended two more times in the same HEPES buffer as described above at a volume of 100 μ l and 750 μ l, respectively. Resuspended cells were then distributed into 40 μ l aliquots, quickly frozen in liquid nitrogen, and stored at -80° C.

5

10

15

20

25

30

35

Agrobacterium cells were transformed with plasmids prepared as described above following the protocol described by Nagel et al. For each DNA construct to be transformed, 50-100 ng DNA (generally resuspended in 10 mM Tris-HCl, 1 mM EDTA, pH 8.0) was mixed with 40 µl of Agrobacterium cells. The DNA/cell mixture was then transferred to a chilled cuvette with a 2mm electrode gap and subject to a 2.5 kV charge dissipated at 25 µF and 200 µF using a Gene Pulser II apparatus (Bio-Rad). After electroporation, cells were immediately resuspended in 1.0 ml LB and allowed to recover without antibiotic selection for 2-4 hours at 28° C in a shaking incubator. After recovery, cells were plated onto selective medium of LB broth containing 100 µg/ml spectinomycin (Sigma) and incubated for 24-48 hours at 28° C. Single colonies were then picked and inoculated in fresh medium. The presence of the plasmid construct was verified by PCR amplification and sequence analysis.

EXAMPLE IV. TRANSFORMATION OF ARABIDOPSIS PLANTS WITH AGROBACTERIUM TUMEFACIENS WITH EXPRESSION VECTOR

After transformation of Agrobacterium tumefaciens with plasmid vectors containing the gene, single Agrobacterium colonies were identified, propagated, and used to transform Arabidopsis plants. Briefly, 500 ml cultures of LB medium containing 50 mg/l kanamycin were inoculated with the colonies and grown at 28° C with shaking for 2 days until an absorbance (A_{600}) of > 2.0 is reached. Cells were then harvested by centrifugation at 4,000 x g for 10 min, and resuspended in infiltration medium (1/2 X Murashige and Skoog salts (Sigma), 1 X Gamborg's B-5 vitamins (Sigma), 5.0% (w/v) sucrose (Sigma), 0.044 μ M benzylamino purine (Sigma), 200 μ l/L Silwet L-77 (Lehle Seeds) until an absorbance (A_{600}) of 0.8 was reached.

Prior to transformation, Arabidopsis thaliana seeds (ecotype Columbia) were sown at a density of ~10 plants per 4" pot onto Pro-Mix BX potting medium (Hummert International) covered with fiberglass mesh (18 mm X 16 mm). Plants were grown under continuous illumination (50-75 μE/m²/sec) at 22-23° C with 65-70% relative humidity. After about 4 weeks, primary inflorescence stems (bolts) are cut off to encourage growth of

multiple secondary bolts. After flowering of the mature secondary bolts, plants were prepared for transformation by removal of all siliques and opened flowers.

The pots were then immersed upside down in the mixture of Agrobacterium infiltration medium as described above for 30 sec, and placed on their sides to allow draining into a 1' x 2' flat surface covered with plastic wrap. After 24 h, the plastic wrap was removed and pots are turned upright. The immersion procedure was repeated one week later, for a total of two immersions per pot. Seeds were then collected from each transformation pot and analyzed following the protocol described below.

EXAMPLE V. IDENTIFICATION OF ARABIDOPSIS PRIMARY TRANSFORMANTS

Seeds collected from the transformation pots were sterilized essentially as follows. Seeds were dispersed into in a solution containing 0.1% (v/v) Triton X-100 (Sigma) and sterile H₂O and washed by shaking the suspension for 20 min. The wash solution was then drained and replaced with fresh wash solution to wash the seeds for 20 min with shaking. After removal of the second wash solution, a solution containing 0.1% (v/v) Triton X-100 and 70% ethanol (Equistar) was added to the seeds and the suspension was shaken for 5 min. After removal of the ethanol/detergent solution, a solution containing 0.1% (v/v) Triton X-100 and 30% (v/v) bleach (Clorox) was added to the seeds, and the suspension was shaken for 10 min. After removal of the bleach/detergent solution, seeds were then washed five times in sterile distilled H₂O. The seeds were stored in the last wash water at 4° C for 2 days in the dark before being plated onto antibiotic selection medium (1 X Murashige and Skoog salts (pH adjusted to 5.7 with 1M KOH), 1 X Gamborg's B-5 vitamins, 0.9% phytagar (Life Technologies), and 50 mg/l kanamycin). Seeds were germinated under continuous illumination (50-75 µE/m²/sec) at 22-23° C. After 7-10 days of growth under these conditions, kanamycin resistant primary transformants (T1 generation) were visible and obtained. These seedlings were transferred first to fresh selection plates where the seedlings continued to grow for 3-5 more days, and then to soil (Pro-Mix BX potting medium).

Primary transformants were crossed and progeny seeds (T₂) collected; kanamycin resistant seedlings were selected and analyzed. The expression levels of the recombinant polynucleotides in the transformants varies from about a 5% expression level increase to a least a 100% expression level increase. Similar observations are made with respect to polypeptide level expression.

EXAMPLE VI. IDENTIFICATION OF ARABIDOPSIS PLANTS WITH TRANSCRIPTION FACTOR GENE KNOCKOUTS

5

10

15

20

25

The screening of insertion mutagenized *Arabidopsis* collections for null mutants in a known target gene was essentially as described in Krysan et al (1999) <u>Plant Cell</u> 11:2283-2290. Briefly, gene-specific primers, nested by 5-250 bases to each other, were designed from the 5' and 3' regions of a known target gene. Similarly, nested sets of primers were also created specific to each of the T-DNA or transposon ends (the "right" and "left" borders). All possible combinations of gene specific and T-DNA/transposon primers were used to detect by PCR an insertion event within or close to the target gene. The amplified DNA fragments were then sequenced which allows the precise determination of the T-DNA/transposon insertion point relative to the target gene. Insertion events within the coding or intervening sequence of the genes were deconvoluted from a pool comprising a plurality of insertion events to a single unique mutant plant for functional characterization. The method is described in more detail in Yu and Adam, US Application Serial No. 09/177,733 filed October 23, 1998.

15 EXAMPLE VII. IDENTIFICATION OF PATHOGEN INDUCED GENES

5

10

20

25

30

In some instances, expression patterns of the pathogen induced genes (such as defense genes) was monitored by microarray experiments. cDNAs were generated by PCR and resuspended at a final concentration of ~ 100 ng/ul in 3X SSC or 150mM Na-phosphate (Eisen and Brown (1999) *Meth. in Enzymol.* 303:179-205). The cDNAs were spotted on microscope glass slides coated with polylysine. The prepared cDNAs were aliquoted into 384 well plates and spotted on the slides using an x-y-z gantry (OmniGrid) purchased from GeneMachines (Menlo Park, CA) outfitted with quill type pins purchased from Telechem International (Sunnyvale, CA). After spotting, the arrays were cured for a minimum of one week at room temperature, rehydrated and blocked following the protocol recommended by Eisen and Brown (1999).

Sample total RNA (10 ug) samples were labeled using fluorescent Cy3 and Cy5 dyes. Labeled samples were resuspended in 4X SSC/0.03% SDS/4 ug salmon sperm DNA/2 ug tRNA/50mM Na-pyrophosphate, heated for 95°C for 2.5 minutes, spun down and placed on the array. The array was then covered with a glass coverslip and placed in a sealed chamber. The chamber was then kept in a water bath at 62°C overnight. The arrays were washed as described in Eisen and Brown (1999) and scanned on a General Scanning 3000 laser scanner. The resulting files are subsequently quantified using Imagene a software purchased from BioDiscovery (Los Angeles, CA).

EXAMPLE VIII. IDENTIFICATION OF PATHOGEN TOLERANCE PHENOTYPE IN OVEREXPRESSOR OR GENE KNOCKOUT PLANTS

5

10

15

20

25

30

35

Experiments were performed to identify those transformants or knockouts that exhibited an improved pathogen tolerance. For such studies, the transformants were exposed to biotropic fungal pathogens, such as Erisyphe orontii, and necrotropic fungal pathogens, such as Fusarium oxysporum. Fusarium oxysporum isolates cause vascular wilts and damping off of various annual vegetables, perennials and weeds (Mauch-Mani and Slusarenko (1994) Molecular Plant-Microbe Interactions 7: 378-383). For Fusarium oxysporum experiments, plants grown on petri dishes were sprayed with a fresh spore suspension of F. oxysporum. The spore suspension was prepared as follows: A plug of fungal hyphae from a plate culture was placed on a fresh potato dextrose agar plate and allowed to spread for one week. 5 ml sterile water was then added to the plate, swirled, and pipetted into 50 ml Armstrong Fusarium medium. Spores were grown overnight in Fusarium medium and then sprayed onto plants using a Preval paint sprayer. Plant tissue was harvested and frozen in liquid nitrogen 48 hours post infection.

Erysiphe orontii is a causal agent of powdery mildew. For Erysiphe orontii experiments, plants were grown approximately 4 weeks in a greenhouse under 12 hour light (20 C, ~30% relative humidity (rh)). Individual leaves were infected with E. orontii spores from infected plants using a camel's hair brush, and the plants were transferred to a Percival growth chamber (20 C, 80% rh.). Plant tissue was harvested and frozen in liquid nitrogen 7 days post infection.

Botrytis cinerea is a necrotrophic pathogen. Botrytis cinerea was grown on potato dextrose agar in the light. A spore culture was made by spreading 10 ml of sterile water on the fungus plate, swirling and transferring spores to 10 ml of sterile water. The spore inoculum (approx. 105 spores/ml) was used to spray 10 day-old seedlings grown under sterile conditions on MS (-sucrose) media. Symptoms were evaluated every day up to approximately 1 week.

Infection with bacterial pathogens Pseudomonas syringae pv maculicola strain 4326 and pv maculicola strain 4326 was performed by hand inoculation at two doses. Two inoculation doses allows the differentiation between plants with enhanced susceptibility and plants with enhanced resistance to the pathogen. Plants were grown for 3 weeks in the greenhouse, then transferred to the growth chamber for the remainder of their growth. Psm ES4326 was hand inoculated with 1 ml syringe on 3 fully-expanded leaves per plant (4 1/2 wk old), using at least 9 plants per overexpressing line at two inoculation doses, OD=0.005 and OD=0.0005. Disease scoring occurred at day 3 post-inoculation with pictures of the plants and leaves taken in parallel.

Table 3 shows the phenotypes observed for particular overexpressor or knockout plants and provides the SEQ ID No., the internal reference code (GID), whether a knockout or overexpressor plant was analyzed and the observed phenotype.

Table 3

SEQ ID No.	GID	Knockout (KO) or overexpressor (OE)	Phenotype
1	G188	KO	Increased susceptibility to Fusarium
3	G616	OE	Increased tolerance to Erysiphe
5	G19	OE	Increased tolerance to Erysiphe
7.	Ġ261	OE	Increased susceptibility to Botrytis
9	G28	OE	Increased resistance to Erysiphe
11	G869	OE	Increased susceptibility to Fusarium
13	G237	OE	Increased tolerance to Erysiphe
15	G409	OE	Increased tolerance to Erysiphe
17	G418	OE	Increased tolerance to Pseudomonas
19	G591	OE	Increased tolerance to Erysiphe
21	G525	OE	Increased tolerance to Pseudomonas
23	G545	OE	Increased susceptibility to Pseudomonas, Erysiphe and Fusarium
25	G865	OE	Increased susceptibility to Erysiphe and Botrytis
27	G881	OE	Increased susceptibility to Erysiphe and Botrytis
29	G896	KO	Increased susceptibility to Fusarium
31	G378	OE	Increased resistance to Erysiphe
33	G569	OE .	Decreased expression of defense genes
35	G558	OE	Increased expression of defense genes

For a particular overexpressor that shows an increased susceptibility to a pathogen, it may be more useful to select a plant with a decreased expression of the particular transcription factor. For a particular knockout that shows an increased susceptibility to a pathogen, it may be more useful to select a plant with an increased expression of the particular transcription factor.

Other than Fusarium oxysporum, Erysyphe orontii, the transgenic plants are more tolerant to Sclerotinia spp., soil-borne oomycetes, foliar oomycetes, Botrytis spp., Rhizoctonia spp, Verticillium dahliae/albo-atrum, Alternaria spp., rusts, Mycosphaerella spp, Fusarium solani, or the like. The transgenic plants are more resistant to fungal diseases such as rusts, smuts, wilts, yellows, root rot, leaf drop, ergot, leaf blight of potato, brown spot of rice, leaf blight, late blight, powdery mildew, downy mildew, and the like; viral diseases such as sugarcane mosaic, cassava mosaic, sugar beet yellows, plum pox, barley yellow dwarf, tomato yellow leaf curl, tomato spotted wilt virus, and the like; bacterial diseases such as citrus canker, bacterial leaf blight, bacterial will, soft rot of vegetables, and the like; nematode diseases such as root knot, sugar beet cyst nematode or the like.

5

10

15

EXAMPLE IX. IDENTIFICATION OF HOMOLOGOUS SEQUENCES

5

10

15

20

25

30

35

Homologous sequences from *Arabidopsis* and plant species other than *Arabidopsis* were identified using database sequence search tools, such as the Basic Local Alignment Search Tool (BLAST) (Altschul et al. (1990) <u>J. Mol. Biol.</u> 215:403-410; and Altschul et al. (1997) <u>Nucl. Acid Res.</u> 25: 3389-3402). The tblastx sequence analysis programs were employed using the BLOSUM-62 scoring matrix (Henikoff, S. and Henikoff, J. G. (1992) <u>Proc. Natl. Acad. Sci. USA</u> 89: 10915-10919).

Identified *Arabidopsis* homologous sequences are provided in Figure 2 and included in the Sequence Listing. The percent sequence identity among these sequences is as low as 47% sequence identity. Additionally, the entire NCBI GenBank database was filtered for sequences from all plants except *Arabidopsis thaliana* by selecting all entries in the NCBI GenBank database associated with NCBI taxonomic ID 33090 (Viridiplantae; all plants) and excluding entries associated with taxonomic ID 3701 (*Arabidopsis thaliana*). These sequences were compared to sequences representing genes of SEQ IDs Nos. 1-58 on 9/26/2000 using the Washington University TBLASTX algorithm (version 2.0a19MP). For each gene of SEQ IDs Nos. 1-58, individual comparisons were ordered by probability score (P-value), where the score reflects the probability that a particular alignment occurred by chance. For example, a score of 3.6e-40 is 3.6 x 10⁻⁴⁰. For up to ten species, the gene with the lowest P-value (and therefore the most likely homolog) is listed in Figure 3.

In addition to P-values, comparisons were also scored by percentage identity. Percentage identity reflects the degree to which two segments of DNA or protein are identical over a particular length. The ranges of percent identity between the non-Arabidopsis genes shown in Figure 3 and the Arabidopsis genes in the sequence listing are: SEQ ID No. 1: 38%-76%; SEQ ID No. 3: 36%-72%; SEQ ID No. 5: 51%-75%; SEQ ID No. 7: 37%-76%; SEQ ID No. 9: 48%-75%; SEQ ID No. 11: 31%-68%; SEQ ID No. 13: 59%-81%; SEQ ID No. 15: 49%-81%; SEQ ID No. 17: 53%-87%; SEQ ID No. 19: 48%-84%; SEQ ID No. 21: 73%-89%; SEQ ID No. 23: 52%-64%; SEQ ID No. 25: 48%-83%; SEQ ID No. 27: 35%-92%; SEQ ID No. 29: 56%-89%; SEQ ID No. 31: 50%-90%; SEQ ID No. 33: 50%-93%; SEQ ID No. 35: 52%-81%; SEQ ID No. 37: 75%-81%; SEQ ID No. 39: 35%-72%; SEQ ID No. 41: 55%-89%; SEQ ID No. 43: 56%-77%; SEQ ID No. 45: 34%-72%; SEQ ID No. 47: 51%-86%; SEQ ID No. 49: 46%-86%; SEQ ID No. 51: 58%-80%; SEQ ID No. 53: 46%-55%; SEQ ID No. 55: 84%-89%; and SEQ ID No. 57: 43%-71%.

The polynucleotides and polypeptides in the Sequence Listing and the identified homologous sequences may be stored in a computer system and have associated or linked with the sequences a function, such as that the polynucleotides and polypeptides are useful for modifying the pathogen tolerance of a plant.

All references, publications, patents and other documents herein are incorporated by reference in their entirety for all purposes. Although the invention has been described with reference to the embodiments and examples above, it should be understood that various modifications can be made without departing from the spirit of the invention.

What is claimed is:

15

20

30

1. A transgenic plant with modified pathogen tolerance, which plant comprises a recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- 5 (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-29, or a complementary nucleotide sequence thereof:
 - (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a);
- 10 (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID Nos. 2N-1, where N=1-29, or a complementary nucleotide sequence thereof;
 - (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of (c);
 - (e) a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence of one or more of: (a), (b), (c), or (d);
 - (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e);
 - (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide that modifies a plant's pathogen tolerance;
 - (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g);
 - (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide sequence of any of (a)-(g);
- 25 (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29;
 - (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29; and
 - (l) a nucleotide sequence which encodes a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-29.
 - 2. The transgenic plant of claim 1, further comprising a constitutive, inducible, or tissue-active promoter operably linked to said nucleotide sequence.
- 35 3. The transgenic plant of claim 1, wherein the plant is selected from the group consisting of: soybean, wheat, corn, potato, cotton, rice, oilseed rape, sunflower, alfalfa, sugarcane, turf, banana, blackberry, blueberry, strawberry, raspberry, cantaloupe, carrot,

cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits, and vegetable brassicas.

- 5 4. An isolated or recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of:
 - (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-29, or a complementary nucleotide sequence thereof;
- 10 (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a);

15

30

- (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID Nos. 2N-1, where N=1-29, or a complementary nucleotide sequence thereof;
- (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of (c);
- (e) a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence of one or more of: (a), (b), (c), or (d);
- (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e);
- 20 (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide having a biological activity that modifies a plant's pathogen tolerance;
 - (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g);
- 25 (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide sequence of any of (a)-(g);
 - (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29;
 - (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29; and
 - (1) a nucleotide sequence which encodes a conserved domain of a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-29.
- The isolated or recombinant polynucleotide of claim 4, further comprising a constitutive, inducible, or tissue-active promoter operably linked to the nucleotide sequence.

6. A cloning or expression vector comprising the isolated or recombinant polynucleotide of claim 4.

- 7. A cell comprising the cloning or expression vector of claim 6.
- 8. A transgenic plant comprising the isolated or recombinant polynucleotide of claim 4.
 - 9. A composition produced by one or more of:

5

15

20

25

30

- (a) incubating one or more polynucleotide of claim 4 with a nuclease;
- 10 (b) incubating one or more polynucleotide of claim 4 with a restriction enzyme;
 - (c) incubating one or more polynucleotide of claim 4 with a polymerase;
 - (d) incubating one or more polynucleotide of claim 4 with a polymerase and a primer;
 - (e) incubating one or more polynucleotide of claim 4 with a cloning vector, or
 - (f) incubating one or more polynucleotide of claim 4 with a cell.
 - 10. A composition comprising two or more different polynucleotides of claim 4.
 - 11. An isolated or recombinant polypeptide comprising a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4.
 - 12. A plant comprising an isolated polypeptide of claim 11.
 - 13. A method for producing a plant having a modified pathogen tolerance, the method comprising altering the expression of the isolated or recombinant polynucleotide of claim 4 or the expression levels or activity of a polypeptide of claim 11 in a plant, thereby producing a modified plant, and selecting the modified plant for improved pathogen tolerance thereby providing the modified plant with a modified pathogen tolerance.
 - 14. The method of claim 13, wherein the polynucleotide is a polynucleotide of claim 4.
 - 15. A method of identifying a factor that is modulated by or interacts with a polypeptide encoded by a polynucleotide of claim 4, the method comprising:
 - (a) expressing a polypeptide encoded by the polynucleotide in a plant; and
- (b) identifying at least one factor that is modulated by or interacts with the polypeptide.

16. The method of claim 15, wherein the identifying is performed by detecting binding by the polypeptide to a promoter sequence, or detecting interactions between an additional protein and the polypeptide in a yeast two hybrid system.

- 5 17. The method of claim 15, wherein the identifying is performed by detecting expression of a factor by hybridization to a microarray, subtractive hybridization or differential display.
 - 18. A method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest, the method comprising:
 - (a) placing the molecule in contact with a plant comprising the polynucleotide or polypeptide encoded by the polynucleotide of claim 4; and,
 - (b) monitoring one or more of:

10

15

20

25

method comprising:

- (i) expression level of the polynucleotide in the plant;
- (ii) expression level of the polypeptide in the plant;
- (iii) modulation of an activity of the polypeptide in the plant; or
- (iv) modulation of an activity of the polynucleotide in the plant.
- 19. An integrated system, computer or computer readable medium comprising one or more character strings corresponding to a polynucleotide of claim 4, or to a polypeptide encoded by the polynucleotide.
- 20. The integrated system, computer or computer readable medium of claim 19, further comprising a link between said one or more sequence strings to a modified plant pathogen tolerance phenotype.
- 21. A method of identifying a sequence similar or homologous to one or more polynucleotides of claim 4, or one or more polypeptides encoded by the polynucleotides, the
 - (a) providing a sequence database; and,
- 30 (b) querying the sequence database with one or more target sequences corresponding to the one or more polynucleotides or to the one or more polypeptides to identify one or more sequence members of the database that display sequence similarity or homology to one or more of the one or more target sequences.
- The method of claim 21, wherein the querying comprises aligning one or more of the target sequences with one or more of the one or more sequence members in the sequence database.

23. The method of claim 21, wherein the querying comprises identifying one or more of the one or more sequence members of the database that meet a user-selected identity criteria with one or more of the target sequences.

5

- 24. The method of claim 21, further comprising linking the one or more of the polynucleotides of claim 4, or encoded polypeptides, to a modified plant pathogen tolerance phenotype.
- 10 25. A plant comprising altered expression levels of an isolated or recombinant polynucleotide of claim 4.
 - 26. A plant comprising altered expression levels or the activity of an isolated or recombinant polypeptide of claim 11.

15

27. A plant lacking a nucleotide sequence encoding a polypeptide of claim 11.

Figure 1

SEQ ID No.	GID	cDNA or protein	conserved domain
1	G188	cDNA	
2	G188	protein	175-222
3	G616	cDNA	
4	G616	protein	39-95
5	G19	cDNA	
6	G19	protein	76-145
7	G261	cDNA	•
8	G261	protein	16-104
9	G28	cDNA	
10	G28	protein	145-213
11	G869	cDNA	
12	G869	protein	109-177
13	G237	cDNA	
14	G237	protein	11-113
15	G409	cDNA	
16	G409	protein	64-124
17	G418	cDNA	,
18	G418	protein	500-560
19	G591	cDNA	
20	G591	protein	143-240
21	G525	cDNA	
22	G525	protein	23-167
23	G545	cDNA	
24	G545	protein	82-102, 136-154
25	G865	cDNA	
26	G865	protein	36-103
27	G881	cDNA	
28	G881	protein	176-233
29	G896	cDNA	
30	G896	protein	18-39
31	G378	cDNA	
32	G378	protein	196-237
33	G569	cDNA	
34	G569	protein	90-153
35	G558	cDNA	
36	G558	prótein	45-105

Figure 2

SEQ ID No.	GID	homolog	cDNA or protein	conserved domain
37	G1396	homolog of G1394	cDNA	
38	G1396	homolog of G1394	protein	entire protein
39	G265	homolog of G261	cDNA	
40	G265	homolog of G261	protein	14-105
41	G1006	homolog of G28	cDNA	
42	G1006	homolog of G28	protein	114-182
43	G1309	homolog of G237	cDNA	
44	G1309	homolog of G237	protein	9-114
45	G2550	homolog of G418	cDNA	
46	G2550	homolog of G418	protein	348-408
47	G965	homolog of G418	cDNA	
48	G965	homolog of G418	protein	423-486
49	G793	homolog of G591	cDNA	
50	G793	homolog of G591	protein	151-206
51	G764	homolog of G525	cDNA	
52	G764	homolog of G525	protein	27-171 ·
53	G350	homolog of G545	cDNA	
54	G350	homolog of G545	protein	91-113,150-170
55	G986	homolog of G881	cDNA	
56	G986	homolog of G881	protein	146-203
57	G1349	homolog of G896	cDNA	
58	G1349	homolog of G896	protein	13-63

Figure 3A

1 G188 7779802 5.20E-36 Lotus japonicus	SEQ ID No.	GID	Genbank NID	P-value	Species
1 G188 7284340 2.10E-34 Glycine max 1 G188 9361307 1.20E-27 Triticum aestivum 1 G188 6359152 3.60E-22 Lycopersicon esculentum 1 G188 6529152 3.60E-22 Lycopersicon esculentum 1 G188 8748477 7.70E-21 Medicago truncatula 1 G188 8756433 7.10E-14 Zea mays 1 G188 6896287 4.10E-12 Pinus taeda 1 G188 6696287 4.10E-12 Pinus taeda 1 G188 562242 9.00E-12 Brassica rapa 3 G616 769230 5.90E-33 Glycine mäx 3 G616 7591307 1.10E-21 Gossypium arboreum 3 G616 7591307 1.10E-21 Oryza sativa 3 G616 8658771 1.50E-21 Oryza sativa 3 G616 8058535 7.40E-20 Antirrinum tuberosum 3					
1 G188 9361307 1.20E-27 Triticum aestivum 1 G188 7340336 1.10E-22 Oryza sativa 1 G188 6529152 3.60E-22 Lycopersicon esculentum 1 G188 6456433 7.10E-14 Zea mays 1 G188 6596287 4.10E-12 Sorghum bicolor 1 G188 65696287 4.10E-12 Pinus taeda 1 G188 562242 9.00E-12 Brassisca rapa 3 G616 7719440 8.30E-37 Lotus japonicus 3 G616 77019440 8.30E-37 Lotus japonicus 3 G616 7501307 1.10E-21 Sosypjum abroeum 3 G616 8858771 1.50E-21 Solanum tuberosum 3 G616 8858771 1.50E-21 Gossypjum abroeum 3 G616 6358535 7.40E-20 Antirrhinum majus 3 G616 6358535 7.50E-20 Antirrhinum majus 3<					
1 G188 7340336 1.10E-22 Cyzopersicon esculentum 1 G188 6529152 3.60E-22 Lycopersicon esculentum 1 G188 5456433 7.10E-14 Zea mays 1 G188 5456433 7.10E-12 Sorghum bicolor 1 G188 6696287 4.10E-12 Pinus taeda 1 G188 562242 9.00E-12 Brassica rapa 3 G616 7719440 8.30E-37 Lotus japonicus 3 G616 779230 5.90E-33 Glycine max 3 G616 8071307 1.10E-21 Gossypium arboreum 3 G616 8071309 1.50E-21 Solanum tuberosum 3 G616 8058711 1.50E-21 Solanum tuberosum 3 G616 6358532 5.80E-20 Antirrhinum fautur 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianu					
1 G188 6529152 3.60E-22 Lycopersicon esculentum 1 G188 5456433 7.10E-14 Zea mays 1 G188 5456433 7.10E-14 Zea mays 1 G188 6696287 4.10E-12 Pinus taeda 1 G188 562242 9.00E-12 Brassica rapa 3 G616 7719440 8.30E-37 Lotus japonicus 3 G616 77501307 1.10E-21 Gossypium arboreum 3 G616 7501307 1.10E-21 Gossypium arboreum 3 G616 885871 1.50E-21 Solanum tuberosum 3 G616 885871 1.50E-21 Gossypium hirsutum 3 G616 885871 1.50E-21 Gossypium hirsutum 3 G616 835853 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp					
1 G188 8748477 7.70E-21 Medicago truncatula 1 G188 5456433 7.10E-14 Zea mays 1 G188 698287 1.60E-12 Sorghum bicolor 1 G188 696287 4.10E-12 Pinus taeda 1 G188 562242 9.00E-12 Brassica rapa 3 G616 779220 5.90E-33 Glycine max 3 G616 7501307 1.10E-21 Sossyptium arboreum 3 G616 8071090 1.50E-21 Solanum tuberosum 3 G616 8057131 1.50E-21 Solanum tuberosum 3 G616 6358532 5.80E-20 Antirrhinum grantitum 3 G616 6358535 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antir					
1 G188 5456433 7.10E-14 Zea mays 1 G188 9302479 1.60E-12 Sorghum bicolor 1 G188 6696287 4.10E-12 Pinus taeda 1 G188 562242 9.00E-12 Brassica rapa 3 G616 779240 8.30E-37 Lotus japonicus 3 G616 7692230 5.90E-33 Glycine max 3 G616 7692230 5.90E-33 Glycine max 3 G616 7692731 1.50E-21 Gossypium arboreum 3 G616 8858771 1.50E-21 Gossypium insutur 3 G616 8585771 1.50E-21 Gossypium hirsutur 3 G616 8358532 7.0E-20 Antirrhinum majus suturinum majus 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus 3 G616 63585332 2.80E-34 Citrus x paradisi<					
1 G188 9302479 1.60E-12 Sorghum bicolor 1 G188 6696287 4.10E-12 Pinus taeda 1 G188 6692242 9.00E-12 Prassica rapa 3 G616 7719440 8.30E-37 Lotus japonicus 3 G616 769230 5.90E-33 Glyche max 3 G616 769230 5.90E-33 Glyche max 3 G616 8071090 1.50E-21 Sossypium hirsutum 3 G616 858771 1.50E-21 Gossypium hirsutum 3 G616 5047315 1.50E-21 Gossypium hirsutum 3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antir				_	
1 G188 6696287 4.10E-12 Pinus taeda 1 G188 562242 9.00E-12 Brassica rapa 3 G616 779440 8.30E-37 Lotus japonicus 3 G616 7501307 1.10E-21 Gossypium arboreum 3 G616 8071090 1.50E-21 Solanum tuberosum 3 G616 8858771 1.50E-21 Oryza sativa 3 G616 5047315 1.50E-21 Gossypium hirsutum 3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358539 <td></td> <td></td> <td></td> <td></td> <td></td>					
1 G188 562242 9.00E-12 Brassica rapa 3 G616 7719440 8.30E-37 Lotus japonicus 3 G616 769230 5.90E-33 Glycine max 3 G616 8071090 1.50E-21 Gossypium arboreum 3 G616 8085771 1.50E-21 Oryza sativa 3 G616 5047315 1.50E-21 Gossypium hirsutum 3 G616 5058532 5.80E-20 Antirrhinum graniticum 3 G616 6358532 5.80E-20 Antirrhinum majus subsp. linkianum 3 G616 6358535 7.40E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Litrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7478682 1.30E-30 Glycine max 5 G19 7426302 8.30E-26 Gossypium arbore					
3 G616 7719440 8.30E-37 Lotus japonicus 3 G616 769230 5.90E-33 Glycine max 3 G616 7501307 1.10E-21 Gossypium arboreum 3 G616 8071090 1.50E-21 Solanum tuberosum 3 G616 5047315 1.50E-21 Oryza sativa 3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 6358535 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358535 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 5					
3 G616 7692230 5.90E-33 Glycine max 3 G616 8071090 1.50E-21 Gossyplum arboreum 3 G616 8071090 1.50E-21 Oryza sativa 3 G616 8058771 1.50E-21 Oryza sativa 3 G616 5047315 1.50E-21 Gossyplum hirsuturm 3 G616 63586532 5.80E-20 Antirrhinum graniticum 3 G616 63586532 5.80E-20 Antirrhinum graniticum 3 G616 63586535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7476682 1.30E-30 Glycine max 6 G19 6654934 1.20E-28 Medicago truncatula 5 G19 7624302 8.30E-26 Gossyplum arboreum 5 G19 7624302 8.30E-26 Gossyplum arboreum 5 G19 9425363 2.90E-25 Ricinus communis 5 G19 9419304 6.00E-25 Ricinus communis 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 97720316 8.80E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 733914 3.50E-41 Citrus unshiu 7 G261 7233914 3.50E-41 Citrus unshiu 7 G261 728275 4.20E-62 Mesembryanthemum crystallinum 9 G28 7628275 4.20E-62 Mesembryanthemum crystallinum 9 G28 762876 4.20E-62 Mesembryanthemum crystallinum 9 G28 762601 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Glycine max 11 G869 2213784 1.30E-30 Oryza sativa					
3 G616 7501307 1.10E-21 Gossypium arboreum 3 G616 8858771 1.50E-21 Solanum tuberosum 3 G616 8858771 1.50E-21 Oryza sativa 3 G616 5047315 1.50E-21 Gossypium hirsutum 3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 6358535 7.00E-20 Antirrhinum graniticum 3 G616 6358538 7.50E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 8789223 2.80E-34 Lycopersicon esculentum 5 G19 8789223 2.80E-34 Lycopersicon esculentum 5 G19 8789223 2.80E-28 Medicago Iruncatula 5 G19 7476882 1.30E-26 Medicago Iruncatula 5 G19 7624302 8.30E-25					
3 G616 8071090 1.50E-21 Solanum tuberosum 3 G616 8858771 1.50E-21 Oryza sativa 3 G616 5047315 1.50E-21 Oryza sativa 3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 2826867 7.00E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 9434234 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 3264766 5.50E-26 Prunu					
3 G616 8858771 1.50E-21 Oryza sativa 3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 6358535 7.00E-20 Antirrhinum majus subsp. linkianum 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 5 G19 8789223 2.80E-34 Lycopersicon esculentum 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7478682 1.30E-30 Glycine max 5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 7624302 <td></td> <td></td> <td></td> <td></td> <td></td>					
3 G616 5047315 1.50E-21 Gossypium hirsutum 3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 2826867 7.00E-20 Antirrhinum majus 3 G616 6358538 7.50E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7478682 1.30E-30 Glycine max 5 G19 7624302 8.30E-26 Prunus armeniaca 5 G19 7624302 8.30E-25 Trilicum aestivum					
3 G616 6358532 5.80E-20 Antirrhinum graniticum 3 G616 2826867 7.00E-20 Antirrhinum majus subsp. linkianum 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7478682 1.30E-30 Glycine max 5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 3425302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25	3				
3 G616 2826867 7.00E-20 Antirrhinum majus 3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7478682 1.30E-30 Glycine max 5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Trilicum aestivum 5 G19 9425363 2.90E-25 Trilicum aestivum 5 G19 9449304 6.00E-25 Hordeum vulgare 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 9479304 8.00E-25 Lotus japonicus <td></td> <td></td> <td></td> <td></td> <td></td>					
3 G616 6358535 7.40E-20 Antirrhinum majus subsp. linkianum 3 G616 6358538 7.50E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7478682 1.30E-30 Glycine max 5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25 Ricinus communis 5 G19 9419304 6.00E-25 Ricinus communis 5 G19 9419304 6.00E-25 Ricinus communis 7 G261 5821137 5.10E-93 Nicotiana tabacu					
3 G616 6358538 7.50E-20 Antirrhinum braun-blanquetii 5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7478682 1.30E-30 Glycine max 5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9429304 6.00E-25 Hordeum vulgare 5 G19 9419304 6.00E-25 Hordeum vulgare 7 G261 5821137 5.10E-93 Nicotiana tabacum					
5 G19 8789223 2.80E-34 Citrus x paradisi 5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7478682 1.30E-30 Glycine max 5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25 Ricinus communis 5 G19 9425363 2.90E-25 Hordeum vulgare 5 G19 9425363 2.90E-25 Hordeum vulgare 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 9419304 6.00E-25 Hordeum vulgare 7 G261 5821137 5.10E-93 Nicotiana tabacum <t< td=""><td></td><td></td><td></td><td></td><td></td></t<>					
5 G19 9434234 4.50E-34 Lycopersicon esculentum 5 G19 7476682 1.30E-30 Glycine max 5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossyplum arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25 Ricinus communis 5 G19 9489304 6.00E-25 Hordeum vulgare 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 7720316 8.00E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 758881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7					
5 G19 7478682 1.30E-30 Glycine max 5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 7720316 8.80E-26 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 78881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 886741 1.00E-73 Zea mays 7 G261 7561318 1.20E-46 Medicago truncatula 7 G					
5 G19 6654934 1.20E-28 Medicago truncatula 5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 688579 3.60E-25 Ricinus communis 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 7720316 8.80E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 7818881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 886741 1.00E-73 Zea mays 7 G261 7561318 1.20E-47 Lycopersicon esculentum 7				4.50E-34	
5 G19 3264766 5.50E-26 Prunus armeniaca 5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 949304 6.00E-25 Ricinus communis 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 7720316 8.80E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 78881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 886741 1.00E-73 Zea mays 7 G261 7501318 1.20E-40 Medicago truncatula 7 G261 7501318 1.20E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7			7478682		
5 G19 7624302 8.30E-26 Gossypium arboreum 5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 688579 3.60E-25 Ricinus communis 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 7720316 8.80E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 758881 8.80E-86 Medicago sativa 7 G261 786881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 886741 1.00E-73 Zea mays 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G2				1.20E-28	
5 G19 9425363 2.90E-25 Triticum aestivum 5 G19 688579 3.60E-25 Ricinus communis 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 7720316 8.80E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 758881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 793914 3.50E-41 Glycine max 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 893922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9			3264766		
5 G19 688579 3.60E-25 Ricinus communis 5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 7720316 8.80E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 7158881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum			7624302		Gossypium arboreum
5 G19 9419304 6.00E-25 Hordeum vulgare 5 G19 7720316 8.80E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 7158881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 79491 1.70E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 89251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 7628776 1.20E-57 Medicago truncatula <			9425363	2.90E-25	
5 G19 7720316 8.80E-25 Lotus japonicus 7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 7158881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 19491 1.70E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum <		G19	688579	3.60E-25	Ricinus communis
7 G261 5821137 5.10E-93 Nicotiana tabacum 7 G261 7158881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 19491 1.70E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 8903922 4.00E-39 Hordeum vulgare 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 764776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum			9419304	6.00E-25	
7 G261 7158881 8.80E-86 Medicago sativa 7 G261 886741 1.00E-73 Zea mays 7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 19491 1.70E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 8903922 4.00E-39 Hordeum vulgare 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 6566281 8.40E-47 Glycine max		G19	7720316	8.80E-25	Lotus japonicus
7 G261 886741 1.00E-73 Zea mays 7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 19491 1.70E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 89251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum		G261	5821137	5.10E-93	
7 G261 5900449 5.20E-47 Lycopersicon esculentum 7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 19491 1.70E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 7627061 8.40E-47 Glycine max	7	G261	7158881	8.80E-86	
7 G261 7561318 1.20E-46 Medicago truncatula 7 G261 19491 1.70E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 7627061 8.40E-47 Glycine max 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla	7	G261	886741	1.00E-73	Zea mays
7 G261 19491 1.70E-42 Lycopersicon peruvianum 7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla	7	G261	5900449	5.20E-47	Lycopersicon esculentum
7 G261 7233914 3.50E-41 Glycine max 7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa	7	G261	7561318	1.20E-46	Medicago truncatula
7 G261 4528238 9.00E-41 Citrus unshiu 7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum <td>7</td> <td>G261</td> <td>19491</td> <td>1.70E-42</td> <td>Lycopersicon peruvianum</td>	7	G261	19491	1.70E-42	Lycopersicon peruvianum
7 G261 8903922 4.00E-39 Hordeum vulgare 7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum	7	G261	7233914	3.50E-41	Glycine max
7 G261 9251913 1.90E-36 Solanum tuberosum 9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossyplum arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum	7	G261	4528238	9.00E-41	Citrus unshiu
9 G28 7528275 4.20E-62 Mesembryanthemum crystallinum 9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum	7	G261	8903922	4.00E-39	Hordeum vulgare
9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum	7	G261	9251913	1.90E-36	Solanum tuberosum
9 G28 6654776 1.20E-57 Medicago truncatula 9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum		G28	7528275	4.20E-62	Mesembryanthemum crystallinum
9 G28 790362 2.30E-54 Nicotiana tabacum 9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum	9				
9 G28 8809570 8.00E-54 Nicotiana sylvestris 9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum	9	G28			
9 G28 3342210 8.40E-54 Lycopersicon esculentum 9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum					Nicotiana sylvestris
9 G28 6566281 8.40E-47 Glycine max 9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum	9	G28	3342210	8.40E-54	Lycopersicon esculentum
9 G28 7627061 8.40E-47 Gossypium arboreum 9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum	9				
9 G28 7324479 2.00E-44 Lycopersicon pennellii 9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum					
9 G28 6478844 1.80E-35 Matricaria chamomilla 9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum					
9 G28 7273972 7.80E-29 Oryza sativa 11 G869 2213784 1.30E-19 Lycopersicon esculentum					<u> </u>
11 G869 2213784 1.30E-19 Lycopersicon esculentum					
					

Figure 3B

SEQ ID No.	GID	Genbank NID	P-value	Species
11	G869	8570080	4.20E-18	Oryza sativa
11	G869	7560260	1.50E-17	Medicago truncatula
11	G869	7534890	5.20E-14	Śorghum bicolor
11	G869	6455322	1.10E-13	Glycine max
11	G869	9362061	2.70E-13	Triticum aestivum
11	G869	7788764	5.70E-13	Lotus japonicus
11	G869	7624302	2.50E-12	Gossypium arboreum
11	G869	3858036	2.80E-12	Populus balsamifera subsp. trichocarpa
13	G237	8283916	4.70E-42	Glycine max
13	G237	9361969	8.30E-41	Triticum aestivum
13	G237	4753385	4.10E-39	Zea mays
13	G237	7535969	4.10E-33	Sorghum bicolor
13	G237	7566043	9.30E-33	Medicago truncatula
13	G237	7339127	2.00E-32	Lycopersicon esculentum
13	G237	5860031	1.10E-28	Pinus taeda
13	G237	7776223	2.20E-28	Lotus japonicus
13	G237	6850206	5.10E-28	Oryza sativa
13	G237	5048991	8.50E-28	Gossypium hirsutum
15	G409	6654773	6.10E-57	Medicago truncatula
15	G409	6531235	2.00E-56	Lycopersicon esculentum
15	G409	7924152	1.10E-47	Glycine max
15	G409	5006854	6.50E-43.	Oryza sativa
15	G409	8098529	2.10E-41	Hordeum vulgare
15	G409	767697	1.40E-37	Daucus carota
15	G409	8328991	3.30E-37	Mesembryanthemum crystallinum
15	G409	7415613	1.40E-32	Physcomitrella patens
15	G409	7785121	2.80E-32	Lotus japonicus
15	G409	6916941	4.80E-32	Lycopersicon pennellii
17	G418	7239156	1.90E-123	Malus x domestica
17	G418	5892190	2.00E-62	Lycopersicon esculentum
17	G418	7628137	8.70E-58	Gossypium arboreum
17	G418	9205496	3.90E-51	Glycine max
17	G418	6069643	1.50E-45	Oryza sativa
17	G418	7562931	6.90E-45	Medicago truncatula
17	G418	7781695	5.50E-40	Lotus japonicus
17	G418	9298824	7.80E-34	Sorghum bicolor
17	G418	9428023	3.90E-32	Triticum aestivum
17	G418	7244366	1.30E-31	Mentha x piperita
19	G591	7646333	1.90E-55	Lycopersicon esculentum
19	G591	7924288	4.10E-53	Glycine max
19	G591	7722838	1.10E-41	Lotus japonicus
19	G591	5804781	1.40E-24	Nicotiana tabacum
19	G591	9198126	2.50E-23	Medicago truncatula
19	G591	427677	9.50E-15	Oryza sativa
19	G591	7624745	1.80E-14	Gossypium arboreum
19	G591	7535578	8.70E-14	Sorghum bicolor
19	G591	5915205	1.30E-11	Zea mays
19	G591	9249806	2.60E-11	Solanum tuberosum
21	G525	4384535	5.60E-61	Lycopersicon esculentum
21	G525	6454868	2.00E-58	Glycine max
21	G525	6066594	9.30E-54	Petunia x hybrida
21	G525	4977542	8.60E-51	Oryza sativa
41	3323	7311342	U.UUL-U I	101720 00010

Figure 3C

SEQ ID No. GID Genbank NID P-value Species 21 G525 9361647 2.50E-50 Triticum aestivum 21 G525 4218536 5.20E-50 Triticum sp. 21 G525 6732159 5.20E-50 Triticum monococcum 21 G525 5343151 2.70E-49 Zea mays 21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 7206360 3.10E-44 Medicago struncatula 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7266808 9.60E-40 Gossypium arboreum 23 G545 438492 3.90E-39 Petunia x hybrida <th></th>	
21 G525 4218536 5.20E-50 Triticum sp. 21 G525 6732159 5.20E-50 Triticum monococcum 21 G525 5343151 2.70E-49 Zea mays 21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula <	
21 G525 6732159 5.20E-50 Triticum monococcum 21 G525 5343151 2.70E-49 Zea mays 21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus	
21 G525 5343151 2.70E-49 Zea mays 21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 7728328 3.70E-52 Medicago sativa 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 438492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus	
21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 438492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 4382658 1.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 7206394 4.90E-29 Medicago tr	
21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 4382658 1.70E-38 Lycopersicon hirsutum 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine m	
23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 7788764 4.10E-23 Lotus japonicus	
23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 7796858 5.70E-27 Glycine max 25 G865 7788764 4.10E-23 Lotus japonicus <td></td>	
23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 7796858 5.70E-27 Glycine max 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 7788764 4.10E-23 Lotus japonicus <td></td>	
23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanth	
23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum cry	
23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniac	
23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
23 G545 7322653 6.80E-37 Lycopersicon hirsutum 23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
23 G545 7785845 1.10E-33 Lotus japonicus 25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 9417297 1.70E-32 Triticum aestivum 25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 7206394 4.90E-29 Medicago truncatula 25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 7796858 5.70E-27 Glycine max 25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 4387560 9.20E-25 Lycopersicon esculentum 25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 569065 1.50E-23 Oryza sativa 25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 7788764 4.10E-23 Lotus japonicus 25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 790362 8.40E-22 Nicotiana tabacum 25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 7528275 5.90E-21 Mesembryanthemum crysta 25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 3264766 8.80E-20 Prunus armeniaca	allinum
25 G865 8098026 2.00E-19 Hordeum vulgare 27 G881 5820418 9.80E-29 Glycine max	
27 G881 8440065 1.00E-27 Gossypium hirsutum 27 G881 4380578 1.50E-27 Lycopersicon esculentum	
27 G881 9199620 2.70E-27 Medicago truncatula	
27 G881 6472584 2.20E-24 Nicôtiana tabacum	
27 G881 9250698 3.20E-24 Solanum tuberosum	
27 G881 8205146 5.20E-21 Oryza sativa	
27 G881 1159878 8.20E-17 Avena fatua	
27 G881 9299778 2.70E-16 Sorghum bicolor	
27 G881 9444636 1.10E-14 Triticum aestivum	
29 G896 9410462 1.90E-101 Hordeum vulgare	
29 G896 7628908 3.60E-82 Gossypium arboreum	
29 G896 7244408 1.80E-79 Mentha x piperita	
29 G896 5046180 2.10E-73 Gossypium hirsutum	
29 G896 7678652 1.10E-63 Lotus japonicus	
29 G896 8286031 1.40E-60 Glycine max	
29 G896 5888938 4.50E-58 Lycopersicon esculentum	
29 G896 9298238 9.20E-54 Sorghum bicolor	
29 G896 7566414 8.00E-52 Medicago truncatula	
29 G896 8845076 1.00E-46 Triticum aestivum	
31 G378 5270028 5.10E-73 Lycopersicon esculentum	
31 G378 5048335 4.10E-58 Gossypium hirsutum	
31 G378 7239521 5.90E-42 Oryza sativa	
31 G378 5606120 6.80E-36 Glycine max	
31 G378 3853800 3.20E-30 Populus tremula x Populus	tremuloides
31 G378 7659983 1.70E-23 Sorghum bicolor	

Figure 3D

SEQ ID No.	GID	Genbank NID	P-value	Species
31	G378	6626305	1.10E-21	Zea mays
31	G378	9412941	9.40E-19	Triticum aestivum
31	G378	3242033	4.30E-17	Mesembryanthemum crystallinum
31	G378	7626259	7.70E-13	Gossypium arboreum
33	G229	7337390	6.60E-51	Lycopersicon esculentum
33	G229	9823237	3.60E-50	Hordeum vulgare
33	G229	7244424	4.90E-50	Mentha x piperita
33	G229	7776053	1.70E-49	Lotus japonicus
33	G229	2921335	5.80E-48	Gossypium hirsutum
33	G229	1491932	4.50E-47	Zea mays
33	G229	6455590	2.80E-44	Glycine max
33	G229	6020191	2.00E-41	Pinus taeda
33	G229	10697236	4.20E-41	Oryza sativa
33	G229	7765706	5.10E-41	Medicago truncatula
35	G663	7673087	5.10E-43	Petunia integrifolia
35	G663	9508051	3.00E-41	Lycopersicon esculentum
35	G663	7673091	3.30E-41	Petunia x hybrida
35	G663	7673097	2.40E-36	Petunia axillaris
35	G663	5048991	1.20E-33	Gossypium hirsutum
35	G663	6455590	2.50E-31	Glycine max
35	G663	7560175	1.90E-27	Medicago truncatula
35	G663	7244424	4.10E-26	Mentha x piperita
35	G663	9954117	3.40E-25	Solanum tuberosum
35	G663	6020191	3.60E-25	Pinus taeda
37	G1396	498704	5.20E-22	Spinacia oleracea
37	G1396	7502400	1.20E-21	Gossypium arboreum
37	G1396	3857536	3.40E-21	Populus balsamifera subsp. trichocarpa
37	G1396	4385300	1.20E-20	Lycopersicon esculentum
37	G1396	6917249	1.50E-20	Lycopersicon pennellii
37	G1396	6915979	1.70E-20	Glycine max
37	G1396	7674530	2.70E-20	Medicago truncatula
37	G1396	8090319	3.40E-20	Sorghum bicolor
37	G1396	3592182		Oryza sativa
37	G1396	6654124	9.10E-20 1.10E-19	Zea mays
39	G265	5821137	6.50E-83	Nicotiana tabacum
39	G265	7158881	3.80E-79	Medicago sativa
39	G265	886741	1.60E-70	Zea mays
39	G265		5.60E-43	Lycopersicon esculentum
		5900449		
39 39	G265 G265	8903922 7561318	8.20E-43 2.10E-41	Hordeum vulgare Medicago truncatula
39	G265	9204445	5.30E-36	Glycine max
. 39			5.40E-36	Citrus unshiu
39	G265	4528238	2.10E-35	
39	G265	19489	2.10E-35 2.00E-32	Lycopersicon peruvianum Solanum tuberosum
41	G265 G1006	9251913 7528275		Mesembryanthemum crystallinum
			2.70E-51 4.90E-49	Lycopersicon esculentum
41	G1006	3342210 6654776		Medicago truncatula
41	G1006	6654776	1.90E-48	Nicotiana tabacum
41	G1006	790362	2.30E-47	
41	G1006	8809570	2.00E-46	Nicotiana sylvestris
41	G1006	7627061	6.40E-41	Gossypium arboreum
41	G1006	7324479	1.20E-35	Lycopersicon pennellii
41	G1006	6478844	1.80E-35	Matricaria chamomilla

Figure 3E

SEQ ID No.	GID	Genbank NID	P-value	Species
41	G1006	6566281	1.30E-34	Glycine max
41	G1006	4716624	3.80E-28	Oryza sativa
43	G1309	9361969	2.40E-45	Triticum aestivum
43	G1309	7566043	9.60E-35	Medicago truncatula
43	G1309	5891104	2.20E-31	Lycopersicon esculentum
43	G1309	5860031	2.10E-30	Pinus taeda
43	G1309	5049507	6.20E-30	Gossypium hirsutum
43	G1309	5139805	1.30E-29	Glycine max
43	G1309	6850206	2.50E-29	Oryža sativa
43	G1309	7721017	3.40E-29	Lotus japonicus
43	G1309	8368245	5.20E-28	Zea mays
43	G1309	20560	9.50E-27	Petunia x hybrida
45	G2550	4380729	2.80E-51	Lycopersicon esculentum
45	G2550	5667196	2.20E-49	Oryza sativa
45	G2550	8669454	1.40E-48	Glycine max
45	G2550	9298824	1.50E-48	Sorghum bicolor
45	G2550	7239156	9.90E-46	Malus x domestica
45	G2550	7570704	5.70E-45	Medicago truncatula
45	G2550	7628137	3.30E-42	Gossypium arboreum
45	G2550	7244366	6.00E-41	Mentha x piperita
45	G2550	9428023	4.70E-40	Triticum aestivum
45	G2550	9250642	3.50E-39	Solanum tuberosum
47	G965	7239156	3.10E-126	Malus x domestica
47	G965	5892190	2.00E-62	Lycopersicon esculentum
47	G965	7628137	1.60E-56	Gossypium arboreum
47	G965	9205496	2.60E-49	Glycine max
47	G965	6069643	1.70E-45	Oryza sativa
47	G965	7562931	2.50E-44	Medicago truncatula
47	G965	7781695	1.60E-41	Lotus japonicus
47	G965	9298824	6.30E-33	Sorghum bicolor
47	G965	9428023	1.50E-31	Triticum aestivum
47	G965	7244366	1.20E-29	Mentha x piperita
49	G793	6976712	3.60E-43	Lycopersicon esculentum
49	G793	7924288	2.00E-41	Glycine max
49	G793	7614163	3.90E-34	Lotus japonicus
49	G793	9198126	5.70E-23	Medicago truncatula
49	G793	5804781	1.10E-22	Nicotiana tabacum
49	G793	7535578	1.60E-14	Sorghum bicolor
49	G793	427677	6.10E-14	Oryza sativa
49 49	G793	5915205	2.90E-10	Zea mays
49	G793 G793	9249806	4.20E-10	Solanum tuberosum Gossypium arboreum
51	G764	7624745 4384535	1.30E-09 7.00E-70	Lycopersicon esculentum
51	G764	5049217	1.80E-65	Gossypium hirsutum
51	G764	6454868		Glycine max
51	G764 G764	6066594	1.90E-64 5.20E-59	Petunia x hybrida
51	G764	4218536	2.30E-52	· · · · · · · · · · · · · · · · · · ·
51	G764 G764	6732159	2.30E-52 2.30E-52	Triticum sp. Triticum monococcum
51	G764	9361647	7.50E-52	Triticum aestivum
51	G764	4977542	4.10E-49	Oryza sativa
51	G764	6799764	4.40E-49	Medicago truncatula
51	G764	9296257	1.00E-48	Sorghum bicolor
<u> </u>	<u> </u>	JE30231	1.00L-40	Looi Sugit propior

Figure 3F

SEQ ID No.	GID	Genbank NID	P-value	Species
53	G350	439492	5.20E-53	Petunia x hybrida
53	G350	7228328	8.90E-51	Medicago sativa
53	G350	4666359	3.10E-48	Datisca glomerata
53	G350	1763062	8.30E-48	Glycine max
53	G350	7626808	9.10E-44	Gossypium arboreum
53	G350	7206360	2.20E-43	Medicago truncatula
53	G350	2981168	2.10E-38	Nicotiana tabacum
53	G350	7322653	2.00E-37	Lycopersicon hirsutum
53	G350	5276755	2.40E-37	Lycopersicon esculentum
53	G350	2058503	1.10E-31	Brassica rapa
55	G986	6472584	1.00E-34	Nicotiana tabacum
55	G986	8440065	8.80E-33	Gossypium hirsutum
55	G986	4385167	1.50E-32	Lycopersicon esculentum
55	G986	8205146	5.50E-30	Oryza sativa
55	G986	5820418	8.80E-26	Glycine max
55	G986	1159878	2.30E-23	Avena fatua
55	G986	9250698	4.60E-22	Solanum tuberosum
55	G986	9413507	7.90E-21	Triticum aestivum
55	G986	7748539	2.30E-20	Lotus japonicus
55	G986	9199620	1.30E-16	Medicago truncatula
57	G1349	8904043	1.50E-47	Hordeum vulgare
57	G1349	7244408	2.40E-47	Mentha x piperita
57	G1349	8286031	3.60E-46	Glycine max
57	G1349	9298238	9.10E-36	Sorghum bicolor
57	G1349	7628908	4.70E-34	Gossypium arboreum
57	G1349	5046180	1.50E-33	Gossypium hirsutum
57	G1349	5888938	1.30E-30	Lycopersicon esculentum
57	G1349	5043924	6.20E-30	Pinus taeda
57	G1349	8845076	4.40E-29	Triticum aestivum
57	G1349	7678652	4.20E-27	Lotus japonicus

MBI15 Sequence Listing.ST25 SEQUENCE LISTING

<pre><110> Heard, Jacqueline Keddie, James Yu, Guo-Liang Ratcliffe, Oliver Creelman, Robert Jiang, Cai-zhong Pineda, Omaira Reuber, Lynne Adam, Luc</pre>
<120> Pathogen Resistance Genes
<130> MBI-0015
<150> 60/166,228 <151> 1999-11-17
<150> Plant Trait Modification III <151> 2000-08-22
<150> 60/197,899 <151> 2000-04-17
<160> 58
<170> PatentIn version 3.0
<210> 1 <211> 1187 <212> DNA <213> Arabidopsis thaliana
<220> <221> CDS <222> (50)(1096) <223> G188
<pre><400> 1 ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc</pre>
ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc 58 Met Ser Ser
gag gat tgg gat ctc ttc gcc gtc gtc aga agc tgc agc tct tct gtt Glu Asp Trp Asp Leu Phe Ala Val Val Arg Ser Cys Ser Ser Ser Val
ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc Met Ser Ser 1 gag gat tgg gat ctc ttc gcc gtc gtc aga agc tgc agc tct tct gtt Glu Asp Trp Asp Leu Phe Ala Val Val Arg Ser Cys Ser Ser Ser Val 5 tcc acc acc aat tct tgt gct ggt cat gaa gac gac ata gga aac tgt Ser Thr Thr Asn Ser Cys Ala Gly His Glu Asp Asp Ile Gly Asn Cys
ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc Met Ser Ser 1 gag gat tgg gat ctc ttc gcc gtc gtc aga agc tgc agc tct tct gtt Glu Asp Trp Asp Leu Phe Ala Val Val Arg Ser Cys Ser Ser Ser Val 5 tcc acc acc act tct tgt gct ggt cat gaa gac gac ata gga aac tgt Ser Thr Thr Asn Ser Cys Ala Gly His Glu Asp Asp Ile Gly Asn Cys 20 aaa caa caa caa gat cct cct cct cct cct ctg ttt caa gct tct tct Lys Gln Gln Gln Asp Pro Pro Pro Pro Leu Phe Gln Ala Ser Ser
ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc Met Ser Ser gag gat tgg gat ctc ttc gcc gtc gtc aga agc tgc agc tct tct gtt Ser Trp Asp Leu Phe Ala Val Val Arg Ser Cys Ser Ser Ser Val tcc acc acc act tct tgt gct ggt cat gaa gac gac ata gga aac tgt Ser Thr Thr Asn Ser Cys Ala Gly His Glu Asp Asp Ile Gly Asn Cys 25 aaa caa caa caa gat cct cct cct cct cct ctg ttt caa gct tct tct Lys Gln Gln Gln Asp Pro Pro Pro Pro Pro Leu Phe Gln Ala Ser Ser 40 tct tgc aac gag tta caa gat tct tgc aaa cca ttt tta ccc gtt act Ser Cys Asn Glu Leu Gln Asp Ser Cys Lys Pro Phe Leu Pro Val Thr
ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc Met Ser Ser gag gat tgg gat ctc ttc gcc gtc gtc aga agc tgc agc tct tct gtt Glu Asp Trp Asp Leu Phe Ala Val Val Arg Ser Cys Ser Ser Ser Val tcc acc acc aat tct tgt gct ggt cat gaa gac gac ata gga aac tgt Ser Thr Thr Asn Ser Cys Ala Gly His Glu Asp Asp Ile Gly Asn Cys 20 30 35 aaa caa caa caa gat cct cct cct cct cct ctg ttt caa gct tct tct Lys Gln Gln Gln Asp Pro Pro Pro Pro Pro Pro Pro Gln Ala Ser Ser 40 202 tct tgc aac gag tta caa gat tct tgc aaa cca ttt tta ccc gtt act Ser Cys Asn Glu Leu Gln Asp Ser Cys Lys Pro Phe Leu Pro Val Thr 55 act act act act act tgg tct cct cct cct ctc ctc ctc ctc c
gag gat tgg gat ctc ttc gcc gtc gtc aga agc tgc agc tct tct gtt Glu Asp Trp Asp Leu Phe Ala Val Val Arg Ser Cys Ser Ser Ser Val 15 tcc acc acc aat tct tgt gct ggt cat gaa gac gac ata gga aac tgt Ser Thr Thr Asn Ser Cys Ala Gly His Glu Asp Asp Ile Gly Asn Cys 30 aaa caa caa caa gat cct cct cct cct cct ctg ttt caa gct tct tct Lys Gln Gln Gln Asp Pro Pro Pro Pro Pro Leu Phe Gln Ala Ser Ser Ser Cys Asn Glu Leu Gln Asp Ser Cys Lys Pro Phe Leu Pro Val Thr 65 act act act act act act tgg tct cct cct cct cct cct cta ctt cct cct

				120			MBI	15 S	eque: 125	nce	List	ing.	ST25	130		
					caa Gln											490
aag Lys	aat Asn	cag Gln 150	caa Gln	aaa Lys	aga Arg	acc Thr	ata Ile 155	tgt Cys	cat His	gta Val	acg Thr	caa Gln 160	gag Glu	aat Asn	ctt Leu	538
tct Ser	tct Ser 165	gat Asp	ttg Leu	tgg Trp	gct Ala	tgg Trp 170	cgt Arg	aaa Lys	tac Tyr	ggt Gly	caa Gln 175	aaa Lys	ccc Pro	atc Ile	aaa Lys	586
ggc Gly 180	tct Ser	cct Pro	tat Tyr	cca Pro	agg Arg 185	aat Asn	tat Tyr	tac Tyr	aga Arg	tgt Cys 190	agt Ser	agc Ser	tca Ser	aaa Lys	gga Gly 195	634
tgt Cys	tta Leu	gca Ala	cga Arg	aaa Lys 200	caa Gln	gtt Val	gaa Glu	aga Arg	agt Ser 205	aat Asn	tta Leu	gat Asp	cct Pro	aat Asn 210	atc Ile	682
ttc Phe	atc Ile	gtt Val	act Thr 215	tac Tyr	acc Thr	gga Gly	gaa Glu	cac His 220	act Thr	cat His	cca Pro	cgt Arg	cct Pro 225	act Thr	cac His	730
cgg Arg	aac Asn	tct Ser 230	ctc Leu	gcc Ala	gga Gly	agt Ser	act Thr 235	cgt Arg	aac Asn	aaa Lys	tct Ser	cag Gln 240	ccc Pro	gtt Val	aac Asn	778
ccg Pro	gtt Val 245	cct Pro	aaa Lys	ccg Pro	gac Asp	aca Thr 250	tct Ser	cct Pro	tta Leu	tcg Ser	gat Asp 255	aca Thr	gta Val	aaa Lys	gaa Glu	826
gag Glu 260	att Ile	cat His	ctt Leu	tct Ser	ccg Pro 265	acg Thr	aca Thr	ccg Pro	ttg Leu	aaa Lys 270	gga Gly	aac Asn	gat Asp	gac Asp	gtt Val 275	874
caa Gln	gaa Glu	acg Thr	aat Asn	gga Gly 280	gat Asp	gaa Glu	gat Asp	atg Met	gtt Val 285	ggt Gly	caa Gln	gaa Glu	gtc Val	aac Asn 290	atg Met	922
gaa Glu	gag Glu	gaa Glu	gag Glu 295	gag Glu	gaa Glu	gaa Glu	gaa Glu	gtg Val 300	gaa Glu	gaa Glu	gat Asp	gat Asp	gaa Glu 305	gaa Glu	gaa Glu	970
gaa Glu	gat Asp	gat Asp 310	gat Aвр	gac Asp	gtg Val	gat Asp	gat Asp 315	ctt Leu	ttg Leu	ata Ile	cca Pro	aat Asn 320	tta Leu	gcg Ala	gtg Val	1018
aga Arg	gat Asp 325	cga Arg	gat Asp	gat Asp	ttg Leu	ttc Phe 330	ttc Phe	gct Ala	gga Gly	agt Ser	ttt Phe 335	cca Pro	tct Ser	tgg Trp	tcc Ser	1066
gcc Ala 340	gga Gly	tcc Ser	gcc Ala	ggt Gly	gac Asp 345	ggt Gly	ggt Gly	gga Gly	tga	tgaa	aaac	gaa	taaa	atct	ca	1116
att	tacaa	att (taca	aaaa	ga aa	aaaa	gtcag	g tt!	ttaa	atta	tta	tttt	tgt	ttgti	caaaac	1176
ttga	acati	tta (t													1187
<210 <210 <210 <210	l > 1 2 > 1	2 348 PRT Arab:	idop	sis :	thal:	iana										
<40	0 > 2	2														
Met 1	Ser	Ser	Glu	Asp 5	Trp	Asp	Leu	Phe	Ala 10	Val	Val	Arg	Ser	Cys 15	Ser	

Ser Ser Val Ser Thr Thr Asn Ser Cys Ala Gly His Glu Asp Asp Ile Page 2 $\,$

MBI15 Sequence Listing.ST25 25 30

20

Gly Asn Cys Lys Gln Gln Gln Asp Pro Pro Pro Pro Pro Leu Phe Gln 35 40

Ala Ser Ser Cys Asn Glu Leu Gln Asp Ser Cys Lys Pro Phe Leu
50 55 60

Pro Val Thr Thr Thr Thr Thr Thr Thr Trp Ser Pro Pro Pro Leu Leu 65 70 75 80

Pro Pro Pro Lys Ala Ser Ser Pro Ser Pro Asn Ile Leu Leu Lys Gln 85 90 95

Glu Gln Val Leu Leu Glu Ser Gln Asp Gln Lys Pro Pro Leu Ser Val

Arg Val Phe Pro Pro Ser Thr Ser Ser Ser Val Phe Val Phe Arg Gly

Gln Arg Asp Gln Leu Leu Gln Gln Gln Ser Gln Pro Pro Leu Arg Ser 130 135 140

Arg Lys Arg Lys Asn Gln Gln Lys Arg Thr Ile Cys His Val Thr Gln 145 150 150 160

Glu Asn Leu Ser Ser Asp Leu Trp Ala Trp Arg Lys Tyr Gly Gln Lys 165 170 175

Pro Ile Lys Gly Ser Pro Tyr Pro Arg Asn Tyr Tyr Arg Cys Ser Ser 180 185 190

Ser Lys Gly Cys Leu Ala Arg Lys Gln Val Glu Arg Ser Asn Leu Asp

Pro Asn Ile Phe Ile Val Thr Tyr Thr Gly Glu His Thr His Pro Arg 210 215 220

Pro Thr His Arg Asn Ser Leu Ala Gly Ser Thr Arg Asn Lys Ser Gln 225 230 240

Pro Val Asn Pro Val Pro Lys Pro Asp Thr Ser Pro Leu Ser Asp Thr 245 250 255

Val Lys Glu Glu Ile His Leu Ser Pro Thr Thr Pro Leu Lys Gly Asn 260 265 270

Asp Asp Val Glu Glu Thr Asn Gly Asp Glu Asp Met Val Gly Glu Glu 275 280 285

Glu Glu Glu Glu Asp Asp Asp Val Asp Asp Leu Leu Ile Pro Asn 305 310 315 320

MBI15 Sequence Listing.ST25
Leu Ala Val Arg Asp Arg Asp Leu Phe Phe Ala Gly Ser Phe Pro
325 330 335

Ser Trp Ser Ala Gly Ser Ala Gly Asp Gly Gly Gly 340

<210><211><211><212><213>	3 1431 DNA Arabido	psis tha	liana							
<220> <221> <222> <223>	CDS (129) G616	(1211)								
<400> ctttaa	3 atcc caa	accaacc	ctaaagtt	tt gat	ttttaat	tttgggg	ıgta a	accaa	ıaaaaa	60
aaacaa	acc cta	attttt	ttctttag	tg atg	agattat	tggtgat	gat ç	gaaat	gattg	120
gagatc	ta atg a Met L 1	ag aat a ys Asn A	ac aac a sn Asn A 5	at ggc sn Gly	gac gt Asp Va	t gtg ga 1 Val As 10	at aad ap Asr	gaa Glu	gtg Val	170
aac aa Asn As: 15	c cgg tt n Arg Le	a agc cg u Ser Ar 20	g tgg ca g Trp Hi	t cac s His	aat tct Asn Ser 25	tcc cgg Ser Arg	g ata g Ile	att Ile	agg Arg 30	218
gtt tc Val Se	a cga gc r Arg Al	t tcc gg a Ser Gl 35	t ggt aa y Gly Ly	s Asp	cga cac Arg His 40	agc aaa Ser Lys	gtc Val	ttg Leu 45	act Thr	266
tct aa Ser Ly	a gga cc s Gly Pr 50	a cgt ga o Arg As	c cgt cg p Arg Ar	t gtc g Val 55	cgg tta Arg Leu	tca gto Ser Val	s tcc Ser 60	acc Thr	gct Ala	314
ctt ca Leu Gl	a ttc ta n Phe Ty 65	t gat ct r Asp Le	t caa ga u Gln As 70	p Arg	tta ggt Leu Gly	tat gat Tyr Ası 75	caa Gln	cct Pro	agc Ser	362
aaa gc Lys Al 80	t gtt ga a Val Gl	a tgg tt u Trp Le	a atc aa u Ile Ly 85	a gct s Ala	gct gaa Ala Glu	gat to Asp Ser 90	atc Ile	tct. Ser	gag Glu	410
ctt cc Leu Pr 95	t tca ct o Ser Le	c aac aa u Asn As 10	c act ca n Thr Hi 0	t ttt s Phe	ccg acc Pro Thr 105	Asp As	gag Glu	aat Asn	cac His 110	458
cag aa Gln As	t cag ac n Gln Th	a tta ac r Leu Th 115	a aca gt r Thr Va	l Ala	gct aat Ala Asn 120	tcc tto Ser Le	g tct i Ser	aaa Lys 125	tct Ser	506
gct tg Ala Cy	t agt ag s Ser Se 13	r Asn Se	a gac ac r Asp Th	g agc r Ser 135	aag aac Lys Asn	tct tc	ggt Gly 140	ttg Leu	tct Ser	554
tta tc Leu Se	a aga tc r Arg Se 145	g gag ct r Glu Le	t aga ga u Arg As 15	p Lys	gct aga Ala Arg	gag agg Glu Arg 15	g Ala	aga Arg	gag Glu	602
aga ac Arg Th 16	r Ala Ly	a gag ac s Glu Th	c aag ga r Lys Gl 165	a aga u Arg	gat cat Asp His	aac ca Asn Hi 170	c act Thr	tcg Ser	ttt Phe	650
acg ga Thr As 175	t ttg tt p Leu Le	a aat to u Asn Se 18	c ggt tc r Gly Se 0	a gat r Asp	ccg gtt Pro Val 185	Asn Se	a aac r Asn	cgg Arg	caa Gln 190	698
tgg at Trp Me	g gct tc t Ala Se	a gct co r Ala Pr 195	t tot to o Ser Se	a tct r Ser	cca atg Pro Met 200	gag ta Glu Ty	t ttc r Phe	agt Ser 205	tcg Ser	746

MOTIE	Sequence	Ticting	CTOE
MHILS	Seguence	Listing	. ST25

							MBI	15 S	eque	nce :	List	ing.	ST25			
ggt Gly	tta Leu	att Ile	ctc Leu 210	999 Gly	tcg Ser	ggt Gly	caa Gln	caa Gln 215	acc Thr	cat His	ttc Phe	cct Pro	att Ile 220	tca Ser	aca Thr	794
aat Asn	tct Ser	cat His 225	cct Pro	ttc Phe	tca Ser	tca Ser	atc Ile 230	tcc Ser	gat Asp	cat His	cat His	cat His 235	cat His	cat His	cct Pro	842
cat His	cat His 240	cag Gln	cat His	caa Gln	gag Glu	ttt Phe 245	tca Ser	ttc Phe	gtt Val	ccc Pro	gac Asp 250	cat His	ttg Leu	ata Ile	tca Ser	890
ccg Pro 255	gca Ala	gaa Glu	tcc Ser	aac Asn	ggc Gly 260	gga Gly	gca Ala	ttc Phe	aat Asn	ctt Leu 265	gat Asp	ttt Phe	aat Asn	atg Met	tca Ser 270	938
aca Thr	ccc Pro	tcc Ser	ggc Gly	gcc Ala 275	gga Gly	gct Ala	gcc Ala	gtc Val	tcc Ser 280	gcc Ala	gca Ala	tca Ser	ggt Gly	ggt Gly 285	ggc Gly	986
ttc Phe	agt Ser	ggt Gly	ttc Phe 290	aac Asn	agg Arg	ggg Gly	acc Thr	ctt Leu 295	cag Gln	tcc Ser	aat Asn	tca Ser	aca Thr 300	aat Asn	cag Gln	1034
cat His	cag Gln	tca Ser 305	ttc Phe	ctc Leu	gct Ala	aat Asn	cta Leu 310	cag Gln	agg Arg	ttt Phe	cca Pro	aca Thr 315	tca Ser	gaa Glu	agt Ser	1082
Gly	gga Gly 320	ggt Gly	cca Pro	cag Gln	ttc Phe	tta Leu 325	ttc Phe	ggt Gly	gca Ala	ctg Leu	cct Pro 330	gca Ala	gag Glu	aat Asn	cac His	1130
cac His 335	cac His	aat Asn	cac His	cag Gln	ttt Phe 340	cag Gln	ctt Leu	tac Tyr	tat Tyr	gaa Glu 345	aat Asn	gga Gly	tgc Cys	aga Arg	aac Asn 350	1178
tca Ser	tca Ser	gaa Glu	cat His	aag Lys 355	ggt Gly	aaa Lys	ggc Gly	aag Lys	aac Asn 360	tga	tga	tatt	aat '	tatt	gcatct	1231
ttg	gttt	tgt 1	tcaa	atgc	tc a	tttt	gtat	g tt	tatc	tttg	gtt	tatt	tca a	aaac	aaatgt	1291
taa	tctc	ttt	cgtt	gtct	ga t	gtgt	gtta	9 991	tttt	gttt	tat	gtat	tga (gggt	ctttgg	1351
aaa	tctt	ttt	gcat	tgtg	ct t	gtaa	tgtt	g ta	tttg	tgat	aat	agca	ttt	tgtt	tgtgag	1411
tta	aaaa	aaa a	aaaa	aaaa	aa											1431
<21 <21 <21 <21	1 > 2 >	4 360 PRT Arab	idop	sis	thal	iana										

<400> 4

Met Lys Asn Asn Asn Asn Gly Asp Val Val Asp Asn Glu Val Asn Asn 1 5 10 10

Arg Leu Ser Arg Trp His His Asn Ser Ser Arg Ile Ile Arg Val Ser 25 30

Arg Ala Ser Gly Gly Lys Asp Arg His Ser Lys Val Leu Thr Ser Lys 35 40 45

Gly Pro Arg Asp Arg Arg Val Arg Leu Ser Val Ser Thr Ala Leu Gln 50 55

Phe Tyr Asp Leu Gln Asp Arg Leu Gly Tyr Asp Gln Pro Ser Lys Ala 65 70 75

MBI15 Sequence Listing.ST25

Val Glu Trp Leu Ile Lys Ala Ala Glu Asp Ser Ile Ser Glu Leu Pro

Ser Leu Asn Asn Thr His Phe Pro Thr Asp Asp Glu Asn His Gln Asn 100 105 110

Gln Thr Leu Thr Thr Val Ala Ala Asn Ser Leu Ser Lys Ser Ala Cys 115 120 125

Ser Ser Asn Ser Asp Thr Ser Lys Asn Ser Ser Gly Leu Ser Leu Ser 130 135 140

Arg Ser Glu Leu Arg Asp Lys Ala Arg Glu Arg Ala Arg Glu Arg Thr 145 150 160

Ala Lys Glu Thr Lys Glu Arg Asp His Asn His Thr Ser Phe Thr Asp 165 170 175

Leu Leu Asn Ser Gly Ser Asp Pro Val Asn Ser Asn Arg Gln Trp Met
180 195 190

Ala Ser Ala Pro Ser Ser Ser Pro Met Glu Tyr Phe Ser Ser Gly Leu 195 200 205

Ile Leu Gly Ser Gly Gln Gln Thr His Phe Pro Ile Ser Thr Asn Ser 210 215 220

His Pro Phe Ser Ser Ile Ser Asp His His His His His Pro His His 225 230 240

Gln His Gln Glu Phe Ser Phe Val Pro Asp His Leu Ile Ser Pro Ala 245 250 255

Glu Ser Asn Gly Gly Ala Phe Asn Leu Asp Phe Asn Met Ser Thr Pro 260 265 270

Ser Gly Ala Gly Ala Ala Val Ser Ala Ala Ser Gly Gly Phe Ser 275 280 285

Gly Phe Asn Arg Gly Thr Leu Gln Ser Asn Ser Thr Asn Gln His Gln 290 295 300

Ser Phe Leu Ala Asn Leu Gln Arg Phe Pro Thr Ser Glu Ser Gly Gly 305 310 315 320

Gly Pro Gln Phe Leu Phe Gly Ala Leu Pro Ala Glu Asn His His His 325 330 335

Asn His Gln Phe Gln Leu Tyr Tyr Glu Asn Gly Cys Arg Asn Ser Ser 340 345 350

Glu His Lys Gly Lys Gly Lys Asn 355 360

<210> 5

MBI15 Sequence Listing.ST25

MBI15 Sequence Listing.ST25	
<211> 1055 <212> DNA <213> Arabidopsis thaliana	
<220> <221> CDS <222> (70)(816) <223> G19	
<400> 5 ataaaggcat ttcagctcca ccgtaggaaa ctttctcttg aaagaaaccc acagcaacaa	60
acagagaaa atg tgt ggc ggt gct att att tcc gat tat gcc cct ctc gtc Met Cys Gly Gly Ala Ile Ile Ser Asp Tyr Ala Pro Leu Val 1 5 10	111
acc aag gcc aag ggc cgt aaa ctc acg gct gag gaa ctc tgg tca gag Thr Lys Ala Lys Gly Arg Lys Leu Thr Ala Glu Glu Leu Trp Ser Glu 15 20 25 30	159
ctc gat gct tcc gcc gcc gac gac ttc tgg ggt ttc tat tcc acc tcc Leu Asp Ala Ser Ala Ala Asp Asp Phe Trp Gly Phe Tyr Ser Thr Ser 35 40 45	207
aaa ctc cat ccc acc aac caa gtt aac gtg aaa gag gag gca gtg aag Lys Leu His Pro Thr Asn Gln Val Asn Val Lys Glu Glu Ala Val Lys 50 55 60	255
aag gag cag gca aca gag ccg ggg aaa cgg agg a	303
tat aga ggg ata cgt aag cgt cca tgg gga aaa tgg gcg gct gag att Tyr Arg Gly Ile Arg Lys Arg Pro Trp Gly Lys Trp Ala Ala Glu Ile 80 85 90	351
cga gat cca cga aaa ggt gtt aga gtt tgg ctt ggt acg ttc aac acg Arg Asp Pro Arg Lys Gly Val Arg Val Trp Leu Gly Thr Phe Asn Thr 95 100 105 110	399
gcg gag gaa gct gcc atg gct tat gat gtt gcg gcc aag cag atc cgt Ala Glu Glu Ala Ala Met Ala Tyr Asp Val Ala Ala Lys Gln Ile Arg 115 120 125	447
ggt gat aaa gcc aag ctc aac ttc cca gat ctg cac cat cct cct Gly Asp Lys Ala Lys Leu Asn Phe Pro Asp Leu His His Pro Pro Pro 130 135 140	495
cct aat tat act cct ccg ccg tca tcg cca cga tca acc gat cag cct Pro Asn Tyr Thr Pro Pro Pro Ser Ser Pro Arg Ser Thr Asp Gln Pro 145 150 155	543
ccg gcg aag aag gtc tgc gtt gtc tct cag agt gag agc gag tta agt Pro Ala Lys Lys Val Cys Val Val Ser Gln Ser Glu Ser Glu Leu Ser 160 165 170	591
cag ccg agt ttc ccg gtg gag tgt ata gga ttt gga aat ggg gac gag Gln Pro Ser Phe Pro Val Glu Cys Ile Gly Phe Gly Asn Gly Asp Glu 175 180 185 190	639
ttt cag aac ctg agt tac gga ttt gag ccg gat tat gat ctg aaa cag Phe Gln Asn Leu Ser Tyr Gly Phe Glu Pro Asp Tyr Asp Leu Lys Gln 195 200 205	687
cag ata tcg agc ttg gaa tcg ttc ctt gag ctg gac ggt aac acg gcg Gln Ile Ser Ser Leu Glu Ser Phe Leu Glu Leu Asp Gly Asn Thr Ala 210 215 220	735
gag caa ccg agt cag ctt gat gag tcc gtt tcc gag gtg gat atg tgg Glu Gln Pro Ser Gln Leu Asp Glu Ser Val Ser Glu Val Asp Met Trp 225 230 235	783
atg ctt gat gat gtc att gcg tcg tat gag taa aagaaaaaaa ataagtttaa Met Leu Asp Asp Val Ile Ala Ser Tyr Glu Page 7	836

	MBI15	Sequence	Listing	.ST25
L.				

240 245 aaaaagttaa ataaagtctg taatatatat gtaaccgccg ttacttttaa aaggttttta 956 ccgtcgcatt ggactgctga tgatgtctgt tgtgtaatgt gtagaatgtg accaaatgga cgttatatta cggtttgtgg tattattagt ttcttagatg gaaaaactta catgtgtaaa 1016 taagatttgt aatgtaagac gaagtactta taacttctt 1055 <210>

<211> 248 <212> PRT

<213> Arabidopsis thaliana

Met Cys Gly Gly Ala Ile Ile Ser Asp Tyr Ala Pro Leu Val Thr Lys
1 10 15

Ala Lys Gly Arg Lys Leu Thr Ala Glu Glu Leu Trp Ser Glu Leu Asp 20 25 30

Ala Ser Ala Ala Asp Asp Phe Trp Gly Phe Tyr Ser Thr Ser Lys Leu 35 40 45

His Pro Thr Asn Gln Val Asn Val Lys Glu Glu Ala Val Lys Lys Glu 50 60

Gln Ala Thr Glu Pro Gly Lys Arg Arg Lys Arg Lys Asn Val Tyr Arg 65 70 75 80

Gly Ile Arg Lys Arg Pro Trp Gly Lys Trp Ala Ala Glu Ile Arg Asp 85 90 95

Pro Arg Lys Gly Val Arg Val Trp Leu Gly Thr Phe Asn Thr Ala Glu

Glu Ala Ala Met Ala Tyr Asp Val Ala Ala Lys Gln Ile Arg Gly Asp 115 120 125

Lys Ala Lys Leu Asn Phe Pro Asp Leu His His Pro Pro Pro Pro Asn 130 135 140

Tyr Thr Pro Pro Pro Ser Ser Pro Arg Ser Thr Asp Gln Pro Pro Ala

Lys Lys Val Cys Val Val Ser Gln Ser Glu Ser Glu Leu Ser Gln Pro

Ser Phe Pro Val Glu Cys Ile Gly Phe Gly Asn Gly Asp Glu Phe Gln
180 185 190

Asn Leu Ser Tyr Gly Phe Glu Pro Asp Tyr Asp Leu Lys Gln Gln Ile 195 200 205

Ser Ser Leu Glu Ser Phe Leu Glu Leu Asp Gly Asn Thr Ala Glu Gln 210 215 220

Pro Ser Gln Leu Asp Glu Ser Val Ser Glu Val Asp Met Trp Met Leu Page 8

240

MBI15 Sequence Listing.ST25
225 230 235

Asp Asp Val Ile Ala Ser Tyr Glu 245

<210>

1857 <211> 185' <212> DNA Arabidopsis thaliana <220> <221> CDS (458) .. (1663) <222> G261 <223> <400> 7 gtttaggttc gagaagcaga gagggttcga gaagctaata agggtttctt ctttttgatt 60 ttaatgctaa aagggttcta gattcgttga attttacaag ggttttaggg gttcttagaa 120 gettttgett gattgtettt tatttagaaa eagtggtgag tttttagtet tteaetttgt 180 240 tcaagttcga agctttttt ggagggaatt ttgggcttct gattttgatc gaaacttact 300 gatagtaagt tetttgagte eteettaaet gtagtttetg tgtaetgaag ttattgaatt 360 gaaagttttt atcttttttg gttattgaaa ctttcatagt ttgatcaaaa gagtctcttg ctctgttttt ggctctgttt ttgtgagtgt gattgtaagc tttgttgtga gtagattgaa 420 tcaaggagtg tgagagttgt taaaagtgtt ttcagag atg gat gag aat aat cat Met Asp Glu Asn Asn His 475 gga gtt tca tca agc tca ctt cca cct ttc ctc acc aaa aca tat gag Gly Val Ser Ser Ser Leu Pro Pro Phe Leu Thr Lys Thr Tyr Glu 523 atg gtt gat gat tct tca tcc gat tct atc gtc tct tgg agt cag agc Met Val Asp Asp Ser Ser Ser Asp Ser Ile Val Ser Trp Ser Gln Ser 571 aat aag agt ttc atc gtt tgg aat ccg ccg gag ttt tct aga gat ctt Asn Lys Ser Phe Ile Val Trp Asn Pro Pro Glu Phe Ser Arg Asp Leu 619 667 ctt ccg aga ttc ttc aag cac aat aac ttc tct agc ttt atc cgc cag Leu Pro Arg Phe Phe Lys His Asn Asn Phe Ser Ser Phe Ile Arg Gln ctt aac aca tat ggt ttt aga aaa gct gat cct gag caa tgg gaa ttt Leu Asn Thr Tyr Gly Phe Arg Lys Ala Asp Pro Glu Gln Trp Glu Phe 715 gcg aat gat gat ttt gtg aga ggt caa cct cat ctt atg aag aac att Ala Asn Asp Asp Phe Val Arg Gly Gln Pro His Leu Met Lys Asn Ile 763 811 cat aga cgc aaa cca gtt cat agc cac tct tta ccg aat ctt caa gct His Arg Arg Lys Pro Val His Ser His Ser Leu Pro Asn Leu Gln Ala cag tta aac ccg ttg acg gat tca gaa cga gtg aga atg aat aat cag Gln Leu Asn Pro Leu Thr Asp Ser Glu Arg Val Arg Met Asn Asn Gln 859 att gag aga ttg aca aaa gag aaa gaa gga ttg ctt gaa gag tta cat lle Glu Arg Leu Thr Lys Glu Lys Glu Gly Leu Leu Glu Glu Leu His 907 aaa caa gac gag gaa cga gaa gtg ttt gag atg caa gtg aaa gaa ctt 955 Lys Gln Asp Glu Glu Arg Glu Val Phe Glu Met Gln Val Lys Glu Leu

	MBI15	Sequence	Listing.ST25	
155		160		1

155 160 165	
aaa gaa cga tta caa cac atg gag aag cgt cag aaa aca atg gtt tcg Lys Glu Arg Leu Gln His Met Glu Lys Arg Gln Lys Thr Met Val Ser 170 175 180	1003
ttt gtt tct caa gta ttg gaa aag cca ggg ctt gct ttg aac cta tcg Phe Val Ser Gln Val Leu Glu Lys Pro Gly Leu Ala Leu Asn Leu Ser 185 190 195	1051
ccg tgt gtt ccc gaa aca aac gag agg aaa aga agg ttc cct agg atc Pro Cys Val Pro Glu Thr Asn Glu Arg Lys Arg Arg Phe Pro Arg Ile 200 205 210	1099
gag ttc ttt ccc gat gaa ccg atg ttg gaa gag aac aaa act tgt gtt Glu Phe Phe Pro Asp Glu Pro Met Leu Glu Glu Asn Lys Thr Cys Val 215 220 225 230	1147
gtt gtg aga gag gaa ggt tct aca agc cct tct tca cac aca aga gag Val Val Arg Glu Glu Gly Ser Thr Ser Pro Ser Ser His Thr Arg Glu 235 240 245	1195
cat caa gtg gaa cag tta gag tca tcg ata gcg att tgg gag aat ctt His Gln Val Glu Gln Leu Glu Ser Ser Ile Ala Ile Trp Glu Asn Leu 250 255 260	1243
gta tcg gat tct tgt gag agt atg tta caa tca aga agt atg atg aca Val Ser Asp Ser Cys Glu Ser Met Leu Gln Ser Arg Ser Met Met Thr 265 270 275	1291
ctt gat gtg gat gaa tca tct act ttt cca gag agc cct cct ctt tct Leu Asp Val Asp Glu Ser Ser Thr Phe Pro Glu Ser Pro Pro Leu Ser 280 285 290	1339
tgc ata cag tta agt gtc gat tca cgt ctc aaa tct cct cct tct cca Cys Ile Gln Leu Ser Val Asp Ser Arg Leu Lys Ser Pro Pro Ser Pro 295 300 305 310	1387
agg atc atc gat atg aac tgt gag ccc gat ggt tcg aaa gaa cag aac Arg Ile Ile Asp Met Asn Cys Glu Pro Asp Gly Ser Lys Glu Gln Asn 315 320 325	1435
act gtt gct gct cct cct cct cca gta gca gga gcg aat gat ggc Thr Val Ala Ala Pro Pro Pro Pro Pro Val Ala Gly Ala Asn Asp Gly 330 335 340	1483
ttc tgg cag cag ttt ttc tca gag aat cct ggc tca acc gag caa cgg Phe Trp Gln Gln Phe Phe Ser Glu Asn Pro Gly Ser Thr Glu Gln Arg 345 350 355	1531
gaa gtt caa tta gag agg aaa gac gat aaa gat aaa gcc gga gta cgt Glu Val Gln Leu Glu Arg Lys Asp Asp Lys Asp Lys Ala Gly Val Arg 360 365 370	1579
act gag aaa tgt tgg tgg aat tcg aga aat gtt aat gca att aca gaa Thr Glu Lys Cys Trp Trp Asn Ser Arg Asn Val Asn Ala Ile Thr Glu 375 380 385 390	1627
cag ctt gga cat ctg act tct tca gag aga agt tga tatgtcaaag Gln Leu Gly His Leu Thr Ser Ser Glu Arg Ser 395 400	1673
attaaatttc tagtctgttt tagttacttg taaaataggg tttctcagtt ttattgtttt	1733
cgattccagt acttaggtat ggttcagctg tttatttatc acttgtatga tctttcccag	1793
ttcattgtag cagacttcaa tggtaatgat.aagctagagc ttatggatag tattcataaa	1853
aaaa	1857

<210> 8 <211> 401 <212> PRT <213> Arabidopsis thaliana

MBI15 Sequence Listing.ST25

<400> 8

Met Asp Glu Asn Asn His Gly Val Ser Ser Ser Ser Leu Pro Pro Phe 1 5 10 15

Leu Thr Lys Thr Tyr Glu Met Val Asp Asp Ser Ser Ser Asp Ser Ile 20 25 30

Val Ser Trp Ser Gln Ser Asn Lys Ser Phe Ile Val Trp Asn Pro Pro 35 40 45

Glu Phe Ser Arg Asp Leu Leu Pro Arg Phe Phe Lys His Asn Asn Phe 50 55 60

Ser Ser Phe Ile Arg Gln Leu Asn Thr Tyr Gly Phe Arg Lys Ala Asp 65 70 75 80

Pro Glu Gln Trp Glu Phe Ala Asn Asp Asp Phe Val Arg Gly Gln Pro 85 90 95

His Leu Met Lys Asn Ile His Arg Arg Lys Pro Val His Ser His Ser 100 105 110

Leu Pro Asn Leu Gln Ala Gln Leu Asn Pro Leu Thr Asp Ser Glu Arg 115 120 125

Val Arg Met Asn Asn Gln Ile Glu Arg Leu Thr Lys Glu Lys Glu Gly 130 135 140

Leu Leu Glu Glu Leu His Lys Gln Asp Glu Glu Arg Glu Val Phe Glu 145 150 150 160

Met Gln Val Lys Glu Leu Lys Glu Arg Leu Gln His Met Glu Lys Arg 165 170 175

Gln Lys Thr Met Val Ser Phe Val Ser Gln Val Leu Glu Lys Pro Gly 180 185 190

Leu Ala Leu Asn Leu Ser Pro Cys Val Pro Glu Thr Asn Glu Arg Lys

Arg Arg Phe Pro Arg Ile Glu Phe Phe Pro Asp Glu Pro Met Leu Glu 210 215 220

Glu Asn Lys Thr Cys Val Val Val Arg Glu Glu Gly Ser Thr Ser Pro 225 230 235 240

Ser Ser His Thr Arg Glu His Gln Val Glu Gln Leu Glu Ser Ser Ile 245 250 255

Ala Ile Trp Glu Asn Leu Val Ser Asp Ser Cys Glu Ser Met Leu Gln 260 265 270

Ser Arg Ser Met Met Thr Leu Asp Val Asp Glu Ser Ser Thr Phe Pro

MBI15 Sequence Listing.ST25 Glu Ser Pro Pro Leu Ser Cys Ile Gln Leu Ser Val Asp Ser Arg Leu Lys Ser Pro Pro Ser Pro Arg Ile Ile Asp Met Asn Cys Glu Pro Asp Gly Ser Lys Glu Gln Asn Thr Val Ala Ala Pro Pro Pro Pro Val 325 Ala Gly Ala Asn Asp Gly Phe Trp Gln Gln Phe Phe Ser Glu Asn Pro Gly Ser Thr Glu Gln Arg Glu Val Gln Leu Glu Arg Lys Asp Asp Lys Asp Lys Ala Gly Val Arg Thr Glu Lys Cys Trp Trp Asn Ser Arg Asn Val Asn Ala Ile Thr Glu Gln Leu Gly His Leu Thr Ser Ser Glu Arg 395 390 385 Ser <210> 9 <211> 964 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (63)..(869) <223> G28 <400> 9 gaaatctcaa caagaaccaa accaaacaac aaaaaaaacat tcttaataat tatctttctg 60 tt atg tcg atg acg gcg gat tct caa tct gat tat gct ttt ctt gag Met Ser Met Thr Ala Asp Ser Gln Ser Asp Tyr Ala Phe Leu Glu 107 155 tcc ata cga cga cac tta cta gga gaa tcg gag ccg ata ctc agt gag Ser Ile Arg Arg His Leu Leu Gly Glu Ser Glu Pro Ile Leu Ser Glu tcg aca gcg agt tcg gtt act caa tct tgt gta acc ggt cag agc att Ser Thr Ala Ser Ser Val Thr Gln Ser Cys Val Thr Gly Gln Ser Ile 203 251 aaa ccg gtg tac gga cga aac cct agc ttt agc aaa ctg tat cct tgc Lys Pro Val Tyr Gly Arg Asn Pro Ser Phe Ser Lys Leu Tyr Pro Cys ttc acc gag agc tgg gga gat ttg ccg ttg aaa gaa aac gat tct gag Phe Thr Glu Ser Trp Gly Asp Leu Pro Leu Lys Glu Asn Asp Ser Glu 65 70 75 299 gat atg tta gtt tac ggt atc ctc aac gac gcc ttt cac ggc ggt tgg
Asp Met Leu Val Tyr Gly Ile Leu Asn Asp Ala Phe His Gly Gly Trp 347 395 gag ccg tct tct tcg tct tcc gac gaa gat cgt agc tct ttc ccg agt Glu Pro Ser Ser Ser Ser Asp Glu Asp Arg Ser Ser Phe Pro Ser gtt aag atc gag act ccg gag agt ttc gcg gcg gtg gat tct gtt ccg 443 Page 12

Val	Lys	Ile	Glu 115	Thr	Pro	Glu					List Val				Pro	
gtc Val	aag Lys	aag Lys 130	gag Glu	aag Lys	acg Thr	agt Ser	cct Pro 135	gtt Val	tcg Ser	gcg Ala	gcg Ala	gtg Val 140	acg Thr	gcg Ala	gcg Ala	491
											ccg Pro 155					539
											gct Ala					587
											gct Ala					635
gct Ala	ttc Phe	agg Arg	atg Met 195	cgt Arg	ggt Gly	tcc Ser	cgc Arg	gct Ala 200	ttg Leu	ttg Leu	aat Asn	ttt Phe	ccg Pro 205	ttg Leu	aga Arg	683
gtt Val	aat Asn	tca Ser 210	gga Gly	gaa Glu	ccc Pro	gac Asp	ccg Pro 215	gtt Val	cga Arg	atc Ile	aag Lys	tcc Ser 220	aag Lys	aga Arg	tct Ser	731
											aag Lys 235					779
gtg Val 240	gcc Ala	gcc Ala	ggt Gly	ggt Gly	gga Gly 245	atg Met	gat Asp	aag Lys	gga Gly	ttg Leu 250	acg Thr	gtg Val	aag Lys	tgc Cys	gag Glu 255	827
											gtt Val		taa			869
ttt	gatt	tt t	cttt	gttg	g at	gatt	atat	gat	tctt	caa	aaaa	gaag	gaa o	gtta	ataaa	929
aaaa	ttcg	gtt t	atta	ittaa	a aa	aaaa	aaaa	a aaa	aaa							964
<210 <211 <212 <213	> 2 > E	10 268 PRT Arabi	idops	sis t	hali:	ana										
<400)> 1	١٥														
Met l	Ser	Met	Thr	Ala 5	Asp	Ser	Gln	Ser	Asp 10	Tyr	Ala	Phe	Leu	Glu 15	Ser	
Ile	Arg	Arg	His 20	Leu	Leu	Gly	Glu	Ser 25	Glu	Pro	Ile	Leu	Ser 30	Glu	Ser	
Chr	Ala	Ser 35	Ser	Val	Thr	Gln	Ser 40	Cys	Val	Thr	Gly	Gln 45	Ser	Ile	Lys	
Pro	Val 50	Tyr	Gly	Arg	Asn	Pro 55	Ser	Phe	Ser	Lys	Leu 60	Tyr	Pro	Сув	Phe	
Thr 55	Glu	Ser	Trp	Gly	Asp 70	Leu	Pro	Leu	Lys	Glu 75	Asn	Asp	Ser	Glu	Asp 80	
let	Leu	Val	Tyr	Gly 85	Ile	Leu	Asn	Asp	Ala 90	Phe	His	Gly	Gly	Trp 95	Glu .	

MBI15 Sequence Listing.ST25 Pro Ser Ser Ser Ser Asp Glu Asp Arg Ser Ser Phe Pro Ser Val 100 105 Lys Ile Glu Thr Pro Glu Ser Phe Ala Ala Val Asp Ser Val Pro Val Lys Lys Glu Lys Thr Ser Pro Val Ser Ala Ala Val Thr Ala Ala Lys Gly Lys His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly Lys Phe Ala Ala Glu Ile Arg Asp Pro Ala Lys Asn Gly Ala Arg Val Trp Leu Gly 165 170 175 Thr Phe Glu Thr Ala Glu Asp Ala Ala Leu Ala Tyr Asp Arg Ala Ala Phe Arg Met Arg Gly Ser Arg Ala Leu Leu Asn Phe Pro Leu Arg Val Asn Ser Gly Glu Pro Asp Pro Val Arg Ile Lys Ser Lys Arg Ser Ser Phe Ser Ser Ser Asn Glu Asn Gly Ala Pro Lys Lys Arg Arg Thr Val Ala Ala Gly Gly Met Asp Lys Gly Leu Thr Val Lys Cys Glu Val Val Glu Val Ala Arg Gly Asp Arg Leu Leu Val Leu 265 <210> 11 <211> 1571 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (428)..(1402) <223> G869 <400> 11 aggaacagtg aaaggttcgg tttttttgggt ttcgatctga taatcaacaa gaaaaaaggg 60 tttgatttat gtcggctggg tttgaatcga ctgtgatttt gtctttgatt catatctctt 120 ctccgatttc atcatcatct tccccatcat cgtcgtcttt gaaatcttgt cttctcaacg 180 ctcttcactt ctgctgtaat aagcagaggc ttgttctgga gactccttct ctttccatgc 240 gcttaagacc caaaaggact tgttctagtg ttgaagtctt tgggggtttt cacataaagc 300 agcaaaagtt ttctttttc atagttcgct gagagttttg agttttgata ccaaaaaagt 360 tttgaccttt tagagtgatt ttttgttctt tctgttttct gggtattttt gaggagtggg 420 tttaaca atg gtt gcg att aga aag gaa cag tct ttg agt ggt gtt agt Met Val Ala Ile Arg Lys Glu Gln Ser Leu Ser Gly Val Ser 469 age gag att aag aag aga get aag aga aac act eta teg tee ett eet 517

								15 S								
Ser 15	Glu	Ile	Lys	Lys	Arg 20	Ala	Lys	Arg	Asn	Thr 25	Leu	Ser	Ser	Leu	Pro 30	
caa Gln	gaa Glu	acc Thr	caa Gln	cct Pro 35	ttg Leu	agg Arg	aaa Lys	gtc Val	cgt Arg 40	att Ile	att Ile	gtg Val	aat Asn	gat Asp 45	cct Pro	565
tat Tyr	gct Ala	act Thr	gat Asp 50	gat Asp	tcc Ser	tct Ser	agt Ser	gat Asp 55	gag Glu	gaa Glu	gag Glu	ctt Leu	aag Lys 60	gtt Val	cct Pro	613
aag Lys	cca Pro	agg Arg 65	aaa Lys	atg Met	aaa Lys	cgt Arg	atc Ile 70	gtt Val	cgt Arg	gag Glu	att Ile	aac Asn 75	ttt Phe	cct Pro	tct Ser	661
atg Met	gaa Glu 80	gtt Val	tct Ser	gaa Glu	cag Gln	cct Pro 85	tct Ser	gag Glu	agt Ser	tct Ser	tct Ser 90	cag Gln	gac Asp	agt Ser	act Thr	709
aaa Lys 95	act Thr	gat Asp	ggc Gly	aag Lys	ata Ile 100	gct Ala	gtg Val	tca Ser	gct Ala	tct Ser 105	cct Pro	gct Ala	gtt Val	cct Pro	agg Arg 110	757
aag Lys	aag Lys	cct Pro	gtt Val	ggt Gly 115	gtt Val	agg Arg	caa Gln	agg Arg	aaa Lys 120	tgg Trp	ggg Gly	aaa Lys	tgg Trp	gct Ala 125	gct Ala	805
gag Glu	att Ile	aga Arg	gat Asp 130	cct Pro	att Ile	aag Lys	aaa Lys	act Thr 135	agg Arg	act Thr	tgg Trp	ttg Leu	ggt Gly 140	act Thr	ttt Phe	853
gat Asp	act Thr	ctt Leu 145	gaa Glu	gaa Glu	gct Ala	gct Ala	aaa Lys 150	gct Ala	tat Tyr	gat Asp	gct Ala	aag Lys 155	aag Lys	ctt Leu	gag Glu	901
ttt Phe	gat Asp 160	gct Ala	att Ile	gtt Val	gct Ala	gga Gly 165	aat Asn	gtg Val	tcc Ser	act Thr	act Thr 170	aaa Lys	cgt Arg	gat Asp	gtt Val	949
tct Ser 175	tca Ser	tct Ser	gag Glu	act Thr	agc Ser 180	caa Gln	tgc Cys	tct Ser	cgt Arg	tct Ser 185	tca Ser	cct Pro	gtt Val	gtt Val	cct Pro 190	997
gtt Val	gag Glu	caa Gln	gat Asp	gac Asp 195	act Thr	tct Ser	gca Ala	tca Ser	gct Ala 200	ctc Leu	act Thr	tgt Cys	gtc Val	aac Asn 205	aac Asn	1045
cct Pro	gat Asp	gac Asp	gtc Val 210	tcg Ser	acc Thr	gtt Val	gct Ala	cca Pro 215	act Thr	gct Ala	cca Pro	act Thr	cca Pro 220	aat Asn	gtt Val	1093
cct Pro	gct Ala	ggt Gly 225	gga Gly	aac Asn	aag Lys	gaa Glu	acg Thr 230	ttg Leu	ttc Phe	gat Asp	ttc Phe	gac Asp 235	ttt Phe	act Thr	aat Asn	1141
cta Leu	cag Gln 240	atc Ile	cct Pro	gat Asp	ttt Phe	ggt Gly 245	ttc Phe	ttg Leu	gca Ala	gag Glu	gag Glu 250	caa Gln	caa Gln	gac Asp	cta Leu	1189
gac Asp 255	ttc Phe	gat Asp	tgt Cys	ttc Phe	ctc Leu 260	gcg Ala	gat Asp	gat Asp	cag Gln	ttt Phe 265	gat Asp	gat Asp	ttc Phe	ggc Gly	ttg Leu 270	1237
ctt Leu	gat Asp	gac Asp	att Ile	caa Gln 275	gga Gly	ttc Phe	gaa Glu	gat Asp	aac Asn 280	ggt Gly	cca Pro	agt Ser	gcg Ala	tta Leu 285	cca Pro	1285
gat Asp	ttc Phe	gac Asp	ttt Phe 290	gcg Ala	gat Asp	gtt Val	gaa Glu	gat Asp 295	ctt Leu	cag Gln	cta Leu	gct Ala	gac Asp 300	tct Ser	agt Ser	1333
ttc Phe	ggt Gly	ttc Phe 305	ctt Leu	gat Asp	caa Gln	ctt Leu	gct Ala 310	cct Pro	atc Ile	aac Asn	atc Ile	tct Ser 315	tgc Cys	cca Pro	tta Leu	1381

Page 15

MBI15 Sequence Listing.ST25 aaa agt ttt gca gct tca tag gatcttgctt agtaatgtta agtgagaaga Lys Ser Phe Ala Ala Ser 320
gtgttttgtt ttttcgttta tgctttagta atttaagaca tacaaaagtg tgtgttccgg
attgtagtaa gatcttaaga cataaagccg ggttttgcaa ttaggaatcg agttttaatg
aagttttagt ttatgtttg
<210> 12 <211> 324 <212> PRT <213> Arabidopsis thaliana
<400> 12
Met Val Ala Ile Arg Lys Glu Gln Ser Leu Ser Gly Val Ser Ser Glu 1 10 15
Ile Lys Lys Arg Ala Lys Arg Asn Thr Leu Ser Ser Leu Pro Gln Glu 20 25 30
Thr Gln Pro Leu Arg Lys Val Arg Ile Ile Val Asn Asp Pro Tyr Ala 35 40 45
Thr Asp Asp Ser Ser Ser Asp Glu Glu Leu Lys Val Pro Lys Pro 50 55 60
Arg Lys Met Lys Arg Ile Val Arg Glu Ile Asn Phe Pro Ser Met Glu 65 70 75 80
Val Ser Glu Gln Pro Ser Glu Ser Ser Ser Gln Asp Ser Thr Lys Thr 85 90 95
Asp Gly Lys Ile Ala Val Ser Ala Ser Pro Ala Val Pro Arg Lys Lys
Pro Val Gly Val Arg Gln Arg Lys Trp Gly Lys Trp Ala Ala Glu Ile 115 120 125
Arg Asp Pro Ile Lys Lys Thr Arg Thr Trp Leu Gly Thr Phe Asp Thr 130 135 140
Leu Glu Glu Ala Ala Lys Ala Tyr Asp Ala Lys Lys Leu Glu Phe Asp 145 150 155 160
Ala Ile Val Ala Gly Asn Val Ser Thr Thr Lys Arg Asp Val Ser Ser 165 170 175
Ser Glu Thr Ser Gln Cys Ser Arg Ser Ser Pro Val Val Pro Val Glu 180 185 190
Gln Asp Asp Thr Ser Ala Ser Ala Leu Thr Cys Val Asn Asn Pro Asp 195 200 205
Asp Val Ser Thr Val Ala Pro Thr Ala Pro Thr Pro Asn Val Pro Ala 210 215 220
Gly Gly Asn Lys Glu Thr Leu Phe Asp Phe Asp Phe Thr Asn Leu Gln

MBI15 Sequence Listing.ST25

Ile Pro Asp Phe Gly Phe Leu Ala Glu Glu Gln Gln Asp Leu Asp Phe

230

Asp Cys Phe Leu Ala Asp Asp Gln Phe Asp Asp Phe Gly Leu Leu Asp

Asp Ile Gln Gly Phe Glu Asp Asn Gly Pro Ser Ala Leu Pro Asp Phe 275 280 285

Asp Phe Ala Asp Val Glu Asp Leu Gln Leu Ala Asp Ser Ser Phe Gly

Phe Leu Asp Gln Leu Ala Pro Ile Asn Ile Ser Cys Pro Leu Lys Ser

Phe Ala Ala Ser

225

<210> 13 <211> 920 <212> DNA <213> Arabidopsis thaliana

<220> <221> CDS <222> (1)..(852) <223> G237

atg gcg aag acg aaa tat gga gag aga cat agg aaa ggg tta tgg tca Met Ala Lys Thr Lys Tyr Gly Glu Arg His Arg Lys Gly Leu Trp Ser 48 cct gaa gaa gac gag aag cta agg agc ttc atc ctc tct tat ggc cat Pro Glu Glu Asp Glu Lys Leu Arg Ser Phe Ile Leu Ser Tyr Gly His 96 tct tgc tgg acc act gtt ccc atc aaa gct ggg tta caa agg aat ggg Ser Cys Trp Thr Thr Val Pro Ile Lys Ala Gly Leu Gln Arg Asn Gly 144 aag agc tgc aga tta aga tgg att aat tac cta aga cca ggg tta aag Lys Ser Cys Arg Leu Arg Trp Ile Asn Tyr Leu Arg Pro Gly Leu Lys agg gat atg att agt gca gaa gaa gaa gag act atc ttg acg ttt cat Arg Asp Met Ile Ser Ala Glu Glu Glu Glu Thr Ile Leu Thr Phe His 240 70

tct ccc ttg ggt aac aag tgg tcg caa ata gct aaa ttc tta ccg gga Ser Pro Leu Gly Asn Lys Trp Ser Gln Ile Ala Lys Phe Leu Pro Gly 85 90 95 288

aga aca gac aat gag ata aag aac tat tgg cac tct cat ttg aaa aag Arg Thr Asp Asn Glu Ile Lys Asn Tyr Trp His Ser His Leu Lys Lys 336 100 105

aaa tgg ctc aag tct cag agc tta caa gat gca aaa tct att tcc cct Lys Trp Leu Lys Ser Gln Ser Leu Gln Asp Ala Lys Ser Ile Ser Pro 384

cct tcg tct tca tca tca tca ctt gtt gct tgt gga gaa aga aat ccg Pro Ser Ser Ser Ser Ser Leu Val Ala Cys Gly Glu Arg Asn Pro 432

							MOT	15 6		200	T.i.o.		cmac			
gaa Glu 145	acc Thr	ttg Leu	atc Ile	tcg Ser	aat Asn 150	cac His	gtg	ttc	tcc	ctc	List cag Gln	aga	ctt	cta	gag Glu 160	480
aac Asn	aaa Lys	tct Ser	tca Ser	tct Ser 165	ccc Pro	tca Ser	caa Gln	gaa Glu	agc Ser 170	aac Asn	gga Gly	aat Asn	aac Asn	agc Ser 175	cat His	528
				gct Ala												576
ctt Leu	tct Ser	tct Ser 195	tca Ser	tat Tyr	ccc Pro	cac His	acc Thr 200	gat Asp	tat Tyr	tcc Ser	tct Ser	gag Glu 205	ttt Phe	acc Thr	gac Asp	624
tct Ser	aag Lys 210	cac His	agt Ser	caa Gln	gct Ala	cca Pro 215	aat Asn	gtc Val	gaa Glu	gag Glu	act Thr 220	ctc Leu	tca Ser	gct Ala	tat Tyr	672
gaa Glu 225	gaa Glu	atg Met	ggt Gly	gat Asp	gtt Val 230	gat Asp	cag Gln	ttc Phe	cat His	tac Tyr 235	aac Asn	gaa Glu	atg Met	atg Met	atc Ile 240	720
aac Asn	aac Asn	agc Ser	aac Asn	tgg Trp 245	act Thr	ctt Leu	aac Asn	gac Asp	att Ile 250	gtg Val	ttt Phe	ggt Gly	tcc Ser	aaa Lys 255	tgt Cys	768
				cat His												816
				ttt Phe							taa	atto	gcgti	tta		862
ttg	taato	gta a	aatca	aaatt	t ct	aagg	gcaaa	a acc	ggaa	aaaa	aaaa	aaaaa	aaa a	aaaa	aaaa	920
<210 <211 <211 <211	0> 1 1> 2 2> 1	14 283 PRT		aaati			gcaaa	a aco	eggaa	aaaa	aaaa	aaaa	aaa a	aaaa	aaa	920
<210 <210 <210	0> 3 1> 2 2> 1 3> 1	14 283 PRT					gcaaa	a aco	cggaa	aaaa	aaaa	aaaa	aaa a	aaaa	aaa	920
<210 <211 <211 <211	0 >	14 283 PRT Arab:	idops		:hali	iana										920
<210 <211 <211 <211 <400 Met	0> 1 1> 2 2> 1 3> 1 0> 1	14 283 PRT Arab: 14 Lys	idops Thr	sis t	chali Tyr	iana Gly	Glu	Arg	His 10	Arg	Lys	Gly	Leu	Trp 15	Ser	920
<210 <211 <211 <211 <400 Met 1	0> 11> 22> 133> 14 0> 21 Ala Glu	14 283 PRT Arab: 14 Lys	idops Thr Asp 20	sis t Lys 5	thali Tyr Lys	iana Gly Leu	Glu Arg	Arg Ser 25	His 10 Phe	Arg Ile	Lys Leu	Gly Ser	Leu Tyr 30	Trp 15	Ser His	920
<210 <211 <211 <400 Met 1 Pro	0> 1 1> 2 2> 1 3> 1 0> 1 Ala Glu	14 283 PRT Arab: 14 Lys Glu Trp 35	idops Thr Asp 20 Thr	Lys 5 Glu	Tyr Lys Val	iana Gly Leu Pro	Glu Arg Ile 40	Arg Ser 25 Lys	His 10 Phe Ala	Arg Ile Gly	Lys Leu Leu	Gly Ser Gln 45	Leu Tyr 30 Arg	Trp 15 Gly Asn	Ser His Gly	920
<211 <211 <211 <400 Met 1 Pro	0)> :: 1> :: 22> 3> Ala Glu Cys	14 283 PRT Arab: 14 Lys Glu Trp 35	Thr Asp 20 Thr	Lys 5 Glu Thr	Tyr Lys Val	Gly Leu Pro Trp 55	Glu Arg Ile 40	Arg Ser 25 Lys Asn	His 10 Phe Ala	Arg Ile Gly Leu	Lys Leu Leu Arg	Gly Ser Gln 45	Leu Tyr 30 Arg	Trp 15 Gly Asn	Ser His Gly Lys	920
<211 <211 <211 <400 Met 1 Pro Ser Lys	Cys Ser 50 Asp	14 283 PRT Arab: 14 Lys Glu Trp 35 Cys	Thr Asp 20 Thr Arg	Lys 5 Glu Thr	Tyr Lys Val Arg	Gly Leu Pro Trp 55	Glu Arg Ile 40	Arg Ser 25 Lys Asn Glu	His 10 Phe Ala Tyr	Arg Ile Gly Leu Thr	Leu Leu Arg 60	Gly Ser Gln 45 Pro Leu	Leu Tyr 30 Arg Gly	Trp 15 Gly Asn Leu	Ser His Gly Lys His 80	920
<211 <211 <211 <400 Met 1 Pro Ser Lys Arg 65	Cys Ser 50 Asp	14 283 PRT Arab: 14 Lys Glu Trp 35 Cys Met	Thr Asp 20 Thr Arg	Lys 5 Glu Thr Leu Ser	Tyr Lys Val Arg Ala 70	Gly Leu Pro Trp 55 Glu Trp	Glu Arg Ile 40 Ile Glu Ser	Arg Ser 25 Lys Asn Glu	His 10 Phe Ala Tyr Glu	Arg Ile Gly Leu Thr 75	Leu Leu Arg 60 Ile	Gly Ser Gln 45 Pro Leu	Leu Tyr 30 Arg Gly Thr	Trp 15 Gly Asn Leu Phe Pro 95	Ser His Gly Lys His 80	920

MBI15 Sequence Listing.ST25 115 Pro Ser Ser Ser Ser Ser Ser Leu Val Ala Cys Gly Glu Arg Asn Pro 130 135 140 Glu Thr Leu Ile Ser Asn His Val Phe Ser Leu Gln Arg Leu Leu Glu Asn Lys Ser Ser Ser Pro Ser Gln Glu Ser Asn Gly Asn Asn Ser His Gln Cys Ser Ser Ala Pro Glu Ile Pro Arg Leu Phe Phe Ser Glu Trp Leu Ser Ser Ser Tyr Pro His Thr Asp Tyr Ser Ser Glu Phe Thr Asp Ser Lys His Ser Gln Ala Pro Asn Val Glu Glu Thr Leu Ser Ala Tyr 210 215 220 Glu Glu Met Gly Asp Val Asp Gln Phe His Tyr Asn Glu Met Met Ile Asn Asn Ser Asn Trp Thr Leu Asn Asp Ile Val Phe Gly Ser Lys Cys Lys Lys Gln Glu His His Ile Tyr Arg Glu Ala Ser Asp Cys Asn Ser 260 265 270 Ser Ala Glu Phe Phe Ser Pro Pro Thr Thr <210> 15 <211> 1302 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (331)..(1149) G409 <400> 15 60 tgtctctctc tctggctctc tttctcttaa cgtgatcata acgtgattcg aaaattggat 120 atagataggt ttcttgttgg atcttgatcc ctctggaaaa ggaggggaga atagcagttc 180 atgatgggat tttgtatctg cccgttggag tcacctgcga gattactatg gagtacaagc ttcttccgcc ataagatcat gatcttctaa tccttcctac ttcttcccat ctttttaatc 240 300 atcttctcqc tatctctqct tcctctttct ctctqtttcc tctttctcag aactcagaag tagttgttgt tttatttctg ttgatcaaaa atg gaa tcc aat tcg ttt ttc ttc
Met Glu Ser Asn Ser Phe Phe Phe 354 gat cca tct gct tca cac ggc aac agc atg ttc ttc ctt ggg aat ctc Asp Pro Ser Ala Ser His Gly Asn Ser Met Phe Phe Leu Gly Asn Leu 402 aat ccc gtc gtc caa gga gga gga gca aga tcg atg atg aac atg gag Asn Pro Val Val Gln Gly Gly Gly Ala Arg Ser Met Met Asn Met Glu 450

		MBI15 Seque	nce Listing.S		
25	30		35		40
gaa act tcg a Glu Thr Ser I	aag cga agg ccc Lys Arg Arg Pro 45	ttc ttt agc Phe Phe Ser 50	ser Pro Glu	gat ctc Asp Leu 55	tac 498 Tyr
Asp Asp Asp I	ttt tac gac gac Phe Tyr Asp Asp 60	cag ttg cct Gln Leu Pro 65	Glu Lys Lys	cgt cgc Arg Arg 70	ctc 546 Leu
	caa gtg cat ctg Gln Val His Leu				
	cct gaa cgc aag Pro Glu Arg Lys 95				
cag cca agg o Gln Pro Arg (105	caa gtg gct gtc Gln Val Ala Val 110	tgg ttt cag Trp Phe Gln	aat cgc cga (Asn Arg Arg) 115	gct cgt Ala Arg	tgg 690 Trp 120
aaa aca aaa d Lys Thr Lys (cag ctt gag aga Gln Leu Glu Arg 125	gac tac gat Asp Tyr Asp 130	ctt ctc aag : Leu Leu Lys :	tcc act Ser Thr 135	tac 738 Tyr
Asp Gln Leu I	ctt tct aac tac Leu Ser Asn Tyr 140	gac tcc atc Asp Ser Ile 145	Val Met Asp	aac gat Asn Asp 150	aag 786 Lys
ctc aga tcc o Leu Arg Ser (155	gag gtt act tcc Glu Val Thr Ser	ctg acc gaa Leu Thr Glu 160	aag ctt cag Lys Leu Gln 165	ggc aaa Gly Lys	caa 834 Gln
gag aca gct a Glu Thr Ala A 170	aat gaa cca cct Asn Glu Pro Pro 175	ggt caa gtg Gly Gln Val	ccc gaa cca a Pro Glu Pro 1 180	aac caa Asn Gln	ctt 882 Leu
gat ccg gtt t Asp Pro Val 1 185	tac att aat gcg Tyr Ile Asn Ala 190	gca gca atc Ala Ala Ile	aaa acc gag Lys Thr Glu 195	gac cgg Asp Arg	tta 930 Leu 200
agt tca ggg a Ser Ser Gly S	agc gtt ggg agc Ser Val Gly Ser 205	gcg gta cta Ala Val Leu 210	gac	gca cct Ala Pro 215	caa 978 Gln
Leu Leu Asp	agc tgt gac tct Ser Cys Asp Ser 220	tac ttc cca Tyr Phe Pro 225	Ser Ile Val	ccc atc Pro Ile 230	caa 1026 Gln
gac aac agc a Asp Asn Ser A 235	aac gcc agt gat Asn Ala Ser Asp	cat gac aat His Asp Asn 240	gac cgg agc Asp Arg Ser 245	tgt ttc Cys Phe	gcc 1074 Ala
gac gtc ttt g Asp Val Phe V 250	gtg ccc acc act Val Pro Thr Thr 255	Ser Pro Ser	cac gat cat his Asp His 1 260	cac ggt His Gly	gaa 1122 Glu
	ttc tgg gga tgg Phe Trp Gly Trp 270		accactc tgata	ataaa	1169
tgtgtgttta t	ttaagttca agagt	catct tcttgt	tgtt tccatgtt	ga cgata	attgt 1229
tgactcgtgg aa	ataattccg ctgtt	caacg gtattt	ttat cagttgca	tt atatg	ctttt 1289

atgaaaaaaa aaa

<210> 16 <211> 272 <212> PRT <213> Arabidopsis thaliana

<400> 16

1302

MBI15 Sequence Listing.ST25

Met Glu Ser Asn Ser Phe Phe Phe Asp Pro Ser Ala Ser His Gly Asn 1 10 15

Ser Met Phe Phe Leu Gly Asn Leu Asn Pro Val Val Gln Gly Gly

Ala Arg Ser Met Met Asn Met Glu Glu Thr Ser Lys Arg Arg Pro Phe 35 40 45

Phe Ser Ser Pro Glu Asp Leu Tyr Asp Asp Asp Phe Tyr Asp Asp Gln 50 55

Leu Pro Glu Lys Lys Arg Arg Leu Thr Thr Glu Gln Val His Leu Leu 65 70 80

Glu Lys Ser Phe Glu Thr Glu Asn Lys Leu Glu Pro Glu Arg Lys Thr 85 90 95

Gln Leu Ala Lys Lys Leu Gly Leu Gln Pro Arg Gln Val Ala Val Trp

Phe Gln Asn Arg Arg Ala Arg Trp Lys Thr Lys Gln Leu Glu Arg Asp 115 120 125

Tyr Asp Leu Leu Lys Ser Thr Tyr Asp Gln Leu Leu Ser Asn Tyr Asp 130 135 140

Ser Ile Val Met Asp Asn Asp Lys Leu Arg Ser Glu Val Thr Ser Leu 145 150 160

Thr Glu Lys Leu Gln Gly Lys Gln Glu Thr Ala Asn Glu Pro Pro Gly 165 170 175

Gln Val Pro Glu Pro Asn Gln Leu Asp Pro Val Tyr Ile Asn Ala Ala

Ala Ile Lys Thr Glu Asp Arg Leu Ser Ser Gly Ser Val Gly Ser Ala 195 200 205

Val Leu Asp Asp Asp Ala Pro Gln Leu Leu Asp Ser Cys Asp Ser Tyr 210 215 220

Phe Pro Ser Ile Val Pro Ile Gln Asp Asn Ser Asn Ala Ser Asp His

Asp Asn Asp Arg Ser Cys Phe Ala Asp Val Phe Val Pro Thr Thr Ser

Pro Ser His Asp His His Gly Glu Ser Leu Ala Phe Trp Gly Trp Pro 260 265 270

<210> 17 <211> 2545 <212> DNA

<213> Arabidopsis thaliana

<220>

MBI15 Sequence Listing.ST25

<221> CDS <222> (103)..(2322) '<223> G418

<400> 17 acatatgttt taaattottt gtotgaatot tacaggatoo gagagagaga gototggaac 60 gatattaaca tatatcatga agaaaaagat tgaagtattg at atg gga ata act $$\operatorname{\mathtt{Met}}$$ Gly Ile Thr 114 aaa act tot oot aat act aca att oto ttg aag act ttt cac aat aat 162 Lys Thr Ser Pro Asn Thr Thr Ile Leu Leu Lys Thr Phe His Asn Asn tct atg tcc caa gat tat cat cat cat cat cat cat aat caa cac caa 210 Ser Met Ser Gln Asp Tyr His His His His His His Asn Gln His Gln gga ggt atc ttc aac ttc tct aat gga ttc gac cga tca gat tct ccc Gly Gly Ile Phe Asn Phe Ser Asn Gly Phe Asp Arg Ser Asp Ser Pro 258 aat tta aca act cag cag aag caa gag cat caa agg gta gag atg gac Asn Leu Thr Thr Gln Gln Lys Gln Glu His Gln Arg Val Glu Met Asp 306 gag gaa tot toa gto goo gga ggt agg att cog gto tac gaa toa goo Glu Glu Ser Ser Val Ala Gly Gly Arg Ile Pro Val Tyr Glu Ser Ala 70 75 80 354 ggt atg tta tcc gaa atg ttt aat ttc ccc gga agc agc ggt gga gga Gly Met Leu Ser Glu Met Phe Asn Phe Pro Gly Ser Ser Gly Gly Gly 402 aga gat ctc gac ctc ggc caa tct ttc cgg tca aat agg cag ttg ctt Arg Asp Leu Asp Leu Gly Gln Ser Phe Arg Ser Asn Arg Gln Leu Leu 498 gag gag caa cat cag aat att ccg gct atg aat gct acg gat tca gcc Glu Glu Gln His Gln Asn Ile Pro Ala Met Asn Ala Thr Asp Ser Ala acc gcc acc gca gcc gcc atg cag tta ttc ttg atg aat cca ccg cca Thr Ala Thr Ala Ala Ala Met Gln Leu Phe Leu Met Asn Pro Pro Pro 546 ccg caa caa cca ccg tct ccg tca tcc aca act tcc cca agg agc cac Pro Gln Gln Pro Pro Ser Pro Ser Ser Thr Thr Ser Pro Arg Ser His 594 cac aat tot toa act ott cac atg tta ott coa agt coa too acc aac His Asn Ser Ser Thr Leu His Met Leu Leu Pro Ser Pro Ser Thr Asn 642 aca act cac cat cag aac tac act aat cat atg tct atg cat cag ctt Thr Thr His His Gln Asn Tyr Thr Asn His Met Ser Met His Gln Leu 690 190 185 cca cat cag cat cac caa cag ata tcg acg tgg cag tct tct ccc gat Pro His Gln His Gln Gln Ile Ser Thr Trp Gln Ser Ser Pro Asp 738 205 cat cat cat cat cat cac aac agc caa acg gag att ggg acc gtc cac His His His His His Asn Ser Gln Thr Glu Ile Gly Thr Val His 215 220 225 786 gtg gaa aac agc gga gga cac gga gga caa ggc ttg tcc tta tct ctc Val Glu Asn Ser Gly Gly His Gly Gly Gln Gly Leu Ser Leu Ser Leu 834 235 tca tcg tct tta gag gct gca gca aaa gcg gaa gag tat aga aac att Ser Ser Ser Leu Glu Ala Ala Ala Lys Ala Glu Glu Tyr Arg Asn Ile 882

MBI15 Sequence Listing.ST25

						tct Ser										930
						gct Ala										978
						tct Ser										1026
						aac Asn 315										1074
						agt Ser										1122
						aac Asn										1170
						gcc Ala										1218
						gag Glu										1266
						gac Asp 395										1314
atg Met 405	cag Gln	atg Met	gtt Val	gtg Val	aac Asn 410	tct Ser	ttc Phe	gac Asp	ata Ile	gta Val 415	atg Met	ggc Gly	cac His	ggt Gly	gcg Ala 420	1362
						ttg Leu										1410
						gtt Val										1458
						gca Ala										1506
						ttg Leu 475										1554
						atg Met										1602
cca Pro	caa Gln	cgc Arg	ggt Gly	ttg Leu 505	cct Pro	gaa Glu	ege Arg	tcc Ser	gtc Val 510	aat Asn	ata Ile	ctt Leu	aga Arg	gct Ala 515	tgg Trp	1650
						cac His										1698
						ggt Gly										1746
						cgt Arg			ГЛS		Met					1794

		MBI15	Sequence	Listing.ST2	25
50	555			560	

	550					555			-		560					
					aaa Lys 570											1842
					gaa Glu											1890
					gaa Glu											1938
					acc Thr											1986
					cgt Arg											2034
					cta Leu 650											2082
					gac Asp											2130
					gcc Ala											2178
gat Asp	gct Ala	gac Asp 695	atg Met	gat Asp	ggt Gly	gtt Val	aac Asn 700	gtt Val	ata Ile	agg Arg	ttt Phe	999 Gly 705	aca Thr	aac Asn	cct Pro	2226
act Thr	ggt Gly 710	gac Asp	gtg Val	tct Ser	ctc Leu	acg Thr 715	ctt Leu	ggt Gly	tta Leu	cgc Arg	cac His 720	gct Ala	gga Gly	aac Asn	atg Met	2274
cct Pro 725	gac Asp	aaa Lys	gac Asp	gct Ala	tct Ser 730	ttc Phe	tgc Cys	gtt Val	aga Arg	gag Glu 735	ttt Phe	999 Gly	ggt Gly	ttt Phe	tag	2322
tttg	cttt	tg t	cact	ccat	t ta	atta	atta	att	atag	gttt	tcca	attct	ta c	ttat	tttaa	2382
ttga	aaat	ct a	tttt	tgto	t ct	taaa	agto	caa	acaa	atac	atta	gtct	ag c	cct	ctctg	2442
cttt	ttt	tt t	ctat	ctc	gt ga	agag	gaaga	a aaa	cgat	acg	taaa	tccc	tt c	gaaa	actaa	2502
tgta	cgtt	gt a	cgac	ttat	t gt	tttc	cataa	a aaa	aaaa	aaaa	aaa					2545
<210	> 1	.8														

Met Gly Ile Thr Lys Thr Ser Pro Asn Thr Thr Ile Leu Leu Lys Thr 1 15

Phe His Asn Asn Ser Met Ser Gln Asp Tyr His His His His His His 20 25 30

Asn Gln His Gln Gly Gly Ile Phe Asn Phe Ser Asn Gly Phe Asp Arg 35 40

Ser Asp Ser Pro Asn Leu Thr Thr Gln Gln Lys Gln Glu His Gln Arg Page 24

<210> 18 <211> 739 <212> PRT <213> Arabidopsis thaliana

MBI15 Sequence Listing.ST25

Val Glu Met Asp Glu Glu Ser Ser Val Ala Gly Gly Arg Ile Pro Val 65 70 80

Tyr Glu Ser Ala Gly Met Leu Ser Glu Met Phe Asn Phe Pro Gly Ser

Ser Gly Gly Gry Arg Asp Leu Asp Leu Gly Gln Ser Phe Arg Ser Asn 100 105 110

Arg Gln Leu Leu Glu Glu Gln His Gln Asn Ile Pro Ala Met Asn Ala 115 120 125

Thr Asp Ser Ala Thr Ala Thr Ala Ala Ala Met Gln Leu Phe Leu Met 130 135 140

Asn Pro Pro Pro Pro Gln Gln Pro Pro Ser Pro Ser Ser Thr Thr Ser 145 150 155 160

Pro Arg Ser His His Asn Ser Ser Thr Leu His Met Leu Leu Pro Ser 165 170 175

Pro Ser Thr Asn Thr Thr His His Gln Asn Tyr Thr Asn His Met Ser

Met His Gln Leu Pro His Gln His His Gln Gln Ile Ser Thr Trp Gln 195 200 205

Ser Ser Pro Asp His His His His His His Asn Ser Gln Thr Glu Ile 210 215 220

Gly Thr Val His Val Glu Asn Ser Gly Gly His Gly Gly Gln Gly Leu 225 230 235 240

Ser Leu Ser Leu Ser Ser Leu Glu Ala Ala Lys Ala Glu Glu 245 250 255

Tyr Arg Asn Ile Tyr Tyr Gly Ala Asn Ser Ser Asn Ala Ser Pro His 260 265 270

His Gln Tyr Asn Gln Phe Lys Thr Leu Leu Ala Asn Ser Ser Gln His 275 280 285

His His Gln Val Leu Asn Gln Phe Arg Ser Ser Pro Ala Ala Ser Ser 290 295 300

Ser Ser Met Ala Ala Val Asn Ile Leu Arg Asn Ser Arg Tyr Thr Thr 305 310 315 320

Ala Ala Gln Glu Leu Leu Glu Glu Phe Cys Ser Val Gly Arg Gly Phe 325 330 335

Leu Lys Lys Asn Lys Leu Gly Asn Ser Ser Asn Pro Asn Thr Cys Gly 340 345

MBI15 Sequence Listing.ST25
Gly Asp Gly Gly Ser Ser Pro Ser Ser Ala Gly Ala Asn Lys Glu
355 360 365

His Pro Pro Leu Ser Ala Ser Asp Arg Ile Glu His Gln Arg Arg Lys 370 380

Val Lys Leu Eu Thr Met Leu Glu Glu Val Asp Arg Tyr Asn His 385 390 395 400

Tyr Cys Glu Gln Met Gln Met Val Val Asn Ser Phe Asp Ile Val Met 405 410 415

Gly His Gly Ala Ala Leu Pro Tyr Thr Ala Leu Ala Gln Lys Ala Met 420 425 430

Ser Arg His Phe Arg Cys Leu Lys Asp Ala Val Ala Ala Gln Leu Lys 435 440 445

Gln Ser Cys Glu Leu Leu Gly Asp Lys Asp Ala Ala Gly Ile Ser Ser 450 455 460

Ser Gly Leu Thr Lys Gly Glu Thr Pro Arg Leu Arg Leu Leu Glu Gln 465 470 475 480

Ser Leu Arg Gln Gln Arg Ala Phe His Gln Met Gly Met Met Glu Gln 485 490 495

Glu Ala Trp Arg Pro Gln Arg Gly Leu Pro Glu Arg Ser Val Asn Ile 500 505 510

Leu Arg Ala Trp Leu Phe Glu His Phe Leu His Pro Tyr Pro Ser Asp 515 520 525

Ala Asp Lys His Leu Leu Ala Arg Gln Thr Gly Leu Ser Arg Asn Gln 530 535 540

Val Ser Asn Trp Phe Ile Asn Ala Arg Val Arg Leu Trp Lys Pro Met 545 550 560

Val Glu Glu Met Tyr Gln Gln Glu Ser Lys Glu Arg Glu Arg Glu Glu 565 570 575

Glu Leu Glu Glu Asn Glu Glu Asp Gln Glu Thr Lys Asn Ser Asn Asp 580 585 590

Asp Lys Ser Thr Lys Ser Asn Asn Glu Ser Asn Phe Thr Ala Val

Arg Thr Thr Ser Gln Thr Pro Thr Thr Thr Ala Pro Asp Ala Ser Asp

Ala Asp Ala Ala Val Ala Thr Gly His Arg Leu Arg Ser Asn Ile Asn 625 630 635

Ala Tyr Glu Asn Asp Ala Ser Ser Leu Leu Leu Pro Ser Ser Tyr Ser 645 655

PCT/US00/31418 WO 01/35726

MBI15 Sequence Listing.ST25

Asn Ala Ala Ala Pro Ala Ala Val Ser Asp Asp Leu Asn Ser Arg Tyr 665

Gly Gly Ser Asp Ala Phe Ser Ala Val Ala Thr Cys Gln Gln Ser Val

Gly Gly Phe Asp Asp Ala Asp Met Asp Gly Val Asn Val Ile Arg Phe

Gly Thr Asn Pro Thr Gly Asp Val Ser Leu Thr Leu Gly Leu Arg His 705 710 715 720

Ala Gly Asn Met Pro Asp Lys Asp Ala Ser Phe Cys Val Arg Glu Phe 730

Gly Gly Phe

<210> 19 <211> 1240 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS (88)..(1020) <222> <223> G591

<400> 19 60 qtaaatctct ctttgaaggt tcctaactcg ttaatcgtaa ctcacagtga ctcgttcgag tcaaagtctc tgtctttagc tcaaacc atg gct agt aac aac cct cac gac aac Met Ala Ser Asn Asn Pro His Asp Asn 114 ctt tct gac caa act cct tct gat gat ttc ttc gag caa atc ctc ggc Leu Ser Asp Gln Thr Pro Ser Asp Asp Phe Phe Glu Gln Ile Leu Gly 162 ctt cct aac ttc tca gcc tct tct gcc gcc ggt tta tct gga gtt gac Leu Pro Asn Phe Ser Ala Ser Ser Ala Ala Gly Leu Ser Gly Val Asp 210 gga gga tta ggt ggt gga gca ccg cct atg atg ctg cag ttg ggt tcc Gly Gly Leu Gly Gly Gly Ala Pro Pro Met Met Leu Gln Leu Gly Ser 45 50 55 258 gga gaa gaa gga agt cac atg ggt ggc tta gga gga agt gga cca act Gly Glu Glu Gly Ser His Met Gly Gly Leu Gly Gly Ser Gly Pro Thr 306 ggg ttt cac aat cag atg ttt cct ttg ggg tta agt ctt gat caa ggg Gly Phe His Asn Gln Met Phe Pro Leu Gly Leu Ser Leu Asp Gln Gly 354 aaa gga cct ggg ttt ctt aga cct gaa gga gga cat gga agt ggg aaa Lys Gly Pro Gly Phe Leu Arg Pro Glu Gly Gly His Gly Ser Gly Lys 402 aga ttc tca gat gat gtt gtt gat aat cga tgt tct tct atg aaa cct Arg Phe Ser Asp Asp Val Val Asp Asn Arg Cys Ser Ser Met Lys Pro 450 gtt ttc cac ggg cag cct atg caa cag cca cct cca tcg gcc cca cat Val Phe His Gly Gln Pro Met Gln Gln Pro Pro Pro Ser Ala Pro His 125 130 135

498

GRO GOT ROT TOO REAL COST	MBI15 Sequence	Listing.ST25	
cag cct act tca atc cgt ccc a Gln Pro Thr Ser Ile Arg Pro A	agg gtt ega get Arg Val Arg Ala	Arg Arg Gly Gln Ala	546
140	145	150	
act gat cca cat agc atc gct g	ag cgg cta cgt	aga gaa aga ata gca	594
Thr Asp Pro His Ser Ile Ala G	Blu Arg Leu Arg	Arg Glu Arg Ile Ala	
gaa cgg atc agg gcg ctg cag g Glu Arg Ile Arg Ala Leu Gln G			642
170 175	180		
gat aga gct gct atg atc gat g	ag att gtc gat	tat ota aag tit cic	690
Asp Arg Ala Ala Met Ile Asp G	lu Ile Val Asp	Tyr Val Lys Phe Leu	•
190	195	200	
agg ctc caa gtc aag gtt ttg a	gc atg aac cga	ctt ggt gga gcc ggt	738
Arg Leu Gln Val Lys Val Leu S 205	er met Asn Arg	Leu Gly Gly Ala Gly 215	
		h h h	206
gcg gtt gct cca ctt gtt act g Ala Val Ala Pro Leu Val Thr A			786
220 2	25	230	
gat gaa acg ggt gag ggt gga a	gg act ccg caa	cca gcg tgg gag aaa	834
Asp Glu Thr Gly Glu Gly Gly A 235 240	rg Thr Pro Gln	Pro Ala Trp Glu Lys 245	
tgg tct aac gat ggg act gaa c Trp Ser Asn Asp Gly Thr Glu A	gt caa gtg gct	aaa ctg atg gaa gag	882
250 255	260	265	
aac gtt gga gcc gcg atg cag c	tt ctt caa tca	aag oct ott tot atg	930
Asn Val Gly Ala Ala Met Gln L	eu Leu Gln Ser	Lys Ala Leu Cys Met	
270	275	280	
atg cca atc tca ttg gca atg g			978
Met Pro Ile Ser Leu Ala Met A 285	290	Ser Gin Pro Pro Asp	
			1000
aca tct tca gtg gtc aag cct g Thr Ser Ser Val Val Lys Pro G			1020
300 3	05	310	
gatttctgca ataaagagtt tgtacag	cta atccaactgt	ccaacatggg tttttcttct	1080
gctctaatga ctctggtttc ttctctc	ctc tctcaccgac	ttgaaaggta aaaaagtgaa	1140
aaaggctttg tagatggaat caatgta	gga tttgcagtag	agggcaaaaa aatgtcatat	1200
agctcaattg atcaagtctt aaaaaaa	aaa aaaaaaaaaa		1240
<210> 20			
<211> 310			
<pre><212> PRT <213> Arabidopsis thaliana</pre>			
<400> 20			
Met Ala Ser Asn Asn Pro His A	en Aen Leu Ser	Asp Gin Thr Pro Ser	

Met Ala Ser Asn Asn Pro His Asp Asn Leu Ser Asp Gln'Thr Pro Ser 1 10 15

Asp Asp Phe Phe Glu Gln Ile Leu Gly Leu Pro Asn Phe Ser Ala Ser 20 25 30

Ser Ala Ala Gly Leu Ser Gly Val Asp Gly Gly Leu Gly Gly Ala 35 40 45

Pro Pro Met Met Leu Gl
n Leu Gly Ser Gly Glu Glu Gly Ser His Met 50 $$

MBI15 Sequence Listing.ST25

Gly Gly Leu Gly Gly Ser Gly Pro Thr Gly Phe His Asn Gln Met Phe 65 70 75 80

Pro Leu Gly Leu Ser Leu Asp Gln Gly Lys Gly Pro Gly Phe Leu Arg

Pro Glu Gly Gly His Gly Ser Gly Lys Arg Phe Ser Asp Asp Val Val 100 105 110

Asp Asn Arg Cys Ser Ser Met Lys Pro Val Phe His Gly Gln Pro Met

Gln Gln Pro Pro Pro Ser Ala Pro His Gln Pro Thr Ser Ile Arg Pro

Arg Val Arg Ala Arg Gly Gln Ala Thr Asp Pro His Ser Ile Ala 145 150 155 160

Glu Arg Leu Arg Arg Glu Arg Ile Ala Glu Arg Ile Arg Ala Leu Glu 165 170 175

Glu Leu Val Pro Thr Val Asn Lys Thr Asp Arg Ala Ala Met Ile Asp 180 185 190

Glu Ile Val Asp Tyr Val Lys Phe Leu Arg Leu Gln Val Lys Val Leu 195 200 205

Ser Met Asn Arg Leu Gly Gly Ala Gly Ala Val Ala Pro Leu Val Thr 210 215 220

Asp Met Pro Leu Ser Ser Ser Val Glu Asp Glu Thr Gly Glu Gly Gly 225 230 235 240

Arg Thr Pro Gln Pro Ala Trp Glu Lys Trp Ser Asn Asp Gly Thr Glu 245 250 255

Arg Gln Val Ala Lys Leu Met Glu Glu Asn Val Gly Ala Ala Met Gln

Leu Leu Gln Ser Lys Ala Leu Cys Met Met Pro Ile Ser Leu Ala Met 275 280 285

Ala Ile Tyr His Ser Gln Pro Pro Asp Thr Ser Ser Val Val Lys Pro

Glu Asn Asn Pro Pro Gln

<210> 21 <211> 1179 <212> DNA

<213> Arabidopsis thaliana

<220>

CDS

<222> (109)..(966) <223> G525

MBI15 Sequence Listing.ST25

<400> 21 cttctctctt ctcaaaaacc cttccctctt cgtctccaaa caacaacaaa cacaacaaca 60																
acaaaaatct tacaagaaga tcatttttag aaaccctatt aggataaa atg gat tac 1: Met Asp Tyr 1												117				
														ata Ile		165
cta Leu 20	cca Pro	cca Pro	gga Gly	ttc Phe	aga Arg 25	ttt Phe	cac His	cct Pro	act Thr	gat Asp 30	gaa Glu	gaa Glu	ctc Leu	ata Ile	act Thr 35	213
cac His	tac Tyr	ctc Leu	aaa Lys	cca Pro 40	aag Lys	gtt Val	ttc Phe	aac Asn	act Thr 45	ttc Phe	ttc Phe	tct Ser	gct Ala	act Thr 50	gcc Ala	261
att Ile	ggt Gly	gaa Glu	gtt Val 55	gat Asp	ctc Leu	aac Asn	aag Lys	att Ile 60	gag Glu	cct Pro	tgg Trp	gac Asp	tta Leu 65	cca Pro	tgg Trp	309
aag Lys	gct Ala	aag Lys 70	atg Met	gga Gly	gaa Glu	aaa Lys	gaa Glu 75	tgg Trp	tat Tyr	ttc Phe	ttc Phe	tgt Cys 80	gtg Val	aga Arg	gac Asp	357
cgg Arg	aaa Lys 85	tac Tyr	ccg Pro	acc Thr	ggt Gly	tta Leu 90	agg Arg	aca Thr	aac Asn	cgg Arg	gcg Ala 95	aca Thr	gaa Glu	gcc Ala	ggt Gly	405
tat Tyr 100	tgg Trp	aaa Lys	gcc Ala	aca Thr	gga Gly 105	aaa Lys	gac Asp	aaa Lys	gag Glu	ata Ile 110	ttc Phe	aag Lys	gga Gly	aaa Lys	tca Ser 115	453
ctt Leu	gtg Val	ggt Gly	atg Met	aag Lys 120	aaa Lys	act Thr	ttg Leu	gtt Val	ttc Phe 125	tat Tyr	aaa Lys	gga Gly	aga Arg	gct Ala 130	cct Pro	501
aaa Lys	gga Gly	gtt Val	aaa Lys 135	acc Thr	aat Asn	tgg Trp	gtt Val	atg Met 140	cat His	gaa Glu	tat Tyr	cgt Arg	tta Leu 145	gaa Glu	ggc Gly	549
aaa Lys	tat Tyr	tgt Cys 150	att Ile	gaa Glu	aat Asn	ctt Leu	ccc Pro 155	caa Gln	aca Thr	gct Ala	aag Lys	aac Asn 160	gaa Glu	tgg Trp	gtt Val	597
ata Ile	tgt Cys 165	cgt Arg	gtt Val	ttc Phe	caa Gln	aaa Lys 170	cgt Arg	gcc Ala	gat Asp	ggt Gly	aca Thr 175	aag Lys	gtt Val	cca Pro	atg Met	645
tca Ser 180	atg Met	ctt Leu	gat Asp	cca Pro	cac His 185	att Ile	aac Asn	cga Arg	atg Met	gaa Glu 190	cca Pro	gcc Ala	ggt Gly	tta Leu	cct Pro 195	693
tcg Ser	tta Leu	atg Met	gat Asp	tgt Cys 200	tct Ser	caa Gln	cga Arg	gac Asp	tcc Ser 205	ttc Phe	acc Thr	ggt Gly	tcg Ser	tcg Ser 210	tct Ser	741
cac His	gtg Val	acc Thr	tgc Cys 215	ttc Phe	tcc Ser	gac Asp	caa Gln	gaa Glu 220	acc Thr	gaa Glu	gac Asp	aaa Lys	aga Arg 225	ctt Leu	gtc Val	789
cac His	gag Glu	tcc Ser 230	aaa Lys	gac Asp	ggt Gly	ttt Phe	ggt Gly 235	tct Ser	ctg Leu	ttt Phe	tac Tyr	tcg Ser 240	gat Asp	cct Pro	ctg Leu	837
ttt Phe	tta Leu 245	caa Gln	gac Asp	aat Asn	tat Tyr	tcg Ser 250	cta Leu	atg Met	aag Lys	ctg Leu	ttg Leu 255	ctt Leu	gac Asp	ggt Gly	caa Gln	885
gaa Glu 260	act Thr	caa Gln	ttc Phe	tcc Ser	ggc Gly 265	aaa Lys	cct Pro	ttc Phe	Asp	ggt Gly 270	Arg	gat Asp	tcg Ser	tcc Ser	ggt Gly 275	933

MBI15 Sequence Listing.ST25

aca gaa gaa ttg gat tgc gtt tgg aat ttc tga gttgtataag ttatgttgta

986
Thr Glu Glu Leu Asp Cys Val Trp Asn Phe
285

gacttgtagt agtcatgtgt tcgtgtgtgt gaatgaatat tcttgttaca tttttttgta

1046
aaaaaggaga aaaaaatatg ctagaaagtc aattgctttt gttatgtagc attagtgttt

1106
tttatgtact caatagactt cctaattaaa taaaaatctt aatttatttg ccaaaaaaaa

1179

<210> 22

<211> 285

<212> PRT

<213> Arabidopsis thaliana

<400> 22

Met Asp Tyr Glu Ala Ser Arg Ile Val Glu Met Val Glu Asp Glu Glu 1 5 15

His Ile Asp Leu Pro Pro Gly Phe Arg Phe His Pro Thr Asp Glu Glu 20 25 30

Leu Ile Thr His Tyr Leu Lys Pro Lys Val Phe Asn Thr Phe Phe Ser 35 40 45

Ala Thr Ala Ile Gly Glu Val Asp Leu Asn Lys Ile Glu Pro Trp Asp 50 55 60

Leu Pro Trp Lys Ala Lys Met Gly Glu Lys Glu Trp Tyr Phe Phe Cys 65 70 75 80

Val Arg Asp Arg Lys Tyr Pro Thr Gly Leu Arg Thr Asn Arg Ala Thr 85 90 95

Glu Ala Gly Tyr Trp Lys Ala Thr Gly Lys Asp Lys Glu Ile Phe Lys 100 105 110

Gly Lys Ser Leu Val Gly Met Lys Lys Thr Leu Val Phe Tyr Lys Gly

Arg Ala Pro Lys Gly Val Lys Thr Asn Trp Val Met His Glu Tyr Arg 130 135 140

Leu Glu Gly Lys Tyr Cys Ile Glu Asn Leu Pro Gln Thr Ala Lys Asn 145 150 150 160

Glu Trp Val Ile Cys Arg Val Phe Gln Lys Arg Ala Asp Gly Thr Lys 165 170 175

Val Pro Met Ser Met Leu Asp Pro His Ile Asn Arg Met Glu Pro Ala 180 185 190

Gly Leu Pro Ser Leu Met Asp Cys Ser Gln Arg Asp Ser Phe Thr Gly 195 200 205

Ser Ser Ser His Val Thr Cys Phe Ser Asp Gln Glu Thr Glu Asp Lys 210 215 220

MBI15 Sequence Listing.ST25

Arg 225		Val	His	Glu	Ser 230	Lys	Asp	Gly	Phe	Gly 235	Ser	Leu	Phe	Tyr	Ser 240	
Asp	Pro	Leu	Phe	Leu 245	Gln	Asp	Asn	Tyr	Ser 250	Leu	Met	Lys	Leu	Leu 255	Leu	
Asp	Gly	Gln	Glu 260	Thr	Gln	Phe	Ser	Gly 265	Lys	Pro	Phe	Asp	Gly 270	Arg	Asp	
Ser	Ser	Gly 275	Thr	Glu	Glu	Leu	Asp 280	Сув	Val	Trp	Asn	Phe 285				
<21 <21 <21 <21	1> 2>	23 890 DNA Arab:	idop	sis 1	thali	iana										
<22 <22 <22 <22	1 > (2 >	CDS (55) G545	(7:	38)												
<40 gca		23 tca a	aacta	aaaa	et c <u>e</u>	gagag	gacaa	a gaa	aatco	ctca	gaal	ctti	caa (ctta	atg Met 1	57
gcg Ala	ctc Leu	gag Glu	gct Ala 5	ctt Leu	aca Thr	tca Ser	cca Pro	aga Arg 10	tta Leu	gct Ala	tct Ser	ccg Pro	att Ile 15	cct Pro	cct Pro	105
ttg Leu	ttc Phe	gaa Glu 20	gat Asp	tct Ser	tca Ser	gtc Val	ttc Phe 25	cat His	gga Gly	gtc Val	gag Glu	cac His 30	tgg Trp	aca Thr	aag Lys	153
					aga Arg											201
act Thr 50	gag Glu	gaa Glu	gag Glu	tat Tyr	cta Leu 55	gct Ala	ttt Phe	tgc Cys	ctc Leu	atg Met 60	ctt Leu	ctc Leu	gct Ala	cgc Arg	gac Asp 65	249
aac Asn	cgt Arg	cag Gln	cct Pro	cct Pro 70	cct Pro	cct Pro	ccg Pro	gcg Ala	gtg Val 75	gag Glu	aag Lys	ttg Leu	agc Ser	tac Tyr 80	aag Lys	297
tgt Cys	agc Ser	gtc Val	tgc Cys 85	gac Asp	aag Lys	acg Thr	ttc Phe	tct Ser 90	tct Ser	tac Tyr	caa Gln	gct Ala	ctc Leu 95	ggt Gly	ggt Gly	345
					cgt Arg											393
gga Gly	gat Asp 115	gat Asp	cat His	tca Ser	acc Thr	tcg Ser 120	Ser	gcg Ala	aca Thr	acc Thr	aca Thr 125	tcc Ser	gcc Ala	gtg Val	act Thr	441
act Thr 130	gga Gly	agt Ser	999 Gly	aaa Lys	tca Ser 135	cac His	gtt Val	tgc Cys	acc Thr	atc Ile 140	tgt Cys	aac Asn	aag Lys	tct Ser	ttt Phe 145	489
cct Pro	tcc Ser	ggt Gly	caa Gln	gct Ala 150	ctc Leu	ggc Gly	gga Gly	cac His	aag Lys 155	cgg Arg	tgc Cys	cac His	tac Tyr	gaa Glu 160	gga Gly	537
aac	aac	aac	atc	aac	act	agt	agc	gtg	_	aac age		gaa	ggt	gcg	999	585

MBI15 Sequence Listing.ST25 Asn Asn Asn Tle Asn Thr Ser Ser Val Ser Asn Ser Glu Gly Ala Gly 165 170 175	
tcc act agc cac gtt agc agt agc cac cgt ggg ttt gac ctc aac atc Ser Thr Ser His Val Ser Ser Ser His Arg Gly Phe Asp Leu Asn Ile 180 185 190	633
cct ccg atc cct gaa ttc tcg atg gtc aac gga gac gac gaa gtc atg Pro Pro Ile Pro Glu Phe Ser Met Val Asn Gly Asp Asp Glu Val Met 195 200 205	681
agc cct atg ccg gcg aag aag cct cgg ttt gac ttt ccg gtc aaa ctt Ser Pro Met Pro Ala Lys Lys Pro Arg Phe Asp Phe Pro Val Lys Leu 210 215 220 225	729
caa ctt taa ggaaatttac ttagacgata agatttcgtt tgtatactgt Gln Leu	778
tgagagttgt gtaggaattt gttgactgta cataccaaat tggactttga ctgattccaa	838
ttettettgt tettteattt taaaaattat taaacegatt etttaceaca aa	890
<210> 24 <211> 227 <212> PRT <213> Arabidopsis thaliana	
<400> 24	
Met Ala Leu Glu Ala Leu Thr Ser Pro Arg Leu Ala Ser Pro Ile Pro 1 5 10 15	
Pro Leu Phe Glu Asp Ser Ser Val Phe His Gly Val Glu His Trp Thr 20 25 30	
Lys Gly Lys Arg Ser Lys Arg Ser Arg Ser Asp Phe His His Gln Asn 35 40 45	
Leu Thr Glu Glu Glu Tyr Leu Ala Phe Cys Leu Met Leu Leu Ala Arg 50 55 60	•
Asp Asn Arg Gln Pro Pro Pro Pro Pro Ala Val Glu Lys Leu Ser Tyr 75 80	
Lys Cys Ser Val Cys Asp Lys Thr Phe Ser Ser Tyr Gln Ala Leu Gly 85 90 95	
Gly His Lys Ala Ser His Arg Lys Asn Leu Ser Gln Thr Leu Ser Gly 100 105 110	
Gly Gly Asp Asp His Ser Thr Ser Ser Ala Thr Thr Thr Ser Ala Val	
Thr Thr Gly Ser Gly Lys Ser His Val Cys Thr Ile Cys Asn Lys Ser 130 135 140	
Phe Pro Ser Gly Gln Ala Leu Gly Gly His Lys Arg Cys His Tyr Glu 145 150 155 160	
Gly Asn Asn Asn Ile Asn Thr Ser Ser Val Ser Asn Ser Glu Gly Ala 165 170 175	

MBI15 Sequence Listing.ST25
Gly Ser Thr Ser His Val Ser Ser Ser His Arg Gly Phe Asp Leu Asn
180 185 190

Ile Pro Pro Ile Pro Glu Phe Ser Met Val Asn Gly Asp Asp Glu Val

Met Ser Pro Met Pro Ala Lys Lys Pro Arg Phe Asp Phe Pro Val Lys 210 215 220

Leu Gln Leu

<223> G865

<210> 25 <211> 1126 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (282)..(920)

gat aaa tcg gat caa cat caa cca gac caa gat caa cca aga aga 392 Asp Lys Ser Asp Gln His Gln Pro Asp Gln Asp Gln Pro Arg Arg 25 30 35

cac tat aga ggc gta agg cag aga cca tgg ggt aaa tgg gcg gca gaa
His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly Lys Trp Ala Ala Glu
40
45
440

atc cgc gat cca aag aaa gca gcc cgt gtc tgg ctc ggg act ttc gag 488
Ile Arg Asp Pro Lys Lys Ala Ala Arg Val Trp Leu Gly Thr Phe Glu
55 60 65

acg gca gag gaa gct gct tta gcc tat gac cga gct gcc ctc aaa ttc 536 Thr Ala Glu Glu Ala Ala Leu Ala Tyr Asp Arg Ala Ala Leu Lys Phe 70 85

aaa ggc acc aag gct aaa ctg aac ttc cct gaa cgg gtc caa ggc cct
Lys Gly Thr Lys Ala Lys Leu Asn Phe Pro Glu Arg Val Gln Gly Pro
90 95 100

act acc acc aca acc att tct cat gca cca aga gga gtt agt gaa tcc 632
Thr Thr Thr Thr Thr Ile Ser His Ala Pro Arg Gly Val Ser Glu Ser
105 110 115

atg aac tca cct cct cct cga cct ggt cca cct tca act act act Act Met Asn Ser Pro Pro Pro Arg Pro Gly Pro Pro Ser Thr Thr Thr 120 125 130

tcg tgg cca atg act tat aac cag gac ata ctt caa tac gct cag ttg

Ser Trp Pro Met Thr Tyr Asn Gln Asp Ile Leu Gln Tyr Ala Gln Leu

135

140

728

MBI15 Sequence Listing.ST25

	cg agt hr Ser												776
	gt caa er Gln		e Ser										824
	cg cag hr Gln												872
	ag aag Slu Lys 200				Tyr							taa	920
tctaai	ttatt a	attgttg	gtc ga	atcagt	t ta	taaai	tagc	tato	cata	gtt 1	catt	tttgg	980
tttcc	gtaac c	tttgtt	gca tg	gaaaata	it ga	atga	acga	9998	acat	gtg (aaca	aatttg	1040
tttgtg	gtttc g	gtaaatg	ta gt	tgtatt	g ga	tttg	ctga	agtt	tgai	ttt 1	ctga	agcata	1100
aatcat	tttga c	ggtcaa	aaa aa	aaaa									1126

<210> 26 <211> 212 <212> PRT <213> Arabidopsis thaliana

<400> 26

Met Val Ser Ala Leu Ser Arg Val Ile Glu Asn Pro Thr Asp Pro Pro 1 10 15

Val Lys Gln Glu Leu Asp Lys Ser Asp Gln His Gln Pro Asp Gln Asp 20 25 30

Gln Pro Arg Arg Arg His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly 35 40 45

Lys Trp Ala Ala Glu Ile Arg Asp Pro Lys Lys Ala Ala Arg Val Trp 50 60

Leu Gly Thr Phe Glu Thr Ala Glu Glu Ala Ala Leu Ala Tyr Asp Arg 65 70 75 80

Ala Ala Leu Lys Phe Lys Gly Thr Lys Ala Lys Leu Asn Phe Pro Glu 85 90 95

Arg Val Gln Gly Pro Thr Thr Thr Thr Thr Ile Ser His Ala Pro Arg

Gly Val Ser Glu Ser Met Asn Ser Pro Pro Pro Arg Pro Gly Pro Pro

Ser Thr Thr Thr Thr Ser Trp Pro Met Thr Tyr Asn Gln Asp Ile Leu 130 135 140

Gln Tyr Ala Gln Leu Leu Thr Ser Asn Asn Glu Val Asp Leu Ser Tyr

Tyr Thr Ser Thr Leu Phe Ser Gln Pro Phe Ser Thr Pro Ser Ser Ser 165 170 175

MBI15 Sequence Listing.ST25

Ser Ser Ser Ser Gln Gln Thr Gln Gln Gln Gln Leu Gln Gln Gln

Gln Gln Arg Glu Glu Glu Lys Asn Tyr Gly Tyr Asn Tyr Tyr Asn

Tyr Pro Arg Glu 210

<210> 27 <211> 1152 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (76)..(1008) <223> G881 <400> 27 60 gtgaccgaag aaagcaaatt gagactacgc accaactagt cetttggttt gtatettaag ataaaggttt ctttt atg gac ggt tct tcg ttt ctc gac atc tct ctc gat Met Asp Gly Ser Ser Phe Leu Asp Ile Ser Leu Asp 111 ctc aac acc aat cct ttc tcc gca aaa ctt ccg aag aag gag gtc tca Leu Asn Thr Asn Pro Phe Ser Ala Lys Leu Pro Lys Lys Glu Val Ser 159 20 207 gtt ttg gct tct act cac tta aag agg aaa tgg ttg gag caa gac gag Val Leu Ala Ser Thr His Leu Lys Arg Lys Trp Leu Glu Gln Asp Glu agc gca agt gag tta cga gag gag cta aac aga gtt aat tca gag aac Ser Ala Ser Glu Leu Arg Glu Glu Leu Asn Arg Val Asn Ser Glu Asn 255 303 aag aag cta aca gag atg tta gct aga gtc tgt gag agc tac aac gaa Lys Lys Leu Thr Glu Met Leu Ala Arg Val Cys Glu Ser Tyr Asn Glu 65 70 75cta cat aat cat ttg gag aag ctt cag agt cgc cag agc cct gaa atc Leu His Asn His Leu Glu Lys Leu Gln Ser Arg Gln Ser Pro Glu Ile 351 gag cag acc gat ata ccg ata aag aaa aga aaa caa gac ccg gat gag Glu Gln Thr Asp Ile Pro Ile Lys Lys Arg Lys Gln Asp Pro Asp Glu 399 ttc tta ggc ttt cct att gga ctc agt agt gga aaa act gag aac agc Phe Leu Gly Phe Pro Ile Gly Leu Ser Ser Gly Lys Thr Glu Asn Ser 447 495 tcc agc aac gaa gat cat cat cat cat cat cag caa cat gag cag aaa Ser Ser Asn Glu Asp His His His His His Gln Gln His Glu Gln Lys aat cag ctt ctt tca tgt aaa aga cca gtc act gat agc ttc aac aaa Asn Gln Leu Leu Ser Cys Lys Arg Pro Val Thr Asp Ser Phe Asn Lys 543 gca aaa gtt tcg act gtc tac gtg cct act gaa aca tcg gac aca agc Ala Lys Val Ser Thr Val Tyr Val Pro Thr Glu Thr Ser Asp Thr Ser 591 165 ttg aca gtt aaa gat gga ttt caa tgg agg aaa tac gga caa aag gtt Leu Thr Val Lys Asp Gly Phe Gln Trp Arg Lys Tyr Gly Gln Lys Val 639

Page 36

180

aca aga q Thr Arg A 190	gac aac Asp Asn	ccg tca Pro Ser	cct aga Pro Arg 195	gct (Ala :	tac ttt Tyr Phe	aga to Arg Co 200	gc tcg ys Ser	ttt Phe	gca Ala	687
ccg tct t Pro Ser (205										735
tct tta d Ser Leu I	Leu Val	gcg aca Ala Thr 225	tac gaa Tyr Glu	Gly :	acg cat Thr His 230	aac ca Asn H	ac ttg is Leu	ggt Gly 235	cca Pro	783
aat gct t Asn Ala S										831
act ttg o						Ala Le				879
gaa agg g Glu Arg A 270	gat aat Asp Asn	acg atg Thr Met	caa gag Gln Glu 275	gtt o	ctg att Leu Ile	caa ca Gln Gl 280	aa atg ln Met	gcg Ala	tca Ser	927
tcg tta a Ser Leu T 285										975
tct ggg a Ser Gly A	Arg Leu			Arg 7		acgttt	tttag t	gaat	gtatt	1028
gtttttgtt	t gttta	gaatg at	tettegt	t ttcg	gaattgt	gtcttt	cgat t	agga	gataa	1088
aagatgtat	ta taaat	attat aa	ıgtagatgı	a agaa	aatcgta	taagta	aaaaa a	aaaa	aaaaa	1148
aaaa										1152

Met Asp Gly Ser Ser Phe Leu Asp Ile Ser Leu Asp Leu Asn Thr Asn 1 $$ 15

Pro Phe Ser Ala Lys Leu Pro Lys Lys Glu Val Ser Val Leu Ala Ser 20 30

Thr His Leu Lys Arg Lys Trp Leu Glu Gln Asp Glu Ser Ala Ser Glu 35 40 45

Leu Arg Glu Glu Leu Asn Arg Val Asn Ser Glu Asn Lys Lys Leu Thr 50 60

Glu Met Leu Ala Arg Val Cys Glu Ser Tyr Asn Glu Leu His Asn His 65 70 75 80

Leu Glu Lys Leu Gln Ser Arg Gln Ser Pro Glu Ile Glu Gln Thr Asp

Ile Pro Ile Lys Lys Arg Lys Gln Asp Pro Asp Glu Phe Leu Gly Phe 100 \$105\$

<210> 28 <211> 310 <212> PRT <213> Arabidopsis thaliana

MBI15 Sequence Listing.ST25 Pro Ile Gly Leu Ser Ser Gly Lys Thr Glu Asn Ser Ser Ser Asn Glu 115 120 125 Asp His His His His Gln Gln His Glu Gln Lys Asn Gln Leu Leu Ser Cys Lys Arg Pro Val Thr Asp Ser Phe Asn Lys Ala Lys Val Ser Thr Val Tyr Val Pro Thr Glu Thr Ser Asp Thr Ser Leu Thr Val Lys Asp Gly Phe Gln Trp Arg Lys Tyr Gly Gln Lys Val Thr Arg Asp Asn 180 \$190\$Pro Ser Pro Arg Ala Tyr Phe Arg Cys Ser Phe Ala Pro Ser Cys Pro Val Lys Lys Lys Val Gln Arg Ser Ala Glu Asp Pro Ser Leu Leu Val Ala Thr Tyr Glu Gly Thr His Asn His Leu Gly Pro Asn Ala Ser Glu 225 230 230 235 Gly Asp Ala Thr Ser Gln Gly Gly Ser Ser Thr Val Thr Leu Asp Leu 245 250 255 Val Asn Gly Cys His Arg Leu Ala Leu Glu Lys Asn Glu Arg Asp Asn Thr Met Gln Glu Val Leu Ile Gln Gln Met Ala Ser Ser Leu Thr Lys Asp Ser Lys Phe Thr Ala Ala Leu Ala Ala Ala Ile Ser Gly Arg Leu Met Glu Gln Ser Arg Thr <210> 29 <211> 1276 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (47)..(1150) <400> 29 taatccgatt cgtcttcatc tgattccctc ccttccgaga ataata atg tac ccg Met Tyr Pro cca cct ccc tca agc atc tac gct cct ccg atg ctg gtg aat tgc tcc
Pro Pro Pro Ser Ser Ile Tyr Ala Pro Pro Met Leu Val Asn Cys Ser
5
10 103 ggt tgc cgg acg cct ctc cag ctc cca tcc ggc gcc cga tct att cgc Gly Cys Arg Thr Pro Leu Gln Leu Pro Ser Gly Ala Arg Ser Ile Arg 20 25 30 35 151

MBI15 Sequence Listing.ST25

									-			-				
tgc Cys	gct Ala	ctc Leu	tgc Cys	cag Gln 40	gct Ala	gtt Val	act Thr	cat His	atc Ile 45	gcc Ala	gac Asp	cct Pro	cgc Arg	acc Thr 50	gcc Ala	199
cct Pro	cct Pro	ccg Pro	caa Gln 55	cct Pro	tcc Ser	tcc Ser	gcc Ala	cct Pro 60	tct Ser	ccg Pro	cct Pro	ccc Pro	caa Gln 65	atc Ile	cac His	247
gcg Ala	cct Pro	ccc Pro 70	ggt Gly	cag Gln	ctg Leu	cct Pro	cac His 75	ccc Pro	cat His	ggc Gly	agg Arg	aag Lys 80	agg Arg	gcc Ala	gtg Val	295
atc Ile	tgt Cys 85	ggc Gly	atc Ile	tcg Ser	tat Tyr	cgt Arg 90	ttc Phe	tct Ser	cgc Arg	cac His	gag Glu 95	ctc Leu	aaa Lys	ggc Gly	tgc Cys	343
atc Ile 100	aac Asn	gac Asp	gcc Ala	aag Lys	tgc Cys 105	atg Met	cgt Arg	cac His	ctt Leu	ctc Leu 110	atc Ile	aac Asn	aaa Lys	ttc Phe	aaa Lys 115	391
ttc Phe	tcc Ser	cca Pro	gat Asp	tca Ser 120	att Ile	ctc Leu	atg Met	ctt Leu	acc Thr 125	gag Glu	gaa Glu	gaa Glu	act Thr	gat Asp 130	cca Pro	439
tat Tyr	cgt Arg	atc Ile	ccg Pro 135	acc Thr	aag Lys	caa Gln	aac Asn	atg Met 140	agg Arg	atg Met	gca Ala	ttg Leu	tat Tyr 145	tgg Trp	ctc Leu	487
gta Val	cag Gln	gga Gly 150	tgc Cys	aca Thr	gca Ala	ggc Gly	gac Asp 155	tca Ser	ctt Leu	gtc Val	ttc Phe	cac His 160	tac Tyr	tct Ser	ggt Gly	535
cat His	ggt Gly 165	tcg Ser	cgt Arg	caa Gln	aga Arg	aac Asn 170	tac Tyr	aac Asn	ggt Gly	gat Asp	gaa Glu 175	gtt Val	gat Asp	ggc Gly	tat Tyr	583
gat Asp 180	gaa Glu	aca Thr	ctc Leu	tgt Cys	cct Pro 185	ctg Leu	gat Asp	ttt Phe	gaa Glu	act Thr 190	cag Gln	999 Gly	atg Met	att Ile	gta Val 195	631
gac Asp	gat Asp	gag Glu	atc Ile	aac Asn 200	gca Ala	acc Thr	att Ile	gta Val	cgc Arg 205	cct Pro	ctt Leu	cca Pro	cat His	ggt Gly 210	gtc Val	679
aag Lys	ctc Leu	cat His	tca Ser 215	att Ile	atc Ile	gat Asp	gct Ala	tgc Cys 220	cat His	agt Ser	ggt Gly	acc Thr	gtt Val 225	ctg Leu	gat Asp	727
tta Leu	ccc Pro	ttc Phe 230	cta Leu	tgc Cys	aga Arg	atg Met	aac Asn 235	aga Arg	gct Ala	999 Gly	cag Gln	tat Tyr 240	gtg Val	tgg Trp	gag Glu	775
gat Asp	cat His 245	cgg Arg	cct Pro	agg Arg	tca Ser	ggt Gly 250	ttg Leu	tgg Trp	aaa Lys	gga Gly	act Thr 255	gct Ala	ggt Gly	gga Gly	gaa Glu	823
gcc Ala 260	att Ile	tca Ser	att Ile	agt Ser	gga Gly 265	tgt Cys	gat Asp	gat Asp	gat Asp	cag Gln 270	act Thr	tcg Ser	gcc Ala	gac Asp	aca Thr 275	871
tca Ser	gcg Ala	ctg Leu	tcg Ser	aag Lys 280	atc Ile	acg Thr	tct Ser	acg Thr	ggt Gly 285	gct Ala	atg Met	act Thr	ttc Phe	tgt Cys 290	ttt Phe	919
att Ile	caa Gln	gca Ala	att Ile 295	gaa Glu	cgc Arg	agc Ser	gca Ala	caa Gln 300	ggc Gly	aca Thr	acc Thr	tat Tyr	gga Gly 305	agc Ser	ctt Leu	967
ctg Leu	aat Asn	tct Ser 310	atg Met	cgc Arg	acc Thr	aca Thr	ata Ile 315	agg Arg	aat Asn	aca Thr	G1y 999	aat Asn 320	gat Asp	ggt Gly	ggt Gly	1015
ggt Gly	agt Ser	ggt Gly	gga Gly	gtt Val	gtg Val	acg Thr	act Thr	gtg Val	Leu	agc Ser	Met	ctt Leu	ctg Leu	aca Thr	999 Gly	1063

1111

1160

1220

1276

MBI15 Sequence Listing.ST25 gga agt gcg att ggg gga tta aga cag gag cct caa ctg act gct tgc Gly Ser Ala Ile Gly Gly Leu Arg Gln Glu Pro Gln Leu Thr Ala Cys 340 caa aca ttc gat gtc tat gca aag cct ttc act ctc tag taaaggacaa Gln Thr Phe Asp Val Tyr Ala Lys Pro Phe Thr Leu gtcacttttt atgtatagcg agtgtgattt gagaatccgt ccatataacc accttttgtt tottattttt atttttcttt caaaagaata aaggaaaaca ttgatttggt gattcg <210> 30 <211> 367 <212> PRT <213> Arabidopsis thaliana <400> 30 Met Tyr Pro Pro Pro Pro Ser Ser Ile Tyr Ala Pro Pro Met Leu Val Asn Cys Ser Gly Cys Arg Thr Pro Leu Gln Leu Pro Ser Gly Ala Arg Ser Ile Arg Cys Ala Leu Cys Gln Ala Val Thr His Ile Ala Asp Pro 35 40 45 Arg Thr Ala Pro Pro Pro Gln Pro Ser Ser Ala Pro Ser Pro Pro Pro Gln Ile His Ala Pro Pro Gly Gln Leu Pro His Pro His Gly Arg Lys Arg Ala Val Ile Cys Gly Ile Ser Tyr Arg Phe Ser Arg His Glu Leu 85 90 95 Lys Gly Cys Ile Asn Asp Ala Lys Cys Met Arg His Leu Leu Ile Asn 100 105 110 Lys Phe Lys Phe Ser Pro Asp Ser Ile Leu Met Leu Thr Glu Glu Glu 115 120 125 Thr Asp Pro Tyr Arg Ile Pro Thr Lys Gln Asn Met Arg Met Ala Leu Tyr Trp Leu Val Gln Gly Cys Thr Ala Gly Asp Ser Leu Val Phe His 145 150 155 160 Tyr Ser Gly His Gly Ser Arg Gln Arg Asn Tyr Asn Gly Asp Glu Val Asp Gly Tyr Asp Glu Thr Leu Cys Pro Leu Asp Phe Glu Thr Gln Gly

Met Ile Val Asp Asp Glu Ile Asn Ala Thr Ile Val Arg Pro Leu Pro 195 200 205

His Gly Val Lys Leu His Ser Ile Ile Asp Ala Cys His Ser Gly Thr Page 40

MBI15 Sequence Listing.ST25

210 Val Leu Asp Leu Pro Phe Leu Cys Arg Met Asn Arg Ala Gly Gln Tyr 230 Val Trp Glu Asp His Arg Pro Arg Ser Gly Leu Trp Lys Gly Thr Ala Gly Gly Glu Ala Ile Ser Ile Ser Gly Cys Asp Asp Asp Gln Thr Ser 260 265 270 Ala Asp Thr Ser Ala Leu Ser Lys Ile Thr Ser Thr Gly Ala Met Thr Phe Cys Phe Ile Gln Ala Ile Glu Arg Ser Ala Gln Gly Thr Thr Tyr Gly Ser Leu Leu Asn Ser Met Arg Thr Thr Ile Arg Asn Thr Gly Asn Asp Gly Gly Ser Gly Gly Val Val Thr Thr Val Leu Ser Met Leu Leu Thr Gly Gly Ser Ala Ile Gly Gly Leu Arg Gln Glu Pro Gln Leu Thr Ala Cys Gln Thr Phe Asp Val Tyr Ala Lys Pro Phe Thr Leu <210> <211> 31 726 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (1)..(726) <223> G378 <400> 31 atg gcc tcg tca tca tca tct tat aga ttc caa tct ggg tct tac Met Ala Ser Ser Ser Ser Ser Tyr Arg Phe Gln Ser Gly Ser Tyr 48 cct ctt tcg tca agt cct tct ctt ggg aat ttc gtc gaa cgc att aaa Pro Leu Ser Ser Pro Ser Leu Gly Asn Phe Val Glu Arg Ile Lys 20 25 30 96 gac gct tgt cat ttc ctt gtc tct gct gtt ttg ggt acc att atc tcc Asp Ala Cys His Phe Leu Val Ser Ala Val Leu Gly Thr Ile Ile Ser 40 45144 gcg atc ttg acc ttc ttc ttc gca cta gtg ggc aca ttg cta ggg gca Ala Ile Leu Thr Phe Phe Ala Leu Val Gly Thr Leu Leu Gly Ala 192 ctt aca gga gct ttg ata ggt caa gaa act gag agt ggt ttc att aga Leu Thr Gly Ala Leu Ile Gly Gln Glu Thr Glu Ser Gly Phe Ile Arg 65 70 75 80 240 gga gca gca att gga gcc att tcg gga gct gtt ttc tct atc gag gtc Gly Ala Ala Ile Gly Ala Ile Ser Gly Ala Val Phe Ser Ile Glu Val 288

ttt	gaa	tca	tct	ctg	gat	ctc	tgg	15 S aaa	tcc	qat	qaq	tcq	qqt	ttc	gga		336
Phe	Ğlu	Ser	Ser 100	Leu	Āsp	Leu	Trp	Lys 105	Ser	Āsp	Ğlü	Ser	Gly 110	Phe	Ğİy		
				ttg Leu													384
				cgc Arg													432
				gtg Val													480
gat Asp	aca Thr	gga Gly	ggc Gly	tca Ser 165	aaa Lys	gga Gly	ttg Leu	aca Thr	gga Gly 170	gac Asp	ctt Leu	gtt Val	gag Glu	aaa Lys 175	atc Ile		528
				atc Ile													576
aca Thr	gac Asp	tca Ser 195	tgt Cys	tct Ser	gtt Val	tgt Cys	ctt Leu' 200	cag Gln	gat Asp	ttc Phe	cag Gln	ctc Leu 205	ggt Gly	gaa Glu	aca Thr		624
				cct Pro													672
				ctt Leu													720
att Ile	taa																726
<210 <211 <212 <213	> 2 > I	2 241 RT	dons	sis t	hali	iana											
<400		2	. Lope														
Met 1	Ala	Ser	Ser	Ser 5	Ser	Ser	Ser	Tyr	Arg 10	Phe	Gln	Ser	Gly	Ser 15	Tyr	•	
Pro	Leu	Ser	Ser 20	Ser	Pro	Ser	Leu	Gly 25	Asn	Phe	Val	Glu	Arg 30	Ile	Lys		
Asp	Ala	Сув 35	His	Phe	Leu	Val	Ser 40	Ala	Val	Leu	Gly	Thr 45	Ile	Ile	Ser		
Ala	Ile 50	Leu	Thr	Phe	Phe	Phe 55	Ala	Leu	Val	Gly	Thr 60	Leu	Leu	Gly	Ala		
Leu 65	Thr	Gly	Ala	Leu	Ile 70	Gly	Gln	Glu	Thr	Glu 75	Ser	Gly	Phe	Ile	Arg 80		
Gly	Ala	Ala	Ile	Gly 85	Ala	Ile	Ser	Gly	Ala 90	Val	Phe	Ser	Ile	Glu 95	Val		

Page 42

Phe Glu Ser Ser Leu Asp Leu Trp Lys Ser Asp Glu Ser Gly Phe Gly 100 105 110

PCT/US00/31418 WO 01/35726

MBI15 Sequence Listing.ST25

Cys Phe Leu Tyr Leu Ile Asp Val Ile Val Ser Leu Leu Ser Gly Arg

Leu Val Arg Glu Arg Ile Gly Pro Ala Met Leu Ser Ala Val Gln Ser

Gln Met Gly Ala Val Asp Thr Ala Phe Asp Asp His Thr Ser Leu Phe

Asp Thr Gly Gly Ser Lys Gly Leu Thr Gly Asp Leu Val Glu Lys Ile

Pro Lys Met Thr Ile Thr Gly Asn Asn Asn Thr Asp Ala Ser Glu Asn

Thr Asp Ser Cys Ser Val Cys Leu Gln Asp Phe Gln Leu Gly Glu Thr

Val Arg Ser Leu Pro His Cys His His Met Phe His Leu Pro Cys Ile

Asp Asn Trp Leu Leu Arg His Gly Ser Cys Pro Met Cys Arg Arg Asp

Ile

<210> 33 <211> 1370 <212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (184)..(969)

<223> G569

gtcgacccac gcgtccgggt ttttctttta tcctcttatc gctaatctgg agctctatat 60 agactataaa gggtttttga ttgattcggg agctcgagat ttgacttctt ttagctgatt 120 cggcaagttt gtatctagaa aggatcgatt ggtgaggtca atagtggttg gtgggtttta gta atg gaa gac ggt gag ctt gat ttc tcc aat cag gaa gtg ttt tcg Met Glu Asp Gly Glu Leu Asp Phe Ser Asn Gln Glu Val Phe Ser 1 5 10 228 agt tcg gag atg ggt gaa tta cca cct agc aat tgt tcg atg gat agt Ser Ser Glu Met Gly Glu Leu Pro Pro Ser Asn Cys Ser Met Asp Ser 20 25 30276 ttc ttt gat ggg ctt tta atg gat act aat gct gct tgt acc cac act Phe Phe Asp Gly Leu Leu Met Asp Thr Asn Ala Ala Cys Thr His Thr 324 40 cac acc tgt aac ccc act gga cca gag aac act cat act cac acg tgc His Thr Cys Asn Pro Thr Gly Pro Glu Asn Thr His Thr His Thr Cys 372

ttc cat gtc cac acc aag att ctc ccg gat gag agc gat gaa aaa gtt Phe His Val His Thr Lys Ile Leu Pro Asp Glu Ser Asp Glu Lys Val

420

							MBI	15 S	eque	nce	List	ing.	ST25				
tct Ser 80	act Thr	gat Asp	gat Asp	aca Thr	gct Ala 85	gag Glu	tct	tgt	999	aag	aag	ggt	gaa	aag	aga Arg 95	468	
cct Pro	ttg Leu	gga Gly	aac Asn	cgg Arg 100	gaa Glu	gcg Ala	gtt Val	aga Arg	aag Lys 105	tat Tyr	aga Arg	gag Glu	aag Lys	aag Lys 110	aag Lys	516	•
				tct Ser												564	
aat Asn	cag Gln	cag Gln 130	ctg Leu	gtg Val	aag Lys	agg Arg	ttg Leu 135	caa Gln	aat Asn	cag Gln	gct Ala	acc Thr 140	ttg Leu	gaa Glu	gct Ala	612	
gag Glu	gtt Val 145	tcg Ser	agg Arg	ctt Leu	aag Lys	tgt Cys 150	ttg Leu	ctt Lėu	gtg Val	gat Asp	ttg Leu 155	aga Arg	gga Gly	aga Arg	ata Ile	660	
				gga Gly												708	
att Ile	cct Pro	tct Ser	ttc Phe	tcg Ser 180	cac His	atg Met	atg Met	aat Asn	cct Pro 185	tgt Cys	aat Asn	gta Val	caa Gln	tgt Сув 190	gat Asp	756	
gat Asp	gaa Glu	gtt Val	tat Tyr 195	tgc Cys	cct Pro	cag Gln	aat Asn	gtg Val 200	ttt Phe	gga Gly	gtg Val	aat Asn	agc Ser 205	caa Gln	gaa Glu	804	
				aat Asn												852	
cta Leu	caa Gln 225	tgc Cys	atg Met	gct Ala	aat Asn	cag Gln 230	aac Asn	tta Leu	aat Asn	gga Gly	aat Asn 235	gga Gly	aac Asn	gga Gly	tca Ser	900	
ttc Phe 240	agc Ser	aac Asn	gtc Val	aat Asn	aca Thr 245	tct Ser	gtc Val	tcg Ser	aat Asn	aag Lys 250	aga Arg	aaa Lys	ggt Gly	999 Gly	cat His 255	948	
				gca Ala 260		tga	agca	tcat	ca a	igctt	gtac	t at	ctat	ttcc	!	999	
acca	gcat	ag a	tatt	gtat	t cc	aaat	aagt	tgt	agag	ttc	agct	gcag	ga t	cago	ttcgc	1059	
tcag	cttt	ga g	gggt	tggt	g gt	gtgg	tctt	tct	ttgt	ggc	acga	gtga	iga t	ctat	ggaca	1119	
gaac	ccag	at t	tagt	agta	g ta	gagg	cagg	att	tcga	ctt	ccac	taac	ca t	catg	ttgct	1179	
tggt	gaag	aa c	aagg	tatg	c cc	atga	agca	cac	tgtt	ttg	taca	ttga	gc t	tgag	gggct	1239	
gtct	ctga	tc t	agco	ttac	t gt	aaca	ttgc	aac	gtto	tca	caat	tgtg	at c	ccaa	gttgc	1299	
tttg	ttga	ct t	aaat	gtga	t aa	tata	gctt	aac	tttt	act	tgaa	aaaa	aa a	aaaa	aaaaa	1359	
aaaa	aaaa	aa a	ı		-											1370	

Met Glu Asp Gly Glu Leu Asp Phe Ser Asn Gln Glu Val Phe Ser Ser 1 10 15

Ser Glu Met Gly Glu Leu Pro Pro Ser Asn Cys Ser Met Asp Ser Phe Page 44

<210> 34 <211> 261 <212> PRT <213> Arabidopsis thaliana

MBI15 Sequence Listing.ST25 25 30

20

Phe Asp Gly Leu Leu Met Asp Thr Asn Ala Ala Cys Thr His Thr His 35 40 45

Thr Cys Asn Pro Thr Gly Pro Glu Asn Thr His Thr His Thr Cys Phe

His Val His Thr Lys Ile Leu Pro Asp Glu Ser Asp Glu Lys Val Ser 65 70 75 80

Thr Asp Asp Thr Ala Glu Ser Cys Gly Lys Lys Gly Glu Lys Arg Pro

Leu Gly Asn Arg Glu Ala Val Arg Lys Tyr Arg Glu Lys Lys Lys Ala

Lys Ala Ala Ser Leu Glu Asp Glu Val Ala Arg Leu Arg Ala Val Asn 115 120 125

Gln Gln Leu Val Lys Arg Leu Gln Asn Gln Ala Thr Leu Glu Ala Glu

Val Ser Arg Leu Lys Cys Leu Leu Val Asp Leu Arg Gly Arg Ile Asp 145 150 150 160

Gly Glu Ile Gly Ser Phe Pro Tyr Gln Lys Pro Met Ala Ala Asn Ile 165 170 175

Pro Ser Phe Ser His Met Met Asn Pro Cys Asn Val Gln Cys Asp Asp 180 185 190

Glu Val Tyr Cys Pro Gln Asn Val Phe Gly Val Asn Ser Gln Glu Gly 195 200 205

Ala Ser Ile Asn Asp Gln Gly Leu Ser Gly Cys Asp Phe Asp Gln Leu 210 215 220

Gln Cys Met Ala Asn Gln Asn Leu Asn Gly Asn Gly Asn Gly Ser Phe

Ser Asn Val Asn Thr Ser Val Ser Asn Lys Arg Lys Gly Gly His Arg 245 250 255

Ala Ser Arg Ala Val 260

<210> 35 <211> 1638 <212> DNA <213> Arabidopsis thaliana

<221> CDS

<222> (267)..(1259) <223> G558

<400> 35

	I15 Sequence Listing.ST25
	tt ttccagcgcc cagaatgctc cgggtgacca 120
	gt tcgtattgtc tgtgcacaca agagaaattt 180
caagaagttg ttactaaaag agaggccac	ca agtggatatt gtctttgtta tcaagtgtta 240
	ng gct gat acc agt ccg aga act gat 293 et Ala Asp Thr Ser Pro Arg Thr Asp 5
	c cat cct gat ctt ggg tcg gag gga 341 o His Pro Asp Leu Gly Ser Glu Gly 20 25
	gat tog agt gac cga tog aag gga 389 Asp Ser Ser Asp Arg Ser Lys Gly 35 40
	agg ctt gct caa aac cgt gag gca 437 Arg Leu Ala Gln Asn Arg Glu Ala 50 55
	g aag gct tat gtt cag cag cta gag 485 3 Lys Ala Tyr Val Gln Gln Leu Glu 70
	g ctt gag cag gag ctg caa aga gca 533 Leu Glu Gln Glu Leu Gln Arg Ala 85
	ggc aca gga gac cag gcc cat tct 581 Gly Thr Gly Asp Gln Ala His Ser 100 105
	ttt gat gct gaa cat tca cgg tgg 629 Phe Asp Ala Glu His Ser Arg Trp 115 120
	aac gag ctg agg tct gct ctg aat 677 Asn Glu Leu Arg Ser Ala Leu Asn 130 135
	cga ata ata gtc gat ggt gtg atg 725 Arg Ile Ile Val Asp Gly Val Met 150
	ata aag agc aat gca gct aag aat 773 Ile Lys Ser Asn Ala Ala Lys Asn 165
	e atg tgg aaa aca cca gct gag aga 821 Met Trp Lys Thr Pro Ala Glu Arg 180 185
	e cgt tca tcc gaa ctt cta aag ctt 869 Arg Ser Ser Glu Leu Leu Lys Leu 195 200
	aca gag aga cag ttg atg ggc ata 917 Thr Glu Arg Gln Leu Met Gly Ile 210 215
	cag gct gaa gat gct ttg tct caa 965 Gln Ala Glu Asp Ala Leu Ser Gln 230
	cta gct gat act tta tcg agc ggg 1013 Leu Ala Asp Thr Leu Ser Ser Gly 245
	aat gtc gca agc tac atg ggt cag 1061 Asn Val Ala Ser Tyr Met Gly Gln 260 265

Page 46

MBI15 Sequence Listing.ST25

atg (Met i																1109
cag g																1157
tta a Leu :																1205
ttc (Phe s																1253
gag (Glu 330	tga	aact	gtat	ett t	ggto	cacat	g to	agct	gtad	aaa	atco	ata	tgga	acaca	ıaa	1309
accag	ggag	ag a	ctat	taat	c aa	acact	tgto	aga	ttct	tct	taco	aaat	cc a	atcaa	caaat	1369
aagca	aaat	tt c	tggg	gaaac	a aa	agad	tctt	: tgt	atgt	agg	tttc	ettet	ac a	tggt	tgtgg	1429
taatt	tcat	gt t	gttt	tagt	t gt	agto	atca	gtt	ttta	att	tago	attt	ga a	aagt	tcaat	1489
gttgt	ttta	ta t	agca	tctt	c ga	ittat	ctta	gaa	aggt	tat	tgaa	ittt	gt t	tttt	tttgt	1549
tact	tttg	tg t	gtgg	jtaaa	ig gt	gttt	taac	ctt	gcaa	ctt	ctgt	acto	jta a	tcat	ttaac	1609
aatat	ttaa	ga t	gtto	tatt	t ga	gttt	tgt									1638
<210:	> 3	6														

<211> 330
<212> PRT
<213> Arabidopsis thaliana

<400> 36

Met Ala Asp Thr Ser Pro Arg Thr Asp Val Ser Thr Asp Asp Asp Thr 1 10 15

Asp His Pro Asp Leu Gly Ser Glu Gly Ala Leu Val Asn Thr Ala Ala 20 25 30

Ser Asp Ser Ser Asp Arg Ser Lys Gly Lys Met Asp Gln Lys Thr Leu 35 40 45

Arg Arg Leu Ala Gln Asn Arg Glu Ala Ala Arg Lys Ser Arg Leu Arg 50 55 60

Lys Lys Ala Tyr Val Gln Gln Leu Glu Asn Ser Arg Leu Lys Leu Thr 65 70 75

Gln Leu Glu Gln Glu Leu Gln Arg Ala Arg Gln Gln Gly Val Phe Ile 85 90 95

Ser Gly Thr Gly Asp Gln Ala His Ser Thr Gly Gly Asn Gly Ala Leu $100 \hspace{1cm} 105 \hspace{1cm} 110$

Ala Phe Asp Ala Glu His Ser Arg Trp Leu Glu Glu Lys Asn Lys Gln 115 120 125

Met Asn Glu Leu Arg Ser Ala Leu Asn Ala His Ala Gly Asp Ser Glu 130 135 140

MBI15 Sequence Listing.ST25

Leu 145	Arg	Ile	Ile	Val	Asp 150	Gly	Val	Met	Ala	His 155	Tyr	Glu	Glu	Leu	Phe 160	
Arg	Ile	Lys	Ser	Asn 165	Ala	Ala	Lys	Asn	Asp 170	Val	Phe	His	Leu	Leu 175	Ser	
Gly	Met	Trp	Lys 180	Thr	Pro	Ala	Glu	Arg 185	Сув	Phe	Leu	Trp	Leu 190	Gly	Gly	
Phe	Arg	Ser 195	Ser	Glu	Leu	Leu	Lys 200	Leu	Leu	Ala	Asn	Gln 205	Leu	Glu	Pro	
Met	Thr 210	Glu	Arg	Gln	Leu	Met 215	Gly	Ile	Asn	Asn	Leu 220	Gln	Gln	Thr	Ser	
Gln 225	Gln	Ala	Glu	Asp	Ala 230	Leu	Ser	Gln	Gly	Met 235	Glu	Ser	Leu	Gln	Gln 240	
Ser	Leu	Ala	Asp	Thr 245	Leu	Ser	Ser	Gly	Thr 250	Leu	Gly	Ser	Ser	Ser 255	Ser	
Gly	Asn	Val	Ala 260	Ser	Tyr	Met	Gly	Gln 265	Met	Ala	Met	Ala	Met 270	Gly	Lys	
Leu	Gly	Thr 275	Leu	Glu	Gly	Phe	Ile 280	Arg	Gln	Ala	Asp	Asn 285	Leu	Arg	Leu	
Gln	Thr 290	Leu	Gln	Gln	Met	Ile 295	Arg	Val	Leu	Thr	Thr 300	Arg	Gln	Ser	Ala	
Arg 305	Ala	Leu	Leu	Ala	Ile 310	His	Asp	Tyr	Phe	Ser 315	Arg	Leu	Arg	Ala	Leu 320	
Ser	Ser	Leu	Trp	Leu 325	Ala	Arg	Pro	Arg	Glu 330							
<210 <211 <212 <213	l> 4 2> I	37 136 DNA Arabi	idops	sis t	hali	iana										
<220 <221 <222 <223	L> (}>	CDS (83). G1396		L3)												
<400 tcga		37 cgt t	tect	ttc	et co	etete	ette	tac	catt	agt:	acgt	tact	gg a	agcto	gatete	60
acgt	atat	tt t	ggat	cgta	aa to	ato Met	gad : Asi	ggc Gly	gaa Glu	gat Asp 5	ttt Phe	gco Ala	gga Gly	a aag / Lys	g gcg s Ala 10	112
gct Ala	gct Ala	gaa Glu	gcc Ala	aag Lys 15	gga Gly	ttg Leu	aac Asn	ccg Pro	gga Gly 20	tta Leu	atc Ile	gtg Val	ctg Leu	ctt Leu 25	gtt Val	160
gtt Val	gga Gly	ggt Gly	ccg Pro 30	ctt Leu	ctt Leu	gtg Val	ttc Phe	cta Leu 35	atc Ile	gcc Ala	aac Asn	tac Tyr	gtg Val 40	ctt Leu	tac Tyr	208

MBI15 Sequence Listing.ST25													
gtt tat gct cag aag aac cta cct cca agg aag aag aag ccc gtt tcc Val Tyr Ala Gln Lys Asn Leu Pro Pro Arg Lys Lys Pro Val Ser 45 50 55	256												
aaa aag aag ctc aag cgg gag aag cta aag caa gga gtc cct gtc cct Lys Lys Lys Leu Lys Arg Glu Lys Leu Lys Gln Gly Val Pro Val Pro 60 65 70	304												
gga gaa taa aagccagctt aagctteett eacttgtgee teetteaaag Gly Glu 75	353												
eggtttttgt teggttacca aattteacce ttgegggttt ttttetteet ttacttetgt	413												
catgaggatt atctttgagg cct	436												
<210> 38 <211> 76 <212> PRT <213> Arabidopsis thaliana													
<400> 38													
Met Asp Gly Glu Asp Phe Ala Gly Lys Ala Ala Ala Glu Ala Lys Gly 1 10 15													
Leu Asn Pro Gly Leu Ile Val Leu Leu Val Val Gly Gly Pro Leu Leu 20 25 30													
Val Phe Leu Ile Ala Asn Tyr Val Leu Tyr Val Tyr Ala Gln Lys Asn 35 40 45													
Leu Pro Pro Arg Lys Lys Pro Val Ser Lys Lys Lys Leu Lys Arg 50 60													
Glu Lys Leu Lys Gln Gly Val Pro Val Pro Gly Glu 65 70 75													
<210> 39 <211> 1470 <212> DNA <213> Arabidopsis thaliana													
<220> <221> CDS <222> (280)(1317) <223> G265													
<400> 39 ctttggtctt ggaagccaaa tcaaaccttt ccttcaatcc tcaaattttc gaaaattttc	60												
tettttgett taegttetet caattettat ttgtaagaaa gtttgtteet ttaatcaate	120												
aaatcaaaga gacttttgaa gattgtttcc caatttgcgt caatcgggat cgagtcaaat	180												
ctgaaatett eteeacteat eatetgaeta taagaettaa teaagggaet ttttgttegg	240												
gtttggtttt aaacgtcttg gattcgaagt ggttaaggt atg gat gaa aat aat Met Asp Glu Asn Asn 1 5	294												
gga ggt tca agc tca ctt cca cct ttc ctt act aaa aca tat gaa atg Gly Gly Ser Ser Leu Pro Pro Phe Leu Thr Lys Thr Tyr Glu Met 10 15 20	342												
gtt gat gat tot tot tot gac tog gto gtt got tgg ago gaa aac aac Val Asp Asp Ser Ser Ser Asp Ser Val Val Ala Trp Ser Glu Asn Asn Page 49	390												

MBI15 Sequence Listing.ST25

			25					30				5	35			
aaa Lys	agc Ser	ttc Phe 40	atc Ile	gtc Val	aag Lys	aat Asn	cca Pro 45	gca Ala	gag Glu	ttt Phe	tca Ser	aga Arg 50	gac Asp	ctt Leu	ctt Leu	438
	aga Arg 55															486
	aca Thr															534
	gat Asp															582
	cga Arg															630
	cct Pro															678
aga Arg	ctg Leu 135	aaa Lys	aat Asn	gag Glu	aaa Lys	gaa Glu 140	ggc Gly	ctt Leu	ctt Leu	gcg Ala	gag Glu 145	tta Leu	cag Gln	aac Asn	caa Gln	726
gag Glu 150	caa Gln	gaa Glu	cgg Arg	aaa Lys	gag Glu 155	ttt Phe	gag Glu	ctg Leu	caa Gln	gta Val 160	acg Thr	aca Thr	ttg Leu	aaa Lys	gat Asp 165	774
cgg Arg	tta Leu	caa Gln	cat His	atg Met 170	gaa Glu	caa Gln	cat His	cag Gln	aaa Lys 175	tca Ser	ata Ile	gtg Val	gca Ala	tat Tyr 180	gtt Val	822
	cag Gln															870
	aga Arg															918
	cac His 215															966
gag Glu 230	aat Asn	ctt Leu	gta Val	tcg Ser	gaa Glu 235	tca Ser	tgc Cys	gag Glu	aag Lys	agc Ser 240	ggt Gly	ttg Leu	cag Gln	tca Ser	tca Ser 245	1014
agc Ser	atg Met	gat Asp	cat His	gat Asp 250	gca Ala	gct Ala	gag Glu	tca Ser	agt Ser 255	cta Leu	agt Ser	att Ile	ggc Gly	gat Asp 260	aca Thr	1062
	ccc Pro															1110
	act Thr															1158
caa Gln	tgt Cys 295	ttg Leu	aca Thr	gag Glu	aac Asn	cct Pro 300	gga Gly	tca Ser	acc Thr	gag Glu	caa Gln 305	caa Gln	gaa Glu	gtt Val	cag Gln	1206
	gag Glu															1254
aat	caa	agg	acg	tat	tgg	tgg	aat	tca		aat age		aat	aac	att	aca	1302

1357

Asn Gln Arg Th	r Tyr Trp Trp 330	MBI15 Sequence Asn Ser Gly As 335	Listing.ST25 n Val Asn Asn Ile Thr 340
gag aaa gct tc Glu Lys Ala Se 34	r	gag gtttttgtaa	aatagttttc ttttggttcc

actgagatta ttgtatgtgt tcattattta ttactctgtt tctgtaaaaa caaatctctc 1417 tattgtttga ggcaggagtg acataaatgc atatgcagaa ttggtttcaa aaa 1470

<210> 40 <211> 345 <212> PRT <213> Arabidopsis thaliana

<400> 40

Met Asp Glu Asn Asn Gly Gly Ser Ser Ser Leu Pro Pro Phe Leu Thr 1 5 15

Lys Thr Tyr Glu Met Val Asp Asp Ser Ser Ser Asp Ser Val Val Ala 20 25 30

Trp Ser Glu Asn Asn Lys Ser Phe Ile Val Lys Asn Pro Ala Glu Phe 35 40 45

Ser Arg Asp Leu Leu Pro Arg Phe Phe Lys His Lys Asn Phe Ser Ser 50 55 60

Phe Ile Arg Gln Leu Asn Thr Tyr Gly Phe Arg Lys Val Asp Pro Glu 65 70 75 80

Lys Trp Glu Phe Leu Asn Asp Asp Phe Val Arg Gly Arg Pro Tyr Leu 85 90 95

Met Lys Asn Ile His Arg Arg Lys Pro Val His Ser His Ser Leu Val

Asn Leu Gln Ala Gln Asn Pro Leu Thr Glu Ser Glu Arg Arg Ser Met 115 120 125

Glu Asp Gln Ile Glu Arg Leu Lys Asn Glu Lys Glu Gly Leu Leu Ala 130 135 140

Glu Leu Gln Asn Gln Glu Gln Glu Arg Lys Glu Phe Glu Leu Gln Val 145 150 155 160

Thr Thr Leu Lys Asp Arg Leu Gln His Met Glu Gln His Gln Lys Ser 165 170 175

Ile Val Ala Tyr Val Ser Gln Val Leu Gly Lys Pro Gly Leu Ser Leu 180 185 190

Asn Leu Glu Asn His Glu Arg Arg Lys Arg Arg Phe Gln Glu Asn Ser 195 200 205

Leu Pro Pro Ser Ser Ser His Ile Glu Gln Val Glu Lys Leu Glu Ser 210 215 220

MBI15 Sequence Listing.ST25 Ser Leu Thr Phe Trp Glu Asn Leu Val Ser Glu Ser Cys Glu Lys Ser 230 Gly Leu Gln Ser Ser Ser Met Asp His Asp Ala Ala Glu Ser Ser Leu Ser Ile Gly Asp Thr Arg Pro Lys Ser Ser Lys Ile Asp Met Asn Ser Glu Pro Pro Val Thr Val Thr Ala Pro Ala Pro Lys Thr Gly Val Asn Asp Asp Phe Trp Glu Gln Cys Leu Thr Glu Asn Pro Gly Ser Thr Glu 290 295 300 Gln Gln Glu Val Gln Ser Glu Arg Arg Asp Val Gly Asn Asp Asn Asn Gly Asn Lys Ile Gly Asn Gln Arg Thr Tyr Trp Trp Asn Ser Gly Asn 325 330 335 Val Asn Asn Ile Thr Glu Lys Ala Ser <210> 41 <211> 913 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (52)..(783) <223> G1006 <400> 41 gataaatcaa tcaacaaaac aaaaaaaact ctatagttag tttctctgaa a atg tac Met Tyr gga cag tgc aat ata gaa tcc gac tac gct ttg ttg gag tcg ata aca Gly Gln Cys Asn Ile Glu Ser Asp Tyr Ala Leu Leu Glu Ser Ile Thr 105 cgt cac ttg cta gga gga gga gag aac gag ctg cga ctc aat gag Arg His Leu Leu Gly Gly Gly Gly Glu Asn Glu Leu Arg Leu Asn Glu 153 tca aca ccg agt tcg tgt ttc aca gag agt tgg gga ggt ttg cca ttg Ser Thr Pro Ser Ser Cys Phe Thr Glu Ser Trp Gly Gly Leu Pro Leu 35 40 50 201 aaa gag aat gat tca gag gac atg ttg gtg tac gga ctc ctc aaa gat Lys Glu Asn Asp Ser Glu Asp Met Leu Val Tyr Gly Leu Leu Lys Asp 249 gcc ttc cat ttt gac acg tca tca tcg gac ttg agc tgt ctt ttt gat Ala Phe His Phe Asp Thr Ser Ser Ser Asp Leu Ser Cys Leu Phe Asp 70 75 80 297 ttt ccg gcg gtt aaa gtc gag cca act gag aac ttt acg gcg atg gag Phe Pro Ala Val Lys Val Glu Pro Thr Glu Asn Phe Thr Ala Met Glu 345 gag aaa cca aag aaa gcg ata ccg gtt acg gag acg gca gtg aag gcg Glu Lys Pro Lys Lys Ala Ile Pro Val Thr Glu Thr Ala Val Lys Ala 393 105

MBI15 Sequence Listing.ST25

							MBT	12 2	eque	nce	PIBC	ing.	5125			
aag Lys 115	cat His	tac Tyr	aga Arg	gga Gly	gtg Val 120	agg Arg	cag Gln	aga Arg	ccg Pro	tgg Trp 125	ggg Gly	aaa Lys	ttc Phe	gcg Ala	gcg Ala 130	441
					gcg Ala											489
ttt Phe	gag Glu	acg Thr	gcg Ala 150	gaa Glu	gat Asp	gcg Ala	gct Ala	tta Leu 155	gct Ala	tac Tyr	gat Asp	ata Ile	gct Ala 160	gct Ala	ttt Phe	537
					cgc Arg											585
					ccg Pro											633
					tcc Ser 200											681
ttg Leu	aaa Lys	cga Arg	agg Arg	aga Arg 215	aaa Lys	gca Ala	gag Glu	aat Asn	ctg Leu 220	acg Thr	tcg Ser	gag Glu	gtg Val	gtg Val 225	cag Gln	729
gtg Val	aag Lys	tgt Cys	gag Glu 230	gtt Val	ggt Gly	gat Asp	gag Glu	aca Thr 235	cgt Arg	gtt Val	gat Asp	gag Glu	tta Leu 240	ttg Leu	gtt Val	777
tca Ser	taa	gtti	tgato	ett 9	gtgt	gttt	tg ta	agtt	gaata	a gtt	ttg	ctat	aaa	tgttq	gag	833
gca	ccaa	gta a	aaagi	tgtto	cc c	gtgat	tgtaa	a att	agtt	act	aaa	cagag	gcc a	atata	atcttc	893
aat	caaa	aaa a	aaaa	aaaa	aa											913
<210> 42 <211> 243 <212> PRT <213> Arabidopsis thaliana																
<40	0 > 4	12														
Met 1	Tyr	Gly	Gln	Сув 5	Asn	Ile	Glu	Ser	Asp 10	Tyr	Ala	Leu	Leu	Glu 15	Ser	
Ile	Thr	Arg	His 20	Leu	Leu	Gly	Gly	Gly 25	Gly	Glu	Asn	Glu	Leu 30	Arg	Leu	
Asn	Glu	Ser 35	Thr	Pro	Ser	Ser	Cys 40	Phe	Thr	Glu	Ser	Trp 45	Gly	Gly	Leu	
Pro	Leu 50	Lys	Glu	Asn	Asp	Ser 55	Glu	Asp	Met	Leu	Val 60	Tyr	Gly	Leu	Leu	
Lys 65	Asp	Ala	Phe	His	Phe 70	Asp	Thr	Ser	Ser	Ser 75	Asp	Leu	Ser	Cys	Leu 80	
Phe	Asp	Phe	Pro		Val	Lys	Val	Glu	Pro 90	Thr	Glu	Asn	Phe	Thr 95	Ala	
				85					30							

MBI15 Sequence Listing.ST25

Lys Ala Lys His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly Lys Phe Ala Ala Glu Ile Arg Asp Pro Ala Lys Asn Gly Ala Arg Val Trp Leu Gly Thr Phe Glu Thr Ala Glu Asp Ala Ala Leu Ala Tyr Asp Ile Ala Ala Phe Arg Met Arg Gly Ser Arg Ala Leu Leu Asn Phe Pro Leu Arg Val Asn Ser Gly Glu Pro Asp Pro Val Arg Ile Thr Ser Lys Arg Ser Thr Ser Ser Ser Glu Asn Gly Lys Leu Lys Arg Arg Lys Ala Glu Asn Leu Thr Ser Glu Val 210 215 220 Val Gln Val Lys Cys Glu Val Gly Asp Glu Thr Arg Val Asp Glu Leu Leu Val Ser <210> 43 <211> 912 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (53)..(859) G1309 <400> 43 cgtcgacctc ttaattaaga cgacttgaga gagaaagaaa gatacgtgga ag atg acc 58 Met Thr 106 aaa tot gga gag aga oca aaa cag aga cag agg aaa ggg tta tgg tca Lys Ser Gly Glu Arg Pro Lys Gln Arg Gln Arg Lys Gly Leu Trp Ser cct gaa gaa gac cag aag ctc aag agt ttc atc ctc tct cgt ggc cat 154 Pro Glu Glu Asp Gln Lys Leu Lys Ser Phe Ile Leu Ser Arg Gly His gct tgc tgg acc act gtt ccc atc cta gct gga ttg caa agg aat ggg Ala Cys Trp Thr Thr Val Pro Ile Leu Ala Gly Leu Gln Arg Asn Gly 202 aaa agc tgc aga tta agg tgg att aat tac cta aga cca gga cta aag Lys Ser Cys Arg Leu Arg Trp Ile Asn Tyr Leu Arg Pro Gly Leu Lys 250 agg ggg tcg ttt agt gaa gaa gaa gaa gag acc atc ttg act tta cat Arg Gly Ser Phe Ser Glu Glu Glu Glu Glu Thr Ile Leu Thr Leu His 70 75 80298 346 tet tee ttg ggt aac aag tgg tet egg att gea aaa tat tta eeg gga

MBI15 Sequence Listing.ST25 Ser Ser Leu Gly Asn Lys Trp Ser Arg Ile Ala Lys Tyr Leu Pro Gly 85 90 95	
aga aca gac aac gag att aag aac tat tgg cat tcc tat ctg aag aag Arg Thr Asp Asn Glu Ile Lys Asn Tyr Trp His Ser Tyr Leu Lys Lys	394
aga tgg ctc aaa tct caa cca caa ctc aaa agc caa ata tca gac ctc	442
Arg Trp Leu Lys Ser Gln Pro Gln Leu Lys Ser Gln Ile Ser Asp Leu 115 120 125 130	
aca gaa tot cot tot toa ota ott tot tgo ggg aaa aga aat otg gaa Thr Glu Ser Pro Ser Ser Leu Leu Ser Cys Gly Lys Arg Asn Leu Glu 135 140 145	490
acc gaa acc cta gat cac gtg atc tcc ttc cag aaa ttt tca gag aat Thr Glu Thr Leu Asp His Val Ile Ser Phe Gln Lys Phe Ser Glu Asn 150 155 160	538
CCa act tca tca cca tcc aaa gaa agc aac aac aac atg atc atg aac Pro Thr Ser Ser Pro Ser Lys Glu Ser Asn Asn Asn Met Ile Met Asn 165 170 175	586
aac agt aat aac ttg cct aaa ctg ttc ttc tct gag tgg atc agt tct Asn Ser Asn Asn Leu Pro Lys Leu Phe Phe Ser Glu Trp Ile Ser Ser 180 185 190	634
tca aat cca cac atc gat tac tcc tct gct ttt aca gat tcc aag cac Ser Asn Pro His Ile Asp Tyr Ser Ser Ala Phe Thr Asp Ser Lys His 195 200 205 210	682
att aat gaa act caa gat caa atc aat gaa gag gaa gtg atg atc Ile Asn Glu Thr Gln Asp Gln Ile Asn Glu Glu Glu Val Met Met Ile 215 220 225	730
aat aac aac aac tac tct tca ctt gag gat gtc atg ctc cgt aca gat Asn Asn Asn Tyr Ser Ser Leu Glu Asp Val Met Leu Arg Thr Asp 230 235 240	778
ttt ttg cag cct gat cat gaa tat gca aat tat tat tct tct gga gat Phe Leu Gln Pro Asp His Glu Tyr Ala Asn Tyr Tyr Ser Ser Gly Asp 245 250 255	826
ttc ttc atc aac agt gac caa aat tat gtc taa gaagagtgaa tatgatcgta Phe Phe Ile Asn Ser Asp Gln Asn Tyr Val 260 265	879
agaggaacat aagctagtta cttgtgttac agc	912
<210> 44 <211> 268 <212> PRT <213> Arabidopsis thaliana	
<400> 44	
Met Thr Lys Ser Gly Glu Arg Pro Lys Gln Arg Gln Arg Lys Gly Leu 1 5 10 15	
Trp Ser Pro Glu Glu Asp Gln Lys Leu Lys Ser Phe Ile Leu Ser Arg 20 25 30	
Gly His Ala Cys Trp Thr Thr Val Pro Ile Leu Ala Gly Leu Gln Arg 35 40 45	
Asn Gly Lys Ser Cys Arg Leu Arg Trp Ile Asn Tyr Leu Arg Pro Gly 50 55 60	
Leu Lys Arg Gly Ser Phe Ser Glu Glu Glu Glu Glu Thr Ile Leu Thr 65 70 75 80	

MBI15 Sequence Listing.ST25

Leu His Ser Ser Leu Gly Asn Lys Trp Ser Arg Ile Ala Lys Tyr Leu Pro Gly Arg Thr Asp Asn Glu Ile Lys Asn Tyr Trp His Ser Tyr Leu Lys Lys Arg Trp Leu Lys Ser Gln Pro Gln Leu Lys Ser Gln Ile Ser Asp Leu Thr Glu Ser Pro Ser Ser Leu Leu Ser Cys Gly Lys Arg Asn Leu Glu Thr Glu Thr Leu Asp His Val Ile Ser Phe Gln Lys Phe Ser Glu Asn Pro Thr Ser Ser Pro Ser Lys Glu Ser Asn Asn Asn Met Ile Met Asn Asn Ser Asn Asn Leu Pro Lys Leu Phe Phe Ser Glu Trp Ile 180 185 190 Ser Ser Ser Asn Pro His Ile Asp Tyr Ser Ser Ala Phe Thr Asp Ser Lys His Ile Asn Glu Thr Gln Asp Gln Ile Asn Glu Glu Glu Val Met Met Ile Asn Asn Asn Asn Tyr Ser Ser Leu Glu Asp Val Met Leu Arg Thr Asp Phe Leu Gln Pro Asp His Glu Tyr Ala Asn Tyr Tyr Ser Ser Gly Asp Phe Phe Ile Asn Ser Asp Gln Asn Tyr Val <210> 45 <211> 1575 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (1)..(1575) <223> G2550 atg gct gtg tat tac cct aat agt gtc ggc atg caa tct ctt tac caa Met Ala Val Tyr Tyr Pro Asn Ser Val Gly Met Gln Ser Leu Tyr Gln 1 5 10 15 48 96 tct tcc tct gct gca tct ttc tcc gag att gtt tcc ggt gat gtt cga Ser Ser Ser Ala Ala Ser Phe Ser Glu Ile Val Ser Gly Asp Val Arg 144 aac aac gag atg gta ttt atc cca cca aca agc gac gta gcc gtc aac 192 Page 56

Asn	Asn 50	Glu	Met	Val	Phe	Ile 55	MBI Pro	15 S Pro	eque: Thr	nce : Ser	List: Asp 60	ing.: Val	ST25 Ala	Val	Asn	
gga Gly 65	aac Asn	gta Val	acg Thr	gtg Val	tca Ser 70	agt Ser	aac Asn	gat Asp	cta Leu	agc Ser 75	ttt Phe	cac His	ggt Gly	gga Gly	gga Gly 80	240
ctt Leu	tct Ser	tta Leu	agt Ser	ctt Leu 85	ggt Gly	aat Asn	cag Gln	atc Ile	cag Gln 90	tca Ser	gct Ala	gtc Val	tct Ser	gtt Val 95	tct Ser	288
ccg Pro	ttt Phe	cag Gln	tat Tyr 100	cat His	tac Tyr	cag Gln	aat Asn	ctt Leu 105	tcg Ser	aac Asn	caa Gln	ttg Leu	agt Ser 110	tac Tyr	aat Asn	336
aat Asn	ctt Leu	aat Asn 115	cct Pro	tct Ser	act Thr	atg Met	tct Ser 120	gat Asp	gag Glu	aat Asn	Gly ggg	aag Lys 125	agc Ser	ttg Leu	agt Ser	384
gtt Val	cat His 130	cag Gln	cat His	cac His	tct Ser	gat Asp 135	caa Gln	att Ile	tta Leu	cct Pro	tcc Ser 140	tct Ser	gtt Val	tac Tyr	aac Asn	432
aac Asn 145	aat Asn	ggt Gly	aat Asn	aat Asn	ggt Gly 150	gtt Val	gga Gly	ttc Phe	tac Tyr	aac Asn 155	aat Asn	tac Tyr	cgt Arg	tac Tyr	gag Glu 160	480
aca Thr	tca Ser	Gly ggg	ttt Phe	gtg Val 165	agt Ser	agt Ser	gta Val	ctg Leu	aga Arg 170	tct Ser	cgt Arg	tac Tyr	ctt Leu	aaa Lys 175	cca Pro	528
aca Thr	caa Gln	caa Gln	ttg Leu 180	ctt Leu	gat Asp	gaa Glu	gtt Val	gtt Val 185	agt Ser	gta Val	agg Arg	aaa Lys	gat Asp 190	ttg Leu	aaa Lys	576
ttg Leu	ggg Gly	aat Asn 195	aag Lys	aag Lys	atg Met	aag Lys	aat Asn 200	gat Asp	aaa Lys	ggt Gly	caa Gln	gac Asp 205	ttt Phe	cac His	aat Asn	624
999 Gly	tct Ser 210	agt Ser	gat Asp	aac Asn	att Ile	aca Thr 215	gaa Glu	gat Asp	gat Asp	aaa Lys	tct Ser 220	caa Gln	tcg Ser	cag Gln	gag Glu	. 672
Leu 225	Ser	Pro	Ser	Glu	Arg 230	Gln	Glu	cta Leu	Gln	Ser 235	ГÀв	Lys	Ser	Lys	240	720
Leu	Thr	Met	Val	Asp 245	Glu	Val	Asp	aaa Lys	Arg 250	Tyr	Asn	GIn	Tyr	255	HIS	768
caa Gln	atg Met	gaa Glu	gct Ala 260	tta Leu	gca Ala	tcg Ser	tct Ser	ttc Phe 265	gag Glu	atg Met	gta Val	aca Thr	ggt Gly 270	ctt Leu	gga Gly	816
gca Ala	gct Ala	aag Lys 275	cct Pro	tac Tyr	aca Thr	tcc Ser	gta Val 280	gct Ala	ctg Leu	aat Asn	aga Arg	atc Ile 285	tct Ser	cgc Arg	cat His	864
ttc Phe	cgc Arg 290	tgt Cys	tta Leu	cgc Arg	gac Asp	gcg Ala 295	ata Ile	aaa Lys	gaa Glu	cag Gln	att Ile 300	cag Gln	gtg Val	atc Ile	aga Arg	912
999 Gly 305	aag Lys	ctt Leu	Gly Ggg	gag Glu	aga Arg 310	gag Glu	act Thr	tct Ser	gat Asp	gaa Glu 315	caa Gln	gga Gly	gag Glu	agg Arg	ata Ile 320	960
ccg Pro	cgt Arg	ctt Leu	agg Arg	tac Tyr 325	tta Leu	gat Asp	caa Gln	cgg Arg	ttg Leu 330	aga Arg	caa Gln	cag Gln	aga Arg	gct Ala 335	ttg Leu	1008
cat His	caa Gln	caa Gln	ctt Leu 340	gga Gly	atg Met	gtt Val	aga Arg	cca Pro 345	gct Ala	tgg Trp	aga Arg	cca Pro	caa Gln 350	aga Arg	ggc Gly	1056

							ata		cga	gct	tgg	ctc	ttt	gag Glu		11	104
ttc Phe	ctt Leu 370	cat His	cca Pro	tat Tyr	cct Pro	aaa Lys 375	gaa Glu	tca Ser	gag Glu	aaa Lys	atc Ile 380	atg Met	ctt Leu	tca Ser	aag Lys	11	152
														aac Asn		12	200
														gaa Glu 415		12	248
														acc Thr		12	296
														tct Ser		13	344
														aca Thr		13	392
														cgt Arg		14	140
act Thr	act Thr	gga Gly	gat Asp	tac Tyr 485	gac Asp	agc Ser	ttg Leu	atg Met	aac Asn 490	tat Tyr	cat His	Gly ggg	ttt Phe	ggt Gly 495	att Ile	14	188
														ggc Gly		15	536
		aat Asn 515										taa				15	575
<210 <211 <212 <213	L> 5 2> I	6 524 PRT Arabi	ldops	sis t	:hali	.ana											
<400)> 4	6															
Met 1	Ala	Val	Tyr	Tyr 5	Pro	Asn	Ser	Val	Gly 10	Met	Gln	Ser	Leu	Tyr 15	Gln		
Glu	Ser	Ile	Tyr 20	Leu	Asn	Glu	Gln	Gln 25	Gln	Gln	Gln	Gln	Gln 30	Ala	Ser		
Ser	Ser	Ser 35	Ala	Ala	Ser	Phe	Ser 40	Glu	Ile	Val	Ser	Gly 45	Asp	Val	Arg		
Asn	Asn 50	Glu	Met	Val	Phe	Ile 55	Pro	Pro	Thr	Ser	Asp 60	Val	Ala	Val	Asn		
Gly 65	Asn	Val	Thr	Val	Ser 70	Ser	Asn	Asp	Leu	Ser 75	Phe	His	Gly	Gly	Gly 80		
Leu	Ser	Leu	Ser	Leu	Gly	Asn	Gln	Ile	Gln	Ser	Ala	Val	Ser	Val	Ser		

MBI15 Sequence Listing.ST25

Pro Phe Gln Tyr His Tyr Gln Asn Leu Ser Asn Gln Leu Ser Tyr Asn 100 105 110

Asn Leu Asn Pro Ser Thr Met Ser Asp Glu Asn Gly Lys Ser Leu Ser 115 120 125

Val His Gln His His Ser Asp Gln Ile Leu Pro Ser Ser Val Tyr Asn 130 135 140

Asn Asn Gly Asn Asn Gly Val Gly Phe Tyr Asn Asn Tyr Arg Tyr Glu 145 150 155 160

Thr Ser Gly Phe Val Ser Ser Val Leu Arg Ser Arg Tyr Leu Lys Pro

Thr Gln Gln Leu Leu Asp Glu Val Val Ser Val Arg Lys Asp Leu Lys 180 185 190

Leu Gly Asn Lys Lys Met Lys Asn Asp Lys Gly Gln Asp Phe His Asn 195 200 205

Gly Ser Ser Asp Asn Ile Thr Glu Asp Asp Lys Ser Gln Ser Gln Glu 210 215 220

Leu Ser Pro Ser Glu Arg Gln Glu Leu Gln Ser Lys Lys Ser Lys Leu 225 230 235 240

Leu Thr Met Val Asp Glu Val Asp Lys Arg Tyr Asn Gln Tyr His His 245 250 255

Gln Met Glu Ala Leu Ala Ser Ser Phe Glu Met Val Thr Gly Leu Gly 260 265 270

Phe Arg Cys Leu Arg Asp Ala Ile Lys Glu Gln Ile Gln Val Ile Arg 290 295 300

Gly Lys Leu Gly Glu Arg Glu Thr Ser Asp Glu Gln Gly Glu Arg Ile 305 310 315 320

Pro Arg Leu Arg Tyr Leu Asp Gln Arg Leu Arg Gln Gln Arg Ala Leu 325 330 335

His Gln Gln Leu Gly Met Val Arg Pro Ala Trp Arg Pro Gln Arg Gly 340 345 350

Phe Leu His Pro Tyr Pro Lys Glu Ser Glu Lys Ile Met Leu Ser Lys 370 375 380

Gln Thr Gly Leu Ser Lys Asn Gln Val Ala Asn Trp Phe Ile Asn Ala 385 390 395 400

PCT/US00/31418 WO 01/35726

MBI15 Sequence Listing.ST25

Arg Val Arg Leu Trp Lys Pro Met Ile Glu Glu Met Tyr Lys Glu Glu Phe Gly Glu Ser Ala Glu Leu Leu Ser Asn Ser Asn Gln Asp Thr Lys Lys Met Gln Glu Thr Ser Gln Leu Lys His Glu Asp Ser Ser Ser Gln Gln Gln Asn Gln Gly Asn Asn Asn Asn Ile Pro Tyr Thr Ser 455 Asp Ala Glu Gln Asn Leu Val Phe Ala Asp Pro Lys Pro Asp Arg Ala Thr Thr Gly Asp Tyr Asp Ser Leu Met Asn Tyr His Gly Phe Gly Ile Asp Asp Tyr Asn Arg Tyr Val Gly Leu Gly Asn Gln Gln Asp Gly Arg 505 Tyr Ser Asn Pro His Gln Leu His Asp Phe Val Val 520 <210> 47 <211> 1983 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (73)..(1956) <223> G965 <400> 47 60 gattctctgt gtatgtctga atccttacag gatccaagag ctttggaaaa aagatataat gaataacaag at atg ggt tta gct act aca act tct tct atg tca caa gat Met Gly Leu Ala Thr Thr Thr Ser Ser Met Ser Gln Asp 111 tat cat cat cac caa gga atc ttt tcc ttc tct aat gga ttc cac cga
Tyr His His His Gln Gly Ile Phe Ser Phe Ser Asn Gly Phe His Arg
15 20 25 159 207 tca tca tca acc act cat cag gag gaa gta gat gaa tcc gcc gtc gtc Ser Ser Ser Thr Thr His Gln Glu Glu Val Asp Glu Ser Ala Val Val tcc ggt gct caa att ccg gtt tat gaa acc gcc gga atg ttg tct gaa Ser Gly Ala Gln Ile Pro Val Tyr Glu Thr Ala Gly Met Leu Ser Glu 255 atg ttt gct tac cct ggc gga ggt ggc ggc ggt tcc ggt gga gag att Met Phe Ala Tyr Pro Gly Gly Gly Gly Gly Gly Ser Gly Glu Ile 65 70 75303 ctt gat cag tct act aaa cag ttg cta gag caa caa aac cgt cac aac Leu Asp Gln Ser Thr Lys Gln Leu Leu Glu Gln Gln Asn Arg His Asn 351

Page 60

399

aac aac aat aac tca act ctt cat atg tta tta cca aat cat cat caa Asn Asn Asn Asn Ser Thr Leu His Met Leu Leu Pro Asn His His Gln

100

MBI15 Sequence Listing.ST25

									•			_				
ggt Gly 110	ttt Phe	gct Ala	ttc Phe	acc Thr	gac Asp 115	gaa Glu	aac Asn	act Thr	atg Met	cag Gln 120	ccg Pro	cag Gln	caa Gln	caa Gln	caa Gln 125	447
					tct Ser											495
atg Met	atc Ile	gga Gly	acc Thr 145	gtc Val	cac His	gtg Val	gaa Glu	gga Gly 150	gga Gly	aag Lys	ggt Gly	ttg Leu	tct Ser 155	tta Leu	tct Ser	543
ctc Leu	tca Ser	tct Ser 160	tca Ser	tta Leu	gcc Ala	gca Ala	gct Ala 165	aaa Lys	gcc Ala	gag Glu	gaa Glu	tat Tyr 170	aga Arg	agc Ser	att Ile	591
tat Tyr	tgt Cys 175	gca Ala	gcc Ala	gtt Val	gat Asp	gga Gly 180	act Thr	tct Ser	tct Ser	tct Ser	tct Ser 185	aac Asn	gca Ala	tcc Ser	gct Ala	639
					aat Asn 195											687
tct Ser	tct Ser	caa Gln	cat His	cat His 210	cac His	cat His	caa Gln	gtt Val	gtt Val 215	gga Gly	cat His	ttt Phe	ggt Gly	tca Ser 220	tca Ser	735
tca Ser	tca Ser	tct Ser	ccc Pro 225	atg Met	gcg Ala	gct Ala	tct Ser	tca Ser 230	tcc Ser	att Ile	gga Gly	999 Gly	atc Ile 235	tac Tyr	acg Thr	783
ttg Leu	agg Arg	aat Asn 240	tcg Ser	aaa Lys	tat Tyr	acg Thr	aaa Lys 245	ccg Pro	gct Ala	caa Gln	gag Glu	ttg Leu 250	ttg Leu	gaa Glu	gag Glu	831
ttt Phe	tgt Cys 255	agt Ser	gtt Val	gga Gly	aga Arg	gga Gly 260	cat His	ttc Phe	aag Lys	aag Lys	aac Asn 265	aaa Lys	ctt Leu	agt Ser	agg Arg	879
aac Asn 270	aac Asn	tca Ser	aac Asn	cct Pro	aat Asn 275	act Thr	acc Thr	ggt Gly	gga Gly	gga Gly 280	gga Gly	ggc Gly	gga Gly	999 Gly	tcc Ser 285	927
tcg Ser	tca Ser	tcg Ser	gcc Ala	gga Gly 290	aca Thr	gct Ala	aat Asn	gat Asp	agt Ser 295	cct Pro	cct Pro	ttg Leu	tct Ser	ccg Pro 300	gct Ala	975
gat Asp	cgg Arg	att Ile	gaa Glu 305	cat His	caa Gln	aga Arg	aga Arg	aaa Lys 310	gtc Val	aag Lys	cta Leu	cta Leu	tct Ser 315	atg Met	ctt Leu	1023
gaa Glu	g <u>ag</u> Glu	gtg Val 320	gac Asp	cga Arg	cgg Arg	tac Tyr	aac Asn 325	cac His	tac Tyr	tgc Cys	gaa Glu	caa Gln 330	atg Met	caa Gln	atg Met	1071
gta Val	gtg Val 335	aac Asn	tca Ser	ttc Phe	gac Asp	caa Gln 340	gta Val	atg Met	ggt Gly	tac Tyr	ggc Gly 345	gcg Ala	gcg Ala	gtt Val	ccg Pro	1119
tac Tyr 350	acg Thr	aca Thr	tta Leu	gct Ala	caa Gln 355	aag Lys	gca Ala	atg Met	tct Ser	agg Arg 360	cat His	ttc Phe	cgg Arg	tgt Cys	ttg Leu 365	1167
aaa Lys	gac Asp	gcg Ala	gta Val	gcg Ala 370	gtt Val	cag Gln	ctt Leu	aaa Lys	cgc Arg 375	agc Ser	tgt Cys	gag Glu	ctt Leu	cta Leu 380	999 Gly	1215
gat Asp	aaa Lys	gag Glu	gcg Ala 385	gca Ala	GJA GGG	gct Ala	gca Ala	tcc Ser 390	tcg Ser	ggg ggg	tta Leu	acc Thr	aaa Lys 395	ggg Gly	gaa Glu	1263
acg Thr	ccg Pro	cga Arg	ttg Leu	cgt Arg	ttg Leu	cta Leu	gag Glu	cag Gln	Ser	ttg Leu	Arg	cag Gln	caa Gln	cga Arg	gcg Ala	1311

MBI15	Sequence	Listing.ST25
405	-	410

		400					405	15 S	eque	nce	List	ing. 410	ST25			
ttt Phe	cat His 415	cat His	atg Met	ggt Gly	atg Met	atg Met 420	gag Glu	caa Gln	gag Glu	gca Ala	tgg Trp 425	aga Arg	ccg Pro	caa Gln	cgt Arg	1359
		cct Pro														1407
cat His	ttt Phe	ctt Leu	aat Asn	ccg Pro 450	tac Tyr	cca Pro	agc Ser	gat Asp	gct Ala 455	Asp	aag Lys	cac His	ctc Leu	tta Leu 460	gca Ala	1455
cga Arg	cag Gln	act Thr	ggt Gly 465	tta Leu	tcc Ser	aga Arg	aat Asn	cag Gln 470	gtg Val	tca Ser	aat Asn	tgg Trp	ttc Phe 475	ata Ile	aat Asn	1503
		gtt Val 480														1551
gaa Glu	gca Ala 495	aaa Lys	gaa Glu	aga Arg	gaa Glu	gaa Glu 500	gca Ala	gaa Glu	gaa Glu	gaa Glu	aat Asn 505	gaa Glu	aat Asn	caa Gln	caa Gln	1599
		aga Arg														1647
		gaa Glu														1695
atg Met	aca Thr	tcg Ser	aca Thr 545	cat His	cac His	gaa Glu	aac Asn	gac Asp 550	tct Ser	tca Ser	ttc Phe	ctc Leu	tct Ser 555	tcc Ser	gtc Val	1743
		gct Ala 560														1791
cag Gln	caa Gln 575	gac Asp	gtc Val	agt Ser	gac Asp	ttc Phe 580	cac His	gtc Val	gac Asp	gga Gly	gat Asp 585	ggt Gly	gtg Val	aac Asn	gtc Val	1839
ata Ile 590	aga Arg	ttc Phe	ggg ggg	acc Thr	aaa Lys 595	cag Gln	act Thr	ggt Gly	gac Asp	gtg Val 600	tct Ser	ctt Leu	acg Thr	ctt Leu	ggt Gly 605	1887
		cac His														1935
		ttt Phe				tag	tctt	cttt	gt t	tctc	aatt	t at	tcat	:c		1983
<210		8									•					
<211 <212		27 PRT														
<213	> A	rabi	.dops	is t	hali	ana							•			

<213> Arabidopsis thaliana

<400> 48

His Gln Gly Ile Phe Ser Phe Ser Asn Gly Phe His Arg Ser Ser Ser 20 25

Thr Thr His Gln Glu Glu Val Asp Glu Ser Ala Val Val Ser Gly Ala Page 62

MBI15 Sequence Listing.ST25

35

Gln Ile Pro Val Tyr Glu Thr Ala Gly Met Leu Ser Glu Met Phe Ala 50 60

Tyr Pro Gly Gly Gly Gly Gly Ser Gly Gly Glu Ile Leu Asp Gln 65 70 75 80

Ser Thr Lys Gln Leu Leu Glu Gln Gln Asn Arg His Asn Asn Asn Asn Asn 90 95

Asn Ser Thr Leu His Met Leu Leu Pro Asn His His Gln Gly Phe Ala 100 105 110

Phe Thr Asp Glu Asn Thr Met Gln Pro Gln Gln Gln His Phe Thr 115 120 125

Trp Pro Ser Ser Ser Ser Asp His His Gln Asn Arg Asp Met Ile Gly 130 135 140

Thr Val His Val Glu Gly Gly Lys Gly Leu Ser Leu Ser Leu Ser Ser 145 150 150 160

Ser Leu Ala Ala Ala Lys Ala Glu Glu Tyr Arg Ser Ile Tyr Cys Ala 165 170 175

Ala Val Asp Gly Thr Ser Ser Ser Ser Asn Ala Ser Ala His His His 180 185 190

Gln Phe Asn Gln Phe Lys Asn Leu Leu Leu Glu Asn Ser Ser Gln 195 200 205

His His His Gln Val Val Gly His Phe Gly Ser Ser Ser Ser Ser 210 215 220

Pro Met Ala Ala Ser Ser Ser Ile Gly Gly Ile Tyr Thr Leu Arg Asn 225 230 240

Ser Lys Tyr Thr Lys Pro Ala Gln Glu Leu Leu Glu Glu Phe Cys Ser 245 250 255

Val Gly Arg Gly His Phe Lys Lys Asn Lys Leu Ser Arg Asn Asn Ser 260 265 270

Asn Pro Asn Thr Thr Gly Gly Gly Gly Gly Gly Gly Ser Ser Ser Ser 275 280 285

Ala Gly Thr Ala Asn Asp Ser Pro Pro Leu Ser Pro Ala Asp Arg Ile 290 295 300

Glu His Gln Arg Arg Lys Val Lys Leu Leu Ser Met Leu Glu Glu Val 305 310 315 320

Asp Arg Arg Tyr Asn His Tyr Cys Glu Gln Met Gln Met Val Val Asn 325 330 335

MBI15 Sequence Listing.ST25
Ser Phe Asp Gln Val Met Gly Tyr Gly Ala Ala Val Pro Tyr Thr Thr
340 345 350

Leu Ala Gln Lys Ala Met Ser Arg His Phe Arg Cys Leu Lys Asp Ala 355 360 365

Val Ala Val Gln Leu Lys Arg Ser Cys Glu Leu Leu Gly Asp Lys Glu 370 375 380

Ala Ala Gly Ala Ala Ser Ser Gly Leu Thr Lys Gly Glu Thr Pro Arg 385 390 395 400

Leu Arg Leu Leu Glu Gln Ser Leu Arg Gln Gln Arg Ala Phe His His
405 410 415

Met Gly Met Met Glu Gln Glu Ala Trp Arg Pro Gln Arg Gly Leu Pro 420 430

Glu Arg Ser Val Asn Ile Leu Arg Ala Trp Leu Phe Glu His Phe Leu 435 440 445

Asn Pro Tyr Pro Ser Asp Ala Asp Lys His Leu Leu Ala Arg Gln Thr 450 455 460

Gly Leu Ser Arg Asn Gln Val Ser Asn Trp Phe Ile Asn Ala Arg Val 465 470 480

Arg Leu Trp Lys Pro Met Val Glu Glu Met Tyr Gln Gln Glu Ala Lys
485 490 495

Glu Arg Glu Glu Ala Glu Glu Glu Asn Glu Asn Gln Gln Gln Arg 500 505 510

Arg Gln Gln Gln Thr Asn Asn Asn Asp Thr Lys Pro Asn Asn Asn Glu 515 520 525

Asn Asn Phe Thr Val Ile Thr Ala Gln Thr Pro Thr Thr Met Thr Ser 530 540

Thr His His Glu Asn Asp Ser Ser Phe Leu Ser Ser Val Ala Ala Ala 545 550 560

Ser His Gly Gly Ser Asp Ala Phe Thr Val Ala Thr Cys Gln Gln Asp 565 570 575

Val Ser Asp Phe His Val Asp Gly Asp Gly Val Asn Val Ile Arg Phe 580 590

Gly Thr Lys Gln Thr Gly Asp Val Ser Leu Thr Leu Gly Leu Arg His

Ser Gly Asn Ile Pro Asp Lys Asn Thr Ser Phe Ser Val Arg Asp Phe 610 620

Gly Asp Phe 625

MBI15 Sequence Listing.ST25

								- 4			5				
<210><211><212><213>	49 1420 DNA Arab:	idops	sis (hali	iana										
<220> <221> <222> <223>	CDS (138) G793) (:	1046))					ř						
<400>	49 etca o	cctaa	acaa	et co	cacac	caaga	a tto	ette	gcat	ggaa	agti	tgt t	ctto	egactt	60
ctcttct															120
ctttact	cgt 1	teel	ı	atg d Met <i>I</i> L	get a Ala <i>H</i>	aat a Asn i	aac a Asn A	aac a Asn <i>l</i>	aac a Asn 1	atc d Ile I	cca (Pro 1	lis A	gat a Asp S	igc Ser	170
atc tcc Ile Se	gat Asp	cca Pro 15	tct Ser	cct Pro	acc Thr	gac Asp	gat Asp 20	ttc Phe	ttc Phe	gag Glu	cag Gln	atc Ile 25	ctc Leu	999 Gly	218
ctt tcc Leu Sei	aac Asn 30	ttc Phe	tcc Ser	ggt Gly	tct Ser	tca Ser 35	ggt Gly	tct Ser	ggt Gly	ctc Leu	tct Ser 40	gga Gly	atc Ile	ggc Gly	266
ggc gtg Gly Va: 45	g ggt L Gly	cca Pro	cct Pro	ccg Pro	atg Met 50	atg Met	ctt Leu	cag Gln	ctt Leu	ggt Gly 55	tca Ser	ggc Gly	aac Asn	gaa Glu	314
ggg aat Gly Ast 60	cat His	aat Asn	cat His	atg Met 65	ggt Gly	gcc Ala	att Ile	gga Gly	gga Gly 70	ggt Gly	gga Gly	cct Pro	gta Val	999 Gly 75	362
ttt cat Phe His	aat 3 Asn	cag Gln	atg Met 80	ttt Phe	ccg Pro	ttg Leu	gga Gly	tta Leu 85	agt Ser	ctc Leu	gat Asp	caa Gln	999 Gly 90	aaa Lys	410
gga cat Gly His	ggc Gly	ttt Phe 95	ctt Leu	aaa Lys	cct Pro	gat Asp	gaa Glu 100	act Thr	ggt Gly	aaa Lys	cgt Arg	ttc Phe 105	caa Gln	gac Asp	458
gat gti Asp Va	ctt Leu 110	gat Asp	aat Asn	cga Arg	tgt Cys	tcc Ser 115	tct Ser	atg Met	aaa Lys	cct Pro	att Ile 120	ttc Phe	cat His	ggg Gly	506
cag cca Gln Pro) Met	tca Ser	cag Gln	cca Pro	gct Ala 130	cca Pro	cca Pro	atg Met	ccg Pro	cat His 135	caa Gln	cag Gln	tct Ser	act Thr	554
att cgg Ile Arg 140	g cct g Pro	aga Arg	gtt Val	agg Arg 145	gct Ala	agg Arg	cga Arg	ggt Gly	caa Gln 150	gct Ala	acc Thr	gat Asp	cca Pro	cat His 155	602
agc ato Ser Ile	gct Ala	gag Glu	agg Arg 160	ctc Leu	cga Arg	agg Arg	gaa Glu	aga Arg 165	ata Ile	gca Ala	gaa Glu	cgg Arg	atc Ile 170	agg Arg	650
tcg ttg Ser Le	g cag 1 Gln	gaa Glu 175	ctt Leu	gta Val	cct Pro	acc Thr	gtt Val 180	aac Asn	aag Lys	aca Thr	gat Asp	agg Arg 185	gct Ala	gct Ala	698
atg ato Met Ilo	gac Asp 190	gag Glu	att Ile	gtc Val	gat Asp	tat Tyr 195	gta Val	aag Lys	ttt Phe	ctc Leu	agg Arg 200	ctc Leu	caa Gln	gtt Val	746
aag gte Lys Val 20	l Leu	agc Ser	atg Met	agc Ser	cgt Arg 210	ctt Leu	ggt Gly	gga Gly	gcc Ala	ggt Gly 215	gct Ala	gtc Val	gca Ala	cca Pro	794
cta gto Leu Va	c act l Thr	gaa Glu	atg Met	cca Pro	tta Leu	tct Ser	tca Ser	Ser	gtt Val age	Glu	gat Asp	gag Glu	acg Thr	cag Gln	842

MBI15 Sequence Listing.ST25
220 225 230 235
gcc gtg tgg gag aaa tgg tca aac gat ggg aca gag agg caa gtg gct Ala Val Trp Glu Lys Trp Ser Asn Asp Gly Thr Glu Arg Gln Val Ala 240 245 250
aag ctg atg gaa gaa aac gtt gga gca gcg atg caa ctt ttg caa tca Lys Leu Met Glu Glu Asn Val Gly Ala Ala Met Gln Leu Leu Gln Ser 255 260 265
aag gct ctt tgc ata atg ccg atc tca ttg gca atg gcg att tac cat Lys Ala Leu Cys Ile Met Pro Ile Ser Leu Ala Met Ala Ile Tyr His 270 275 280
tct cag cca cca gac aca tct tct tca atc gtc aaa cca gag atg aat Ser Gln Pro Pro Asp Thr Ser Ser Ser Ile Val Lys Pro Glu Met Asn 285 290 295
cct cca ccg tag atttttgttc atccaacggt ccccagctga tgattgacat Pro Pro Pro 300
tttgctctgt ttcccactac tagacttttg tgactcatga aaggtaagta aaaaggcat
ggagatggaa tctaagtagg atttgtgcag taaagaagta aaacgggatc tgtcaaaag
aggaaaaagc tctcgcttgc ttggctagta tttatcattt tgatgaaagt aactctttt
tgttcaaaga ctttagtgtg attttcagga ccaagggctt tgagggtagt gctagctgt
gtaatagtaa tgaaggtgtg ggatcgtgtt tttgaattat gtaaaaaagg aagaaaaaa
aaatgttggt attatattat ggttttgcct gaaa
<210> 50 <211> 302 <212> PRT <213> Arabidopsis thaliana
<400> 50
Met Ala Asn Asn Asn Ile Pro His Asp Ser Ile Ser Asp Pro Ser 1 5 10 15
Pro Thr Asp Asp Phe Phe Glu Gln Ile Leu Gly Leu Ser Asn Phe Ser 20 25 30
Gly Ser Ser Gly Ser Gly Leu Ser Gly Ile Gly Gly Val Gly Pro Pro 35 40 45
Pro Met Met Leu Gln Leu Gly Ser Gly Asn Glu Gly Asn His Asn His 50 55 60
Met Gly Ala Ile Gly Gly Gly Pro Val Gly Phe His Asn Gln Met
65 70 75 80
Phe Pro Leu Gly Leu Ser Leu Asp Gln Gly Lys Gly His Gly Phe Leu 85 90 95
Phe Pro Leu Gly Leu Ser Leu Asp Gln Gly Lys Gly His Gly Phe Leu
Phe Pro Leu Gly Leu Ser Leu Asp Gln Gly Lys Gly His Gly Phe Leu 85 90 95 Lys Pro Asp Glu Thr Gly Lys Arg Phe Gln Asp Asp Val Leu Asp Asn

Pro Ala Pro Pro Met Pro His Gln Gln Ser Thr Ile Arg Pro Arg Val.
Page 66

MBI15 Sequence Listing.ST25 130 Arg Ala Arg Arg Gly Gln Ala Thr Asp Pro His Ser Ile Ala Glu Arg Leu Arg Arg Glu Arg Ile Ala Glu Arg Ile Arg Ser Leu Gln Glu Leu Val Pro Thr Val Asn Lys Thr Asp Arg Ala Ala Met Ile Asp Glu Ile 180 185 190 Val Asp Tyr Val Lys Phe Leu Arg Leu Gln Val Lys Val Leu Ser Met 195 200 205 Ser Arg Leu Gly Gly Ala Gly Ala Val Ala Pro Leu Val Thr Glu Met 210 215 220Pro Leu Ser Ser Val Glu Asp Glu Thr Gln Ala Val Trp Glu Lys Trp Ser Asn Asp Gly Thr Glu Arg Gln Val Ala Lys Leu Met Glu Glu 245 250 255 Asn Val Gly Ala Ala Met Gln Leu Leu Gln Ser Lys Ala Leu Cys Ile Met Pro Ile Ser Leu Ala Met Ala Ile Tyr His Ser Gln Pro Pro Asp 275 280 285 Thr Ser Ser Ser Ile Val Lys Pro Glu Met Asn Pro Pro 290 295 300 <210> 51 <211> 1198 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (96) . . (1052) <223> <400> 51 atcgaattcg cggccgctcg atatctttac aaccattaaa caaaaaattt ggccactaca 60 agttgaaaaa gttttgatta tatctaatcg ctgaa atg gat tac aag gta tca Met Asp Tyr Lys Val Ser 113 161 gat tta cca cct ggt ttc aga ttt cac cca act gat gaa gaa ctt ata Asp Leu Pro Pro Gly Phe Arg Phe His Pro Thr Asp Glu Glu Leu Ile 25 30 209

aca cac tat cta aga cca aag gtt gta aac tct ttt ttc tct gct ata Thr His Tyr Leu Arg Pro Lys Val Val Asn Ser Phe Phe Ser Ala Ile 40 45 50

gct att ggt gaa gtt gat ctc aac aaa gtc gag cct tgg gac ttg cct Ala Ile Gly Glu Val Asp Leu Asn Lys Val Glu Pro Trp Asp Leu Pro

Page 67

257

305

VD715 0 1''	
MBI15 Sequence Listing.ST25 55 60 65 70	
tgg aag gct aag ctt ggg gaa aaa gag tgg tac ttc ttt tgc gta aga Trp Lys Ala Lys Leu Gly Glu Lys Glu Trp Tyr Phe Phe Cys Val Arg 75 80 . 85	353
gac cga aaa tac ccg act ggt tta aga acg aat cgt gct act aaa gcc Asp Arg Lys Tyr Pro Thr Gly Leu Arg Thr Asn Arg Ala Thr Lys Ala 90 95 100	401
ggt tat tgg aaa gct aca ggg aaa gat aaa gag atc ttc aaa ggg aaa Gly Tyr Trp Lys Ala Thr Gly Lys Asp Lys Glu Ile Phe Lys Gly Lys 105 110 115	449
Ser Leu Val Gly Met Lys Lys Thr Leu Val Phe Tyr Lys Gly Arg Ala 120 125 130	497
cct aaa gga gta aaa aca aat tgg gtc atg cat gag tat cga tta gaa Pro Lys Gly Val Lys Thr Asn Trp Val Met His Glu Tyr Arg Leu Glu 135 140 145 150	545
ggc aaa ttc gct atc gat aat ctc tct aaa acc gct aag aac gaa tgt Gly Lys Phe Ala Ile Asp Asn Leu Ser Lys Thr Ala Lys Asn Glu Cys 155 160 165	593
gtt att agt cgt gtt ttt cat aca cgg act gat ggt acg aag gag cat Val Ile Ser Arg Val Phe His Thr Arg Thr Asp Gly Thr Lys Glu His 170 175 180	641
atg tcc gtt ggt tta cct ccg ctg atg gat tct tct cca tat cta aag Met Ser Val Gly Leu Pro Pro Leu Met Asp Ser Ser Pro Tyr Leu Lys 185 190 195	689
agt aga gga caa gac tct tta gcc ggg acc acc ctt ggt ggg ttg ttg Ser Arg Gly Gln Asp Ser Leu Ala Gly Thr Thr Leu Gly Gly Leu Leu 200 205 210	737
tct cac gtt acc tac ttc tcc gac caa aca acc gat gac aag agt ctt Ser His Val Thr Tyr Phe Ser Asp Gln Thr Thr Asp Asp Lys Ser Leu 215 220 225 230	785
gtg gcc gat ttt aaa act acc atg ttt ggt tcc gga tcg act aac ttt Val Ala Asp Phe Lys Thr Thr Met Phe Gly Ser Gly Ser Thr Asn Phe 235 240 245	833
tta cca aac ata ggt tct cta cta gac ttc gat cct ctg ttt cta caa Leu Pro Asn Ile Gly Ser Leu Leu Asp Phe Asp Pro Leu Phe Leu Gln 250 260	881
aac aat tct tca gta ctg aag atg ttg ctt gac aat gaa gaa acc caa Asn Asn Ser Ser Val Leu Lys Met Leu Leu Asp Asn Glu Glu Thr Gln 265 270 275	929
ttt aag aag aat ctt cac aat tca ggt tca tca gag agt gaa cta aca Phe Lys Lys Asn Leu His Asn Ser Gly Ser Ser Glu Ser Glu Leu Thr 280 285 290	977
gcg agt tct tgg caa ggt cac aat tct tat ggt tcc act ggt cca gtg Ala Ser Ser Trp Gln Gly His Asn Ser Tyr Gly Ser Thr Gly Pro Val 295 300 305 310	1025
aat ctt gat tgc gtt tgg aaa ttc tga atttggaaaa tcgaaaattt Asn Leu Asp Cys Val Trp Lys Phe 315	1072
ggatgttaac tagggggtat atagggtttt taaaaacagt gtatatatgc gttatgtgtt	1132
agotttagat totaggatat acaaagatga cactaataga ttottataac attttgtaaa	1192

<210> 52 <211> 318

aaaaaa

1198

MBI15 Sequence Listing.ST25

<212> PRT

<213> Arabidopsis thaliana

<400> 52

Met Asp Tyr Lys Val Ser Arg Ser Gly Glu Ile Val Glu Gly Glu Val 1 5 10 15

Glu Asp Ser Glu Lys Ile Asp Leu Pro Pro Gly Phe Arg Phe His Pro 25 30

Thr Asp Glu Glu Leu Ile Thr His Tyr Leu Arg Pro Lys Val Val Asn 35 40 45

Ser Phe Phe Ser Ala Ile Ala Ile Gly Glu Val Asp Leu Asn Lys Val 50 55 60

Glu Pro Trp Asp Leu Pro Trp Lys Ala Lys Leu Gly Glu Lys Glu Trp 65 70 75 80

Tyr Phe Phe Cys Val Arg Asp Arg Lys Tyr Pro Thr Gly Leu Arg Thr 85 90 95

Asn Arg Ala Thr Lys Ala Gly Tyr Trp Lys Ala Thr Gly Lys Asp Lys

Glu Ile Phe Lys Gly Lys Ser Leu Val Gly Met Lys Lys Thr Leu Val 115 120 125

Phe Tyr Lys Gly Arg Ala Pro Lys Gly Val Lys Thr Asn Trp Val Met 130 135 140

His Glu Tyr Arg Leu Glu Gly Lys Phe Ala Ile Asp Asn Leu Ser Lys 145 150 160

Thr Ala Lys Asn Glu Cys Val Ile Ser Arg Val Phe His Thr Arg Thr 165 170 175

Asp Gly Thr Lys Glu His Met Ser Val Gly Leu Pro Pro Leu Met Asp 180 185 190

Ser Ser Pro Tyr Leu Lys Ser Arg Gly Gln Asp Ser Leu Ala Gly Thr

Thr Leu Gly Gly Leu Leu Ser His Val Thr Tyr Phe Ser Asp Gln Thr 210 215 220

Thr Asp Asp Lys Ser Leu Val Ala Asp Phe Lys Thr Thr Met Phe Gly 225 230 235 240

Ser Gly Ser Thr Asn Phe Leu Pro Asn Ile Gly Ser Leu Leu Asp Phe 245 255

Asp Pro Leu Phe Leu Gln Asn Asn Ser Ser Val Leu Lys Met Leu Leu

Asp Asn Glu Glu Thr Gln Phe Lys Lys Asn Leu His Asn Ser Gly Ser 275 280 285

MBI15 Sequence Listing.ST25

Ser Glu Ser Glu Leu Thr Ala Ser Ser Trp Gln Gly His Asn Ser Tyr 290 295 300

Gly Ser Thr Gly Pro Val Asn Leu Asp Cys Val Trp Lys Phe 305 310 315

<210> 53 <211> 932 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (43)..(759) <223> G350 <400> 53 54 ctttcttctc aatttagaac ttagtagcta gtcttcaaga ta atg gca ctt gaa Met Ala Leu Glu act ctt act tct cca aga tta tct tct ccg atg ccg act ctg ttt caa Thr Leu Thr Ser Pro Arg Leu Ser Ser Pro Met Pro Thr Leu Phe Gln 102 gat toa goa cta ggg ttt cat gga agc aaa ggc aaa cga tot aag cga Asp Ser Ala Leu Gly Phe His Gly Ser Lys Gly Lys Arg Ser Lys Arg 150 tca aga tct gaa ttc gac cgt cag agt ctc acg gag gat gaa'tat atc Ser Arg Ser Glu Phe Asp Arg Gln Ser Leu Thr Glu Asp Glu Tyr Ile 198 gct tta tgt ctc atg ctt ctt gct cgc gac gga gat aga aac cgt gac Ala Leu Cys Leu Met Leu Leu Ala Arg Asp Gly Asp Arg Asn Arg Asp 55 60 65 ctt gac ctg cct tct tct tcg tct tca cct cct ctg ctt cct cct ctt Leu Asp Leu Pro Ser Ser Ser Ser Pro Pro Leu Leu Pro Pro Leu 70 75 80 294 cct act ccg atc tac aag tgt agc gtc tgt gac aag gcg ttt tcg tct Pro Thr Pro Ile Tyr Lys Cys Ser Val Cys Asp Lys Ala Phe Ser Ser 85 90 95 342 390 tac cag get ett ggt gga cac aag gea agt cac egg aaa age ttt teg Tyr Gln Ala Leu Gly Gly His Lys Ala Ser His Arg Lys Ser Phe Ser ctt act caa tct gcc gga gga gat gag ctg tcg aca tcg tcg gcg ata Leu Thr Gln Ser Ala Gly Gly Asp Glu Leu Ser Thr Ser Ser Ala Ile 438 acc acg tct ggt ata tcc ggt ggc ggg gga gga agt gtg aag tcg cac Thr Thr Ser Gly Ile Ser Gly Gly Gly Gly Gly Ser Val Lys Ser His 486 gtt tgc tct atc tgt cat aaa tcg ttc gcc acc ggt caa gct ctc ggc Val Cys Ser Ile Cys His Lys Ser Phe Ala Thr Gly Gln Ala Leu Gly 150 155 160 534 ggc cac aaa cgg tgc cac tac gaa gga aag aac gga ggc ggt gtg agt Gly His Lys Arg Cys His Tyr Glu Gly Lys Asn Gly Gly Gly Val Ser 582 agt agc gtg tcg aat tct gaa gat gtg ggg tct aca agc cac gtc agc Ser Ser Val Ser Asn Ser Glu Asp Val Gly Ser Thr Ser His Val Ser 630 678 agt ggc cac cgt ggg ttt gac ctc aac ata ccg ccg ata ccg gaa ttc

Ser Gly Hi	.s Arg 200	Gly	Phe	Asp		15 S Asn 205							Phe
tcg atg gt Ser Met Va 21	ıl Asn	gga Gly	gac Asp	gaa Glu	gag Glu 220	gtg Val	atg Met	agt Ser	cct Pro	atg Met 225	ccg Pro	gcg Ala	aag Lys
aaa ctc co Lys Leu Ar 230	g ttt g Phe	gac Asp	ttc Phe	ccg Pro 235	gag Glu	aaa Lys	ccc Pro	taa	acat	caaa	ect i	aggaa	aaact
ttacagaatt	catt	ttata	ıg ga	aatt	gttt	tac	ctgta	atat	aca	aatai	cg a	attt	gattg
atgttcttct	tcac	tgaaa	a at	tate	gatto	ttt	gtt	gtat	aati	gate	gtt 1	tctga	aaaag
atataacttt	ttat	taaaa	aa aa	aaaa	aaaa	a aaa	a						
<210> 54 <211> 238 <212> PRT <213> Ara		sis t	hali	lana									
<400> 54													
Met Ala Le 1	u Glu	Thr 5	Leu	Thr	Ser	Pro	Arg 10	Leu	Ser	Ser	Pro	Met 15	Pro
Thr Leu Ph	ie Gln 20	Asp	Ser	Ala	Leu	Gly 25	Phe	His	Gly	Ser	30 TAe	Gly	Lys
Arg Ser Ly		Ser	Arg	Ser	Glu 40	Phe	Asp	Arg	Gln	Ser 45	Leu	Thr	Glu
Asp Glu Ty 50	r Ile	Ala	Leu	Cys 55	Leu	Met	Leu	Leu	Ala 60	Arg	Asp	Gly	Asp
Arg Asn Ar 65	g Asp	Leu	Asp 70	Leu	Pro	Ser	Ser	Ser 75	Ser	Ser	Pro	Pro	Leu 80
Leu Pro Pr	o Leu	Pro 85	Thr	Pro	Ile	туг	Lys 90	Сув	Ser	Val	Сув	Asp 95	Lys
Ala Phe Se	er Ser 100	туг	Gln	Ala	Leu	Gly 105	Gly	His	Lys	Ala	Ser 110	His	Arg
Lys Ser Ph		Leu	Thr	Gln	Ser 120	Ala	Gly	Gly	Авр	Glu 125	Leu	Ser	Thr
Ser Ser Al	a Ile	Thr	Thr	Ser 135	Gly	Ile	Ser	Gly	Gly 140	Gly	Gly	Gly	Ser
Val Lys Se	er His	Val	Cys 150	Ser	Ile	Сув	His	Lys 155	Ser	Phe	Ala	Thr	Gly 160
Gln Ala Le	eu Gly	Gly 165	His	Lys	Arg	Cys	His 170	туг	Glu	Gly	Lys	Asn 175	Gly
Gly Gly Va	l Ser 180	Ser	Ser	Val	Ser	Asn 185	Ser	Glu	Asp	Val	Gly 190	Ser	Thr
Ser His Va		Ser	Gly	His	Arg 200	Gly		Asp age		Asn 205	Ile	Pro	Pro
							P	aut	/ 1				

MBI15 Sequence Listing.ST25

Ile Pro Glu Phe Ser Met Val Asn Gly Asp Glu Glu Val Met Ser Pro 210 215 220

Met Pro Ala Lys Lys Leu Arg Phe Asp Phe Pro Glu Lys Pro

<210> 55 <211> 1022 <212> DNA <213> Arabidopsis thaliana <220> <221> <222> (31) . . (846) G986 <223> <400> 55 54 cattaaattg gctcctgtga acctaaattt atg gac tat gat ccc aac acc aat Met Asp Tyr Asp Pro Asn Thr Asn ccg ttc gac ctt cat ttc tcc ggt aaa ctt ccg aaa aga gaa gtc tcg Pro Phe Asp Leu His Phe Ser Gly Lys Leu Pro Lys Arg Glu Val Ser 10 15 20 102 gct tca gct tct aaa gtt gta gag aag aaa tgg tta gtg aaa gat gag Ala Ser Ala Ser Lys Val Val Glu Lys Lys Trp Leu Val Lys Asp Glu 150 aag aga aat atg cta caa gat gaa ata aac cgg gtt aat tcg gag aac Lys Arg Asn Met Leu Gln Asp Glu Ile Asn Arg Val Asn Ser Glu Asn 198 aag aag cta acc gaa atg tta gca aga gtc tgt gag aag tac tat gct Lys Lys Leu Thr Glu Met Leu Ala Arg Val Cys Glu Lys Tyr Tyr Ala 246 294 ctt aat aat ctt atg gag gag ttg cag agt cga aag agt cct gaa agt Leu Asn Asn Leu Met Glu Glu Leu Gln Ser Arg Lys Ser Pro Glu Ser 80 gtt aac ttt cag aac aaa cag cta acg ggg aaa cga aaa caa gaa ctt Val Asn Phe Gln Asn Lys Gln Leu Thr Gly Lys Arg Lys Gln Glu Leu 90 95 100 342 gat gag ttt gtt agc tcc cca att gga ctc agt ctc gga cca atc gag Asp Glu Phe Val Ser Ser Pro Ile Gly Leu Ser Leu Gly Pro Ile Glu 390 aac atc acc aac gat aaa gcg acg gtt tca acc gct tac ttt gct gct Asn Ile Thr Asn Asp Lys Ala Thr Val Ser Thr Ala Tyr Phe Ala Ala 438 gag aag tot gac aca ago ttg act gtg aaa gat gga tat caa tgg agg Glu Lys Ser Asp Thr Ser Leu Thr Val Lys Asp Gly Tyr Gln Trp Arg aaa tac ggg caa aag att acg aga gat aat cca tct cct aga gct tac Lys Tyr Gly Gln Lys Ile Thr Arg Asp Asn Pro Ser Pro Arg Ala Tyr 155 160 165 534 ttc aga tgc tcg ttt tca ccg tct tgt cta gtc aag aag agg gtg caa Phe Arg Cys Ser Phe Ser Pro Ser Cys Leu Val Lys Lys Lys Val Gln 582 cga agt gca gaa gat cca tct ttc ttg gta gcc act tac gaa ggg aca Arg Ser Ala Glu Asp Pro Ser Phe Leu Val Ala Thr Tyr Glu Gly Thr 630 678 cat aac cac acc gga cca cat gca agt gtg tcc agg aca gtg aaa ctt

PCT/US00/31418 WO 01/35726

His	Asn	His	Thr	Gly 205	Pro	His	MBI Ala	15 S Ser	eque Val 210	nce Ser	List Arg	ing. Thr	ST25 Val	Lys 215	Leu	
														gag Glu		726
G1y 999	acg Thr	att Ile 235	caa Gln	gag Glu	gtt Val	ttg Leu	gtg Val 240	caa Gln	caa Gln	atg Met	gct Ala	tct Ser 245	tcg Ser	ttg Leu	acc Thr	774
														999 Gly		822
	ata Ile						tga	aagt	tct	cta q	gaaca	atgta	at a	ttci	gttt	876
tgt	ctat	tt t	gtt	gctca	at to	ctag	jtaaa	a aag	ggtaa	aaga	tttg	gttt	gat o	ettga	attagg	936
agge	ataç	gat q	gtcaa	attt	aat	gtgt	gtgt	ata	ataat	tac	atca	aaato	cta a	agtat	ccaaa	996
aagg	ggtca	acc o	ccat	ttta	at ct	tate	J									1022
<210 <210 <210 <210	l> 2 2> I	66 271 PRT Arabi	idops	sis t	hali	iana										
<400)> 9	6														
Met 1	Asp	Tyr	Asp	Pro 5	Asn	Thr	Asn	Pro	Phe 10	Asp	Leu	His	Phe	Ser 15	Gly	
Lys	Leu	Pro	Lys 20	Arg	Glu	Val	Ser	Ala 25	Ser	Ala	Ser	Lys	Val 30	Val	Glu	
Lys	Lys	Trp 35	Leu	Val	Lys	Asp	Glu 40	Lys	Arg	Asn	Met	Leu 45	Gln	Asp	Glu	
Ile	Asn 50	Arg	Val	Asn	Ser	Glu 55	Asn	Lys	Lys	Leu	Thr 60	Glu	Met	Leu	Ala	
Arg 65	Val	Суз	Glu	Lys	Tyr 70	Tyr	Ala	Leu	Asn	Asn 75	Leu	Met	Glu	Glu	Leu 80	
Gln	Ser	Arg	Lys	Ser 85	Pro	Glu	Ser	Val	Asn 90	Phe	Gln	Asn	Lys	Gln 95	Leu	
Thr	Gly	Lys	Arg 100	Lys	Gln	Glu	Leu	Asp 105	Glu	Phe	Val	Ser	Ser 110	Pro	Ile	
Gly	Leu	Ser 115	Leu	Gly	Pro	Ile	Glu 120	Asn	Ile	Thr	Asn	Asp 125	Lys	Ala	Thr	
Val	Ser 130	Thr	Ala	Tyr	Phe	Ala 135	Ala	Glu	Lys	Ser	Asp 140	Thr	Ser	Leu	Thr	
Val 145	Lys	Asp	Gly	Tyr	Gln 150	Trp	Arg	Lys	туг	Gly 155	Gln	Lys	Ile	Thr	Arg 160	
Asp	Asn	Pro	Ser	Pro 165	Arg	Ala	Tyr	Phe	170	Сув		Phe	Ser	Pro 175	Ser	

MBI15 Sequence Listing.ST25

Cys Leu Val Lys Lys Lys Val Gln Arg Ser Ala Glu Asp Pro Ser Phe Leu Val Ala Thr Tyr Glu Gly Thr His Asn His Thr Gly Pro His Ala Ser Val Ser Arg Thr Val Lys Leu Asp Leu Val Gln Gly Gly Leu Glu Pro Val Glu Glu Lys Lys Glu Arg Gly Thr Ile Gln Glu Val Leu Val Gln Gln Met Ala Ser Ser Leu Thr Lys Asp Pro Lys Phe Thr Ala Ala Leu Ala Thr Ala Ile Ser Gly Arg Leu Ile Glu His Ser Arg Thr <210> 57 <211> 1230 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (1)..(1089) <223> G1349 <400> 57 atg gct agt cgg aga gaa gta cgg tgc cgg tgc ggc aga cgg atg tgg Met Ala Ser Arg Arg Glu Val Arg Cys Arg Cys Gly Arg Arg Met Trp gtt caa cca gac gcc cgt acc gtc caa tgc tca acc tgc cac acc gtc Val Gln Pro Asp Ala Arg Thr Val Gln Cys Ser Thr Cys His Thr Val 96 acg cag ctc tac tcg cta gtg gac ata gct cgc ggt gca aac cgc ata Thr Gln Leu Tyr Ser Leu Val Asp Ile Ala Arg Gly Ala Asn Arg Ile 35 40 45144 192 att cat ggg ttt caa cag cta ctt aga caa cac caa ccg caa cat cat Ile His Gly Phe Gln Gln Leu Leu Arg Gln His Gln Pro Gln His His gaa caa caa caa caa caa atg atg gct caa ccg cca cca cgg ctg ctt Glu Gln Gln Gln Gln Met Met Ala Gln Pro Pro Pro Arg Leu Leu 240 gag cct ctt ccc tcg ccg ttt ggg aag aag aga gca gtt tta tgc ggc Glu Pro Leu Pro Ser Pro Phe Gly Lys Lys Arg Ala Val Leu Cys Gly 288 gtg aac tat aag gga aaa agt tat agc ttg aaa ggt tgc atc agt gat Val Asn Tyr Lys Gly Lys Ser Tyr Ser Leu Lys Gly Cys Ile Ser Asp 100 110 336 gct aag tcc atg aga tct tta ttg gtt caa caa atg ggt ttc cct att Ala Lys Ser Met Arg Ser Leu Leu Val Gln Gln Met Gly Phe Pro Ile 384 gac tct att ctc atg ctc aca gaa gat gaa gcc agc ccg cag aga ata Asp Ser Ile Leu Met Leu Thr Glu Asp Glu Ala Ser Pro Gln Arg Ile 432 135 480 ccg acg aag aga aac att agg aag gcg atg aga tgg tta gtt gaa ggg Page 74

Pro 145	Thr	Lys	Arg	Asn	Ile	Arg					List Trp			Glu	Gly 160	
aac	aga Arg	gca Ala	agg Arg	gac Asp 165	tca Ser	cta Leu	gtg Val	ttc Phe	cat His 170	ttc	tct Ser	ggt Gly	cat His	gga Gly 175	tct	528
					aac Asn											576
					cat His											624
					gtg Val											672
					tgt Cys 230											720
att Ile	tgc Cys	agg Arg	atg Met	gag Glu 245	agg Arg	aat Asn	ggt Gly	tct Ser	tat Tyr 250	gaa Glu	tgg Trp	gaa Glu	gac Asp	cat His 255	aga Arg	768
tca Ser	gtc Val	aga Arg	gct Ala 260	tac Tyr	aaa Lys	gga Gly	aca Thr	gat Asp 265	ggt Gly	gga Gly	gca Ala	gct Ala	ttc Phe 270	tgt Cys	ttc Phe	. 816
					gat Asp											864
					gcc Ala											912
					ccc Pro 310											960
tct Ser	gca Ala	ata Ile	cga Arg	gag Glu 325	gcc Ala	cag Gln	tct Ser	cgc Arg	ctc Leu 330	gcc Ala	ttt Phe	aac Asn	999 Gly	gac Asp 335	tac Tyr	1008
					tcc Ser											1056
ttt Phe	gac Asp	gtg Val 355	tac Tyr	gcg Ala	aca Thr	aag Lys	ttt Phe 360	gta Val	ctc Leu	tga	atgo	etgta	aca (atga	atgctg	1109
caaa	tago	tc g	ggaaa	cgtt	t ct	atgt	gtat	gta	tcat	gta	atga	attat	gt I	gcat	agcct	1169
ctct	ctto	ctt a	acgaç	gcaat	a aç	gctat	gaaa	a taa	ttga	ttc	gcta	agaa	aat 1	taaa	aatgaa	1229
a																1230
<210 <211 <212 <213	> 3 > E	8 862 PRT Arabi	ldops	sis t	:hali	ana										

Met Ala Ser Arg Arg Glu Val Arg Cys Arg Cys Gly Arg Arg Met Trp 1 5 10 15

Val Gln Pro Asp Ala Arg Thr Val Gln Cys Ser Thr Cys His Thr Val 20 25 30

MBI15 Sequence Listing.ST25

Thr Gln Leu Tyr Ser Leu Val Asp Ile Ala Arg Gly Ala Asn Arg Ile 35 40 45

Ile His Gly Phe Gln Gln Leu Leu Arg Gln His Gln Pro Gln His His 50 55 60

Glu Gln Gln Gln Gln Met Met Ala Gln Pro Pro Pro Arg Leu Leu 65 70 75 80

Glu Pro Leu Pro Ser Pro Phe Gly Lys Lys Arg Ala Val Leu Cys Gly 85 90 95

Val Asn Tyr Lys Gly Lys Ser Tyr Ser Leu Lys Gly Cys Ile Ser Asp 100 105 110

Ala Lys Ser Met Arg Ser Leu Leu Val Gln Gln Met Gly Phe Pro Ile 115 120 125

Asp Ser Ile Leu Met Leu Thr Glu Asp Glu Ala Ser Pro Gln Arg Ile 130 135 140

Pro Thr Lys Arg Asn Ile Arg Lys Ala Met Arg Trp Leu Val Glu Gly 145 150 150 160

Asn Arg Ala Arg Asp Ser Leu Val Phe His Phe Ser Gly His Gly Ser 165 170 175

Gln Gln Asn Asp Tyr Asn Gly Asp Glu Ile Asp Gly Gln Asp Glu Ala

Leu Cys Pro Leu Asp His Glu Thr Glu Gly Lys Ile Ile Asp Asp Glu 195 200 205

Ile Asn Arg Ile Leu Val Arg Pro Leu Val His Gly Ala Lys Leu His 210 215 220

Ala Val Ile Asp Ala Cys Asn Ser Gly Thr Val Leu Asp Leu Pro Phe 225 230 230 235

Ile Cys Arg Met Glu Arg Asn Gly Ser Tyr Glu Trp Glu Asp His Arg 245 250 255

Ser Ala Cys Asp Asp Glu Ser Ser Gly Tyr Thr Pro Val Phe Thr 275 280 285

Gly Lys Asn Thr Gly Ala Met Thr Tyr Ser Phe Ile Lys Ala Val Lys 290 295 300

Thr Ala Gly Pro Ala Pro Thr Tyr Gly His Leu Leu Asn Leu Met Cys 305 310 315

Ser Ala Ile Arg Glu Ala Gln Ser Arg Leu Ala Phe Asn Gly Asp Tyr Page 76

MBI15 Sequence Listing.ST25

Phe Asp Val Tyr Ala Thr Lys Phe Val Leu 355

325

al application No. Interna INTERNATIONAL SEARCH REPORT PCT/US00/31418 CLASSIFICATION OF SUBJECT MATTER IPC(7) : A01H 1/00, 5/00; C12N 5/14, 15/82 435/320.1, 419, 468; 800/278, 279, 287, 301, 305-310, 312, 314, 317, 320, 322 US CL According to International Patent Classification (IPC) or to both national classification and IPC FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S.: 435/320.1, 419, 468; 800/278, 279, 287, 301, 305-310, 312, 314, 317, 320, 322 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EAST, USPAT, STN, Agricola, CaPlus, Biosis, Embase DOCUMENTS CONSIDERED TO BE RELEVANT Relevant to claim No. Citation of document, with indication, where appropriate, of the relevant passages Category * WO 97/47183 A1 (PURDUE RESEARCH FOUNDATION) 18 December 1997 1-9, 12, 13, 25 (18.12.1997), entire reference. 10, 11, 26, 27 Υ 1-9, 12, 13, 25 US 5,939,601 (KLESSIG et al) 17 August 1999 (17.08.1999), entire reference. Х 10, 11, 26, 27 Y Database Genbank on NCBI, US National Library of Medicine, (Bethesda, MD, USA) 1-13, 25-27 A No. AB009055, SATO, S. et al 'Strucural analysis of Arabidopsis thaliana chromosome 5. IV. Sequence features of the regions of 1,456,315 bp covered by nineteen physically assigned P1 and TAC clones. 27 December 2000, DNA RES. 1998, Vol. 5, No. 1, pages 41-54, see bases 16,003-16,490, 16,571-16,683 and 16,780-17,365. Further documents are listed in the continuation of Box C. See patent family annex. later document published after the international filing date or priority Special categories of cited documents: date and not in conflict with the application but cited to understand the principle or theory underlying the invention document defining the general state of the art which is not considered to be "A" "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "E" earlier application or patent published on or after the international filing date when the document is taken alone document which may throw doubts on priority claim(s) or which is cited to -Y" document of particular relevance; the claimed invention cannot be establish the publication date of another citation or other special reason (as considered to involve an inventive step when the document is specified) combined with one or more other such documents, such combination being obvious to a person skilled in the art "O" document referring to an oral disclosure, use, exhibition or other means document member of the same patent family document published prior to the international filing date but later than the **"&"** priority date claimed Date of the actual completion of the international search Date of mailing of the international search report

Authorized officer

PARALEGAL SPECIALIST

Telephone No. 703-308-FEDIMOLOGY CENTER 1600

David Kruse

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Washington, D.C. 20231

23 February 2001 (23.02.2001)

Box PCT

Facsimile No. (703)305-3230

INTERNATIONAL SEARCH REPORT

Inter nal application No.
PCT/US00/31418

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)		
This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:		
1.		Claim Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
2.		Claim Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3.	6.4(a).	Claim Nos.: 14 because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule
Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)		
		ional Searching Authority found multiple inventions in this international application, as follows: ontinuation Sheet
1. 2. 3.		As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. Rema	ark on l	No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-13 & 25-27 and SEQ ID NOs 1&2 Protest

INTERNATIONAL SEARCH REPORT

Ì

Inte.....nal application No.

PCT/US00/31418

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Groups I-XXIX, claim(s) 1-14 and 25-27, drawn to a transgenic plant having modified seed characteristics, polynucleotides and vectors for producing said transgenic plant and a method of making said transgenic plant. Applicant must elect one pair of sequences (one nucleic acid and the corresponding amino acid translation) to be examined, i.e. SEQ ID NO: 1 and 2 in Group I, SEQ ID NO: 3 and 4 in Group II, SEQ ID NO: 5 and 6 in Group III, etc.

Group XXX, claim(s) 15-17, drawn to a method of identifying a factor that is modulated.

Group XXXI, claims(s) 18, drawn to a method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide.

Group XXXII, claims(s) 19 and 20, drawn to an integrated computer system.

Group XXXIII, claim(s) 21-24, drawn to a method for identifying a polynucleotide sequence comprising selecting a nucleic acid sequence from a database that meets a selected sequence criteria.

The inventions listed as Groups I-XXXIII do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The inventions listed as Groups I-XXXIII do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Groups I-XXIX are drawn to a transgenic plant and a method of producing said plant with a nucleic acid sequence. The methods of Groups I-XXIX differ from each other in that they are directed to a plant transformation method and transgenic plant with a structurally and functionally distinct nucleic acid sequence which encodes a structurally and functionally distinct amino acid sequence. In addition, Groups XXX, XXXI and XXXIII are different methods from any of Groups I-XXIX in that they have different method steps and different end products, and Group XXXII requires a computer system. Thus, there is no single special technical feature, which links the inventions of Groups I-XXXIII under PCT Rule 13.2.